



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2021 – 09:25 am BST

PDB ID : 7PUC
Title : CARM1 in complex with EML981
Deposited on : 2021-09-29
Resolution : 2.19 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	92% 5% 5%
1	B	361	90% 5% 5% 2%
1	C	361	91% 5% 5% 4%
1	D	361	88% 7% 5% 3%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	502	-	-	-	X

CONFIDENTIAL VALIDATION REPORT

2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 22350 atoms, of which 10942 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	343	5450	1776	2699	454	507	14	0	0	0
1	B	343	5460	1778	2705	457	506	14	0	0	0
1	C	342	5436	1772	2692	453	505	14	0	0	0
1	D	342	5439	1769	2696	456	504	14	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY	-	expression tag	UNP Q9WVG6
A	128	HIS	-	expression tag	UNP Q9WVG6
A	129	MET	-	expression tag	UNP Q9WVG6
B	127	GLY	-	expression tag	UNP Q9WVG6
B	128	HIS	-	expression tag	UNP Q9WVG6
B	129	MET	-	expression tag	UNP Q9WVG6
C	127	GLY	-	expression tag	UNP Q9WVG6
C	128	HIS	-	expression tag	UNP Q9WVG6
C	129	MET	-	expression tag	UNP Q9WVG6
D	127	GLY	-	expression tag	UNP Q9WVG6
D	128	HIS	-	expression tag	UNP Q9WVG6
D	129	MET	-	expression tag	UNP Q9WVG6

- Molecule 2 is methyl 6-[4-[[{N}-[({E})-3-[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]prop-2-enyl]carbamidoyl]amino]butylcarbomoylamino]-4-oxidanyl-naphthalene-2-carboxylate (three-letter code: 85U) (formula: C₃₀H₃₆N₁₀O₇) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	H	N			O
2	A	1	Total	C	H	N	O	0	0
			83	30	36	10	7		
2	B	1	Total	C	H	N	O	0	0
			83	30	36	10	7		
2	C	1	Total	C	H	N	O	0	0
			83	30	36	10	7		
2	D	1	Total	C	H	N	O	0	0
			83	30	36	10	7		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	10	2	6	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	70	Total	O	0	0
			70	70		
4	B	63	Total	O	0	0
			63	63		
4	C	50	Total	O	0	0
			50	50		
4	D	40	Total	O	0	0
			40	40		

CONFIDENTIAL


VALIDATION

REPORT

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-arginine methyltransferase CARM1

Chain A: 





- Molecule 1: Histone-arginine methyltransferase CARM1

Chain B: 





- Molecule 1: Histone-arginine methyltransferase CARM1

Chain C: 



- Molecule 1: Histone-arginine methyltransferase CARM1

Chain D: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	75.32Å 98.94Å 208.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.19 48.13 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.0 (48.13-2.19) 99.2 (48.13-2.19)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.18Å)	Xtriage
Refinement program	PHENIX 1.19.2, 4158	Depositor
R, R_{free}	0.202 , 0.231 0.200 , 0.227	Depositor DCC
R_{free} test set	3991 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available), (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	22350	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.1217e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 85U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2821	0.62	0/3823
1	B	0.43	0/2825	0.64	0/3827
1	C	0.43	0/2814	0.64	0/3813
1	D	0.42	0/2812	0.62	0/3809
All	All	0.43	0/11272	0.63	0/15272

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2751	2699	2699	4	0
1	B	2755	2705	2705	18	0
1	C	2744	2692	2692	8	0
1	D	2743	2696	2696	23	0
2	A	47	36	0	0	0
2	B	47	36	0	0	0
2	C	47	36	0	1	0
2	D	47	36	0	0	0
3	A	4	6	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	70	0	0	0	0
4	B	63	0	0	0	0
4	C	50	0	0	0	0
4	D	40	0	0	0	0
All	All	11408	10942	10798	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:230:ASN:O	1:D:232:LEU:HD23	1.77	0.85
1:D:182:THR:OG1	1:D:400:ILE:HD11	1.82	0.78
1:D:428:PHE:CZ	1:D:430:LYS:HD3	2.27	0.68
1:A:270:LEU:HD12	1:A:360:PHE:CE1	2.32	0.65
1:D:182:THR:OG1	1:D:400:ILE:CD1	2.50	0.58
1:A:188:ILE:HD11	1:A:209:ARG:HH21	1.69	0.57
1:D:441:LEU:N	1:D:441:LEU:HD12	2.20	0.56
1:D:428:PHE:CZ	1:D:430:LYS:CD	2.88	0.56
1:A:430:LYS:HG2	1:A:433:ASP:OD2	2.06	0.56
1:B:312:ASN:OD1	1:B:315:TYR:CE1	2.60	0.55
1:B:143:GLU:HG3	1:B:445:LYS:HB3	1.89	0.54
1:B:315:TYR:CE1	1:B:328:ARG:HG3	2.43	0.53
1:C:282:PRO:O	1:C:283:SER:OG	2.23	0.52
1:B:315:TYR:CD1	1:B:328:ARG:HG3	2.45	0.52
1:B:318:SER:HB3	1:D:463:LYS:H	1.75	0.52
1:D:188:ILE:HD13	1:D:250:GLU:CD	2.30	0.51
1:C:263:MET:CE	1:C:419:VAL:HG11	2.40	0.51
1:D:428:PHE:HZ	1:D:430:LYS:HD3	1.72	0.51
1:D:230:ASN:O	1:D:232:LEU:CD2	2.52	0.51
1:B:312:ASN:OD1	1:B:315:TYR:HE1	1.95	0.50
1:B:263:MET:CE	1:B:419:VAL:HG11	2.42	0.49
1:D:428:PHE:CE2	1:D:430:LYS:HD3	2.48	0.48
1:B:318:SER:CB	1:D:463:LYS:H	2.26	0.48
1:B:143:GLU:HG3	1:B:445:LYS:CB	2.45	0.47
1:B:430:LYS:HG3	1:C:167:TYR:CE2	2.50	0.46
1:D:337:ARG:HG2	1:D:467:LEU:O	2.15	0.46
1:B:293:VAL:CG1	1:B:356:TYR:HB3	2.46	0.45
1:D:263:MET:CE	1:D:419:VAL:HG11	2.47	0.45
1:D:188:ILE:HD13	1:D:250:GLU:OE1	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:314:TRP:O	1:D:325:SER:HA	2.17	0.45
1:C:263:MET:HE3	1:C:419:VAL:HG11	1.99	0.44
1:C:293:VAL:CG1	1:C:356:TYR:HB3	2.48	0.44
1:D:350:MET:HE2	1:D:350:MET:HA	2.00	0.43
1:B:312:ASN:HA	1:B:315:TYR:HD1	1.83	0.43
1:D:295:LEU:HD21	1:D:373:ILE:HG21	2.01	0.43
1:B:463:LYS:H	1:D:318:SER:HB3	1.84	0.42
1:B:314:TRP:O	1:B:325:SER:HA	2.18	0.42
1:D:439:CYS:SG	1:D:441:LEU:HD11	2.59	0.42
1:C:384:LEU:HD12	1:C:384:LEU:N	2.34	0.42
1:B:370:ARG:HG3	1:B:442:ILE:CD1	2.50	0.42
1:A:453:ILE:O	1:A:465:SER:HA	2.20	0.41
1:C:262:TYR:OH	2:C:501:85U:O4	2.33	0.41
1:C:453:ILE:O	1:C:465:SER:HA	2.20	0.41
1:D:441:LEU:N	1:D:441:LEU:CD1	2.84	0.41
1:D:453:ILE:O	1:D:465:SER:HA	2.21	0.41
1:B:293:VAL:HG12	1:B:356:TYR:HB3	2.03	0.40
1:B:315:TYR:CE1	1:B:328:ARG:CG	3.04	0.40
1:B:463:LYS:H	1:D:318:SER:CB	2.34	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/361 (94%)	333 (98%)	8 (2%)	0	100	100
1	B	341/361 (94%)	332 (97%)	9 (3%)	0	100	100
1	C	340/361 (94%)	333 (98%)	7 (2%)	0	100	100
1	D	340/361 (94%)	331 (97%)	9 (3%)	0	100	100
All	All	1362/1444 (94%)	1329 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/313 (96%)	297 (99%)	2 (1%)	84	91
1	B	299/313 (96%)	297 (99%)	2 (1%)	84	91
1	C	298/313 (95%)	296 (99%)	2 (1%)	84	91
1	D	298/313 (95%)	294 (99%)	4 (1%)	69	81
All	All	1194/1252 (95%)	1184 (99%)	10 (1%)	81	90

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	GLU
1	A	425	SER
1	B	169	ARG
1	B	328	ARG
1	C	169	ARG
1	C	185	LYS
1	D	144	GLU
1	D	169	ARG
1	D	242	LYS
1	D	295	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	GLN
1	B	149	GLN
1	B	160	GLN
1	C	160	GLN
1	D	160	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	85U	C	501	-	46,51,51	0.46	0	59,71,71	0.63	1 (1%)
3	EDO	A	502	-	3,3,3	0.09	0	2,2,2	0.28	0
2	85U	B	501	-	46,51,51	0.45	0	59,71,71	0.64	2 (3%)
2	85U	D	501	-	46,51,51	0.44	0	59,71,71	0.55	1 (1%)
2	85U	A	501	-	46,51,51	0.47	0	59,71,71	0.57	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	85U	C	501	-	-	8/26/46/46	0/5/5/5
3	EDO	A	502	-	-	1/1/1/1	-
2	85U	B	501	-	-	9/26/46/46	0/5/5/5
2	85U	D	501	-	-	6/26/46/46	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	85U	A	501	-	-	10/26/46/46	0/5/5/5

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	85U	C24-C25-N8	2.35	123.92	120.35
2	A	501	85U	C24-C25-N8	2.31	123.86	120.35
2	C	501	85U	C24-C25-N8	2.31	123.86	120.35
2	D	501	85U	C24-C25-N8	2.30	123.85	120.35
2	B	501	85U	C14-N2-C13	2.00	125.75	121.75

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	85U	C1-C11-O2-C12
2	B	501	85U	C15-C14-N2-C13
2	B	501	85U	N4-C17-N3-C28
2	B	501	85U	N5-C17-N3-C28
2	C	501	85U	C1-C11-O2-C12
2	C	501	85U	N2-C13-N1-C8
2	C	501	85U	O4-C13-N1-C8
2	C	501	85U	N4-C17-N3-C28
2	C	501	85U	N5-C17-N3-C28
2	D	501	85U	C1-C11-O2-C12
2	B	501	85U	C1-C11-O2-C12
2	D	501	85U	O3-C11-O2-C12
2	B	501	85U	O3-C11-O2-C12
2	C	501	85U	O3-C11-O2-C12
2	A	501	85U	O3-C11-O2-C12
2	D	501	85U	N2-C13-N1-C8
2	D	501	85U	O4-C13-N1-C8
2	B	501	85U	N2-C13-N1-C8
3	A	502	EDO	O1-C1-C2-O2
2	B	501	85U	O4-C13-N1-C8
2	B	501	85U	C14-C15-C16-C28
2	A	501	85U	C30-C18-C19-O7
2	D	501	85U	C15-C16-C28-N3
2	A	501	85U	N2-C13-N1-C8
2	D	501	85U	N2-C14-C15-C16

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	501	85U	N5-C17-N3-C28
2	A	501	85U	O4-C13-N1-C8
2	A	501	85U	N4-C17-N3-C28
2	A	501	85U	N2-C14-C15-C16
2	A	501	85U	C30-C18-C19-C20
2	A	501	85U	N4-C29-C30-C18
2	C	501	85U	C14-C15-C16-C28
2	B	501	85U	C30-C29-N4-C17
2	C	501	85U	C15-C14-N2-C13


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	85U	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

CONFIDENTIAL VALIDATION REPORT

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	343/361 (95%)	0.19	3 (0%) 84 83	29, 44, 61, 78	0
1	B	343/361 (95%)	0.24	8 (2%) 60 58	34, 49, 63, 97	0
1	C	342/361 (94%)	0.35	16 (4%) 31 30	37, 54, 73, 81	0
1	D	342/361 (94%)	0.28	10 (2%) 51 49	40, 54, 68, 95	0
All	All	1370/1444 (94%)	0.27	37 (2%) 54 52	29, 51, 68, 97	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	347	ARG	4.4
1	B	477	TYR	3.5
1	D	346	ILE	3.4
1	A	283	SER	3.3
1	C	276	ALA	3.2
1	C	178	LEU	3.2
1	D	350	MET	3.1
1	C	136	SER	3.1
1	B	347	ARG	3.0
1	D	380	LEU	3.0
1	C	432	GLY	3.0
1	D	315	TYR	2.9
1	C	204	ALA	2.8
1	B	135	ARG	2.7
1	C	234	ASP	2.7
1	C	477	TYR	2.7
1	C	284	GLY	2.6
1	C	184	PHE	2.6
1	B	364	LYS	2.6
1	D	135	ARG	2.6
1	D	248	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	428	PHE	2.5
1	C	212	TYR	2.5
1	B	188	ILE	2.5
1	D	167	TYR	2.4
1	D	378	HIS	2.4
1	C	249	PRO	2.3
1	C	183	ASP	2.3
1	C	210	LYS	2.3
1	C	203	ALA	2.2
1	B	380	LEU	2.2
1	B	475	PHE	2.1
1	A	477	TYR	2.1
1	B	315	TYR	2.1
1	D	322	VAL	2.0
1	C	200	SER	2.0
1	C	190	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
3	EDO	A	502	4/4	0.78	0.47	76,92,96,100	0
2	85U	C	501	47/47	0.86	0.24	56,86,156,179	0
2	85U	D	501	47/47	0.90	0.21	53,86,118,138	0
2	85U	A	501	47/47	0.90	0.20	45,79,136,159	0
2	85U	B	501	47/47	0.92	0.24	49,83,131,148	0

The following is a graphical depiction of the model fit to experimental electron density of all




instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






CONFIDENTIAL


Electron density around 85U D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 85U A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.