

Supplemental Online Content

Gahm J, Ljung Konstantinidou A, Lagergren J, et al. Effectiveness of single vs multiple doses of prophylactic intravenous antibiotics in implant-based breast reconstruction: a randomized clinical trial. *JAMA Netw Open*. 2022;5(9):e2231583.
doi:10.1001/jamanetworkopen.2022.31583

eTable. Outcome at 6-Month Follow-up for Per-Protocol Analysis, According to Randomization (n = 647)

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable. Outcome at 6-Month Follow-up for Per-Protocol Analysis, According to Randomization (n = 647)

| Outcome | Single-dose (%) N=321 | Multiple-dose (%) N=326 | Odds-ratio ^a (95% confidence interval) | p-value |
|--------------------------------------|--------------------------|----------------------------|--|---------|
| Implant removal ^b | 12 (3.7) | 16 (4.9) | 1.31 (0.61-2.82) | 0.487 |
| Intravenous antibiotics ^c | 18 (5.6) | 24 (7.4) | 1.30 (0.69-2.45) | 0.413 |
| Oral antibiotics ^d | 97 (30.2) | 77 (23.6) | 0.71 (0.50-1.02) | 0.063 |
| | | | | |

^aMultiple-dose versus single-dose.

^b12 patients (1.9%) with missing outcome data (8/4).

^c18 patients (2.8%) with missing outcome data (13/5).

^d18 patients (2.8%) with missing outcome data (8/10).