
Supplemental Online Content



eTable. ICD-9-CM and ICD-10-CM diagnostic codes for identifying suicide attempt, intentional self-harm, and other diagnoses included in the statistical models

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable. ICD-9-CM and ICD-10-CM diagnostic codes for identifying suicide attempt, intentional self-harm, and other diagnoses included in the statistical models

Diagnosis	ICD-9-CM	ICD-10-CM
Intentional self-harm due to:		
Non-Fatal Suicide Attempts and Intentional Self-Harm	E950-E959	T36–T50 with the 6th character of the code = 2 (except for T36.9, T37.9, T39.9, T41.4, T42.7, T43.9, T45.9, T47.9, and T49.9, which are included if the 5th character of the code = 2)
		T51–T65 with the 6th character of the code = 2 (except for T51.9, T52.9, T53.9, T54.9, T56.9, T57.9, T58.0, T58.1, T58.9, T59.9, T60.9, T61.0, T61.1, T61.9, T62.9, T63.9, T64.0, T64.8, and T65.9, which are included if the 5th character of the code = 2)
		T71 with the 6th character of the code = 2
		Т14.91
		X71-X83
Other included diagnoses		
Substance use disorder	291, 292, 303, 304, 305 (except 305.1)	F10-F19 (except F17)
Depressive disorder	296.2, 296.3, 300.4, 311	F32, F33, F34.1
Anxiety/stress/adjustment disorder	300.0, 300.2, 300.3, 308, 309.0, 309.24- 309.9	F40-F43
Attention-deficit/hyperactivity disorder	314	F90
Bipolar/other mood disorder	296 (except 296.2, 296.3)	F30-F39 (except F32, F33, F34.1)
Schizophrenia	295, 297, 298	F20-F29
Sleep disorder	307.4, 327, 347, 780.5	F51.01, F51.03, F51.09, F51.11, F51.12, F51.19, F51.3, F51.8, G47
Pain	339,346,7840,7280, 72881, 7291, 354- 357, 830-854, 860-887,890-897, 900-959	G43, G44, R51, M60, M791, M797, G56-G59, G629, M792, S00-S78 excluding Sx2, S798, S799, S80, S81, S83-S88, S898, S899, S90, S91, S93-S99, T07, T14-T28, T30-T34, T79
Crohn's disease	5559	K5090
Celiac disease	5790	K900