

Impact of the COVID-19 pandemic on the care and outcomes of patients with NAFLD-related cirrhosis

Jesús Rivera-Esteban, Ramiro Manzano-Nuñez, Teresa Broquetas, Isabel Serra-Matamala, Octavi Bassegoda, Agnès Soriano-Varela, Gemma Espín, Joaquín Castillo, Juan Bañares, José Carrión, Pere Ginès, Isabel Graupera,
Juan M Pericàs

Table of contents

Table S1.....	2
---------------	---

Table S1. Baseline characteristics of the cohort according to Child-Pugh classification.

	Overall n=354	Child A n=287	Child B-C n=43	p value
Age, mean years (SD)	67.3 (9.6)	66.9 (9.2)	68.8 (10.5)	0.11
Females, n (%)	171 (48.3)	138 (48.1)	19 (44.2)	0.63
Tobacco use, n (%)	38 (10.8)	32 (11.1)	3 (7.0)	0.21
Alcohol use, n (%)*	76 (21.5)	62 (21.6)	11 (25.6)	0.57
Body Mass Index, median kg/m² (IQR)	31.2 (27.6-35.1)	31.8 (27.8-35.3)	30.2 (26.9-32.9)	0.023
BMI ≥ 25 kg/m², n (%)	302 (92.9)	249 (92.9)	35 (94.6)	0.70
BMI ≥ 30 kg/m², n (%)	188 (57.8)	152 (56.7)	23 (62.2)	0.53
Arterial hypertension, n (%)	251 (70.9)	206 (71.8)	29 (67.4)	0.55
T2D, n (%)	258 (72.9)	214 (74.6)	28 (65.1)	0.19
Dyslipidaemia, n (%)	181 (51.1)	156 (54.4)	15 (34.9)	0.017
Previous stroke, n (%)	16 (4.5)	12 (4.2)	3 (7.0)	0.41
Previous ischemic heart disease, n (%)	35 (9.9)	31 (10.8)	3 (7.0)	0.44
Liver stiffness, mean kPa (SD)**	23.6 (14.8)	22.9 (13.8)	33.9 (25.4)	0.10
CAP, mean dB/m (SD)***	307.0 (58.0)	308.1 (57.6)	281.3 (77.6)	0.43

*Alcohol intake was defined as <20 gr/day and <30gr/day for women and men, respectively.

*Data available in 83 subjects.

**Data available in 75 subjects.