

Supplementary Information

Clinical Pharmacokinetics

Single-dose Pharmacokinetics of Milvexian in Participants with Normal Renal Function and Participants with Moderate or Severe Renal Impairment

Vidya Perera,¹ Grigor Abelian, Danshi Li, Zhaoqing Wang, Liping Zhang, Susan Lubin, Akintunde Bello, Bindu Murthy

¹ **Corresponding author:** Bristol Myers Squibb, 3401 Princeton Pike, Lawrenceville, NJ 08648, USA. E-mail: Vidya.Perera@bms.com

Online Resource 1. Pharmacokinetic (PK) sampling schedule

Study day	Time (relative to milvexian dose), hour:min	PK blood sample for plasma conc.	PK protein binding blood sample	PK urine sample	PD blood sample
1	00:00	X ^a		X ^a	X ^a
1	00:30	X		X (0-8 h)	X
1	1:00	X			
1	1:30	X			
1	2:00	X			
1	3:00	X			X
1	4:00	X	X		X
1	4:30	X			
1	5:00	X			X
1	6:00	X			X
1	7:00	X			X
1	8:00	X		X (8-12 h)	X
1	10:00	X			
1	12:00	X		X (12-24 h)	X
2	18:00	X			
2	24:00	X		X (24-48 h)	X
2	36:00	X			
3	40:00	X			
3	48:00	X		X (48-72 h)	X
4	72:00	X		X (72-96 h)	X
5	96:00	X			

^a Sample obtained prior to milvexian administration.
Conc. concentration, *PD* pharmacodynamics,