Supplemental Table 1. Key Inclusion and Exclusion Criteria.

	Inclusion criteria
٠	Male or female aged 18 years or older
•	Type 1 or type 2 Diabetes Mellitus
•	CI-DME with CST of \geq 320µm on Spectralis [®] SD-OCT
•	BCVA \leq 62 and \geq 23 ETDRS letter score in the study eye
•	History of response to anti-VEGF agents and/or corticosteroids and, in the opinion of the Investigator
	remained responsive to treatment
•	Written informed consent obtained from the subject prior to screening procedures
	Exclusion criteria
٠	Macular edema due to causes other than DME
•	Concurrent disease in the study eye, other than CI-DME, that could compromise BCVA, require
	medical or surgical intervention during the study period or could confound interpretation of the resul
•	Any condition that could confound the ability to detect a change in CST in the study eye
•	Previous confounding treatments / procedures, or expected to require confounding treatments /
	procedures at any time during the study period
•	Presence of neovascularization at the disc in the study eye
٠	Uncontrolled glaucoma in the study eye
•	Any active ocular / intra-ocular infection or inflammation in either eye
•	Any active ocular / intra-ocular infection or inflammation in either eye Poorly controlled Diabetes Mellitus

BCVA = best-corrected visual acuity; CI-DME = center-involved diabetic macular edema; CST = central subfield thickness; DME = diabetic macular edema; ETDRS = international clinical diabetic retinopathy disease severity scale; SD-OCT = spectral domain optical coherence tomography