

Supplemental materials – Mohan et al.

Otolith geochemistry reflects life histories of Pacific bluefin tuna

John A. Mohan^{1,2}, Heidi Dewar³, Owyn E. Snodgrass³, Nathan R. Miller⁴, Yosuke Tanaka⁵, Seiji Ohshimo^{4,5}, Jay R. Rooker¹, Malcom Francis⁶, R. J. David Wells¹

¹ Texas A&M University at Galveston, Department of Marine Biology, 1001 Texas Clipper Rd., Galveston, TX 77553, USA

² University of New England, School of Marine and Environmental Programs, 11 Hills Beach Rd, 04005

³ Southwest Fisheries Science Center, National Marine Fisheries Service, 8901 La Jolla Shores Dr., La Jolla, CA 92037, USA

⁴ The University of Texas at Austin, Jackson School of Geosciences, 2275 Speedway Stop C9000, Austin, TX 78712, USA

⁵ Highly Migratory Resources Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, Fisheries Research and Education Agency, 5-7-1, Orido, Shimizu, Shizuoka, 424-8633, Japan

⁶ Pelagic Fish Resources Division, Fisheries Stock Assessment Center, Fisheries Resources Institute, Fisheries Research and Education Agency, 1551-8 Taira-machi, Nagasaki, 851-2213, Japan

⁷ National Institute of Water and Atmospheric Research (NIWA), Private Bag 14901, Wellington 6021, New Zealand

Keywords: *Thunnus orientalis*, migration, biomineralization, trace elements, otolith chemistry

Supplemental materials – Mohan et al.

Supplemental Figure 1 (below). Elemental profiles of individual PBT otoliths collected from the Eastern Pacific (EPO), Western Pacific (WPO) and South Pacific (SPO); different colored lines represent individual fish from each region. See Table 1 for details on collection. Asterisk (*) indicates visible crystal bond on surface of NZ02 that was removed from Figure 1.











