

Title

Host genetic factors related to innate immunity, environmental sensing and cellular functions are associated with human skin microbiota

Author list

Lucas Moitinho-Silva^{1,2}, Frauke Degenhardt¹, Elke Rodriguez², Hila Emmert², Simonas Juzenas^{1,3}, Lena Möbus², Florian Uellendahl-Werth¹, Nicole Sander², Hansjörg Baurecht⁴, Lukas Tittmann⁵, Wolfgang Lieb⁶, Christian Gieger^{7,8}, Annette Peters⁷, David Ellinghaus¹, Corinna Bang¹, Andre Franke^{1#*}, Stephan Weidinger^{2#*} and Malte Christoph Rühlemann^{1#}

Affiliations

¹Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany

²Department of Dermatology and Allergy, University Hospital Schleswig-Holstein, Kiel, Germany

³Institute of Biotechnology, Life Science Centre, Vilnius University, Vilnius, Lithuania

⁴Department for Epidemiology and Preventive Medicine, University of Regensburg, Regensburg, Germany

⁵Biobank PopGen and Institute of Epidemiology, Kiel University, Kiel, Germany

⁶Institute of Epidemiology, Kiel University, Kiel, Germany

⁷Institute of Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

⁸Research Unit of Molecular Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

#Authors contributed equally.

*Corresponding authors: AF, a.franke@mucosa.de; SW, sweidinger@dermatology.uni-kiel.de.

Supplementary Table

Supplementary Table 1. Samples collected for microbial profiling.

	KORA FF4 (N = 635)	PopGen (N = 1021)
Dry		
Dorsal forearm	-	260 (25)
Volar forearm	-	251 (25)
Moist		
Antecubital fossa	318 (50)	258 (25)
Sebaceous		
Forehead	-	252 (25)
Retroauricular fold	317 (50)	-

Percentages are shown (%).