



Figure S1: The schematic of phage treatment in *S. Typhimurium*-infected mouse model established in this study. After one-week acclimatization, 6-8 week-old female C57BL/6NJcl mice were orally fed with 100 μ L of 200 mg/mL streptomycin sulfate one day before *S. Typhimurium* oral infection with 100 μ L of 10^9 CFU/mL (10^8 CFU/mouse). Infected mice were divided into 4 groups (7 mice per group) and orally fed 100 μ L of 10^{11} PFU/mL (10^{10} PFU/mouse) of ST-W77, SE-W109, ST-W77 combined with SE-W109, and SM buffer as the indicated time point. All mice were euthanized on day 4 post-infection.

Table S1. The list of bacterial and phage strains used in this study

Designation	Genotype and relevant characteristic	Reference
S. Typhimurium IR715 (S. Tm)	Nalidixic acid-resistant derivative of ATCC 14028	(1)
ST-W77	Myovirus within the Viunalikevirus	(2)
SE-W109	Siphovirus within the Jerseylikevirus	(2)

Table S2. The list of primers used in this study

Organism	Target gene	Primers	Reference
<i>Homo sapiens</i>	<i>GAPDH</i>	5'-CCAGGAAATGAGCTTGACAAAGT-3' 5'-CCCACTCCTCCACCTTGAC-3'	(3)
	<i>IL-8</i>	5'-GCCAACACAGAAATTATTGTAAAGCTT-3' 5'-CCTCTGCACCCAGTTTCCTT-3'	(4)
	<i>MIP-3A</i>	5'-CTGCTTGATGTCAGTGCTGCTAC-3' 5'-CTGCCGTGTGAAGCCCACAATAAA-3'	(5)
	<i>IL-1B</i>	5'-AACAGATGAAGTGCTCCTCCAGG-3' 5'-TGGAGAACACCACTTGTGCTCCA-3'	(6)
<i>Mus musculus</i>	<i>Gapdh</i>	5'-TGTAGACCATGTAGTTGAGGTCA-3' 5'-AGGTCGGTGTGAACGGATTG-3'	(7)
	<i>Kc</i>	5'-TGCACCCAAACCGAAGTCAT-3' 5'-TTGTCAGAACGCCAGCGTTCAC-3'	
	<i>Nos2</i>	5'-CCAGCCTTGCATCCTCATTGG-3' 5'-CCAAACACCAAGCTCATGCGG-3'	
	<i>IL-17</i>	5'- GCTCCAGAAGGCCCTCAGA-3' 5'- AGCTTCCCTCCGCATTGA-3'	(8)
	<i>IFN-g</i>	5'- TCAAGTGGCATAGATGTGGAAGAA-3' 5'- TGGCTCTGCAGGATTTCATG-3'	(8)
	<i>Zo-1</i>	5'-CAGGGCTTTGGAGGAA-3' 5'-TACACGATCGTGGCAATAAAC-3'	(9)

Table S3. The criteria for mouse cecal histopathological scoring used in this study (10)

Score	Neutrophils infiltration	Mononuclear leukocyte infiltration	Submucosal edema	Epithelial damage	Exudate
0	No changes (0-5)	No changes (0-5)	No changes	No changes	No changes
1	6-20	6-10	Detectable (<10%)	Desquamation	Slight accumulation
2	21-60	11-20	Mild (10-20%)	Mild erosion and mild loss of goblet cells / undifferentiated enterocytes hyperplasia	Mild accumulation
3	61-100	21-40	Moderate (21-40%)	Marked erosion and moderate loss of goblet cells / undifferentiated hyperplasia	Moderate accumulation
4	>100	>40	Marked (>40%)	Ulceration and marked loss of goblet cells / undifferentiated hyperplasia	Marked accumulation

References:

1. Stojiljkovic I, Baumler AJ, Heffron F. Ethanolamine utilization in *Salmonella typhimurium*: nucleotide sequence, protein expression, and mutational analysis of the cchA cchB eutE eutJ eutG eutH gene cluster. *J Bacteriol.* 1995;177(5):1357-66.
2. Phothaworn P, Supokaivanich R, Lim J, Klumpp J, Imam M, Kutter E, et al. Development of a broad-spectrum *Salmonella* phage cocktail containing Viunalike and Jerseylike viruses isolated from Thailand. *Food Microbiol.* 2020;92:103586.
3. Raffatellu M, Chessa D, Wilson RP, Dusold R, Rubino S, Baumler AJ. The Vi capsular antigen of *Salmonella enterica* serotype Typhi reduces Toll-like receptor-dependent interleukin-8 expression in the intestinal mucosa. *Infection and immunity.* 2005;73(6):3367-74.
4. Stylianou E, Yndestad A, Sikkeland LI, Bjerkeli V, Damas JK, Haug T, et al. Effects of interferon-alpha on gene expression of chemokines and members of the tumour necrosis factor superfamily in HIV-infected patients. *Clin Exp Immunol.* 2002;130(2):279-85.
5. Johanesen PA, Dwinell MB. Flagellin-independent regulation of chemokine host defense in *Campylobacter jejuni*-infected intestinal epithelium. *Infection and immunity.* 2006;74(6):3437-47.
6. Jung HC, Eckmann L, Yang SK, Panja A, Fierer J, Morzycka-Wroblewska E, et al. A distinct array of proinflammatory cytokines is expressed in human colon epithelial cells in response to bacterial invasion. *J Clin Invest.* 1995;95(1):55-65.
7. Winter SE, Thiennimitr P, Winter MG, Butler BP, Huseby DL, Crawford RW, et al. Gut inflammation provides a respiratory electron acceptor for *Salmonella*. *Nature.* 2010;467(7314):426-9.
8. Winter SE, Thiennimitr P, Nuccio SP, Haneda T, Winter MG, Wilson RP, et al. Contribution of flagellin pattern recognition to intestinal inflammation during *Salmonella enterica* serotype typhimurium infection. *Infection and immunity.* 2009;77(5):1904-16.
9. Inagaki-Ohara K, Sawaguchi A, Suganuma T, Matsuzaki G, Nawa Y. Intraepithelial lymphocytes express junctional molecules in murine small intestine. *Biochemical and biophysical research communications.* 2005;331(4):977-83.
10. Buddhasiri S, Sukjoi C, Kaewsakhorn T, Nambunmee K, Nakphaichit M, Nitisinprasert S, et al. Anti-inflammatory Effect of Probiotic *Limosilactobacillus reuteri* KUB-AC5 Against *Salmonella* Infection in a Mouse Colitis Model. *Front Microbiol.* 2021;12:716761.