S3 Fig. Forrest plots for prevalence of ACO among patients with COVID-19.

a. USA

								Prevalence
Study or Subgroup	Prevalence	SE	Incidence	Total	Weight	Prevalence	Year	IV, Random, 95%CI
Wang L 2020	3.3	0.7	21	641	16.7%	3.30 [1.93, 4.67]	2020	•
Hussein MH 2020	16	0.9	292	1827	15.3%	16.00 [14.24, 17.76]	2020	
Zhu Z 2020	1	0.5	5	502	18.0%	1.00 [0.02, 1.98]	2020	•
Ho KS 2021	0.5	0.1	54	10523	19.4%	0.50 [0.30, 0.70]	2021	†
Ferastraoaru D 2021	0.4	0.1	19	4558	19.4%	0.40 [0.20, 0.60]	2021	†
Cao L 2021	7.9	1.5	27	343	11.1%	7.90 [4.96, 10.84]	2021	*
Total (95% CI)			418	18394	100.0%	4.24 [2.74, 5.73]		•
Heterogeneity: $Tau^2 = 2.98$; $Chi^2 = 336.99$, $df = 5$ (P < 0.00001); $I^2 = 99\%$							-	
Test for overall effect: $Z = 5.56$ (P < 0.00001)								0 50 100 Frequency (%)

b. UK

