

Table S1. Distribution of patients according to therapeutic indication and socio-demographic characteristics of patients who accepted (Group 1) and refused (Group 2) to participate in a clinical trial.

	Group 1 (N=80)	Group 2 (N=20)	p-value
Distribution of patients, n (%)			
Cardiology, n (%)	8 (10.0%)	1 (5.0%)	
Myocardial infarction	8 (10.0%)	1 (5.0%)	
Dermatology, n (%)	3 (3.75%)	1 (5.0%)	
Hidradenitis suppurativa	3 (3.75%)	1 (5.0%)	
Endocrinology, n (%)	14 (17.50%)	3 (15.0%)	
Diabetes mellitus	7 (8.75%)	3 (15.0%)	
Hypoparathyroidism	7 (8.75%)	0 (0.0%)	
Gynecology, n (%)	3 (3.75%)	0 (0.0%)	
Contraception	3 (3.75%)	0 (0.0%)	
Neurology, n (%)	46 (57.50%)	14 (70.0%)	
Migraine	5 (6.25%)	0 (0.0%)	
Multiple sclerosis	1 (1.25%)	0 (0.0%)	
Refractory epilepsy	1 (1.25%)	0 (0.0%)	
Stroke	39 (48.75%)	14 (70.0%)	
Oncology, n (%)	6 (7.50%)	1 (5.0%)	
Breast cancer	3 (3.75%)	0 (0.0%)	
Colorectal cancer	0 (0.0%)	1 (5.0%)	
Gastrointestinal cancer	1 (1.25%)	0 (0.0%)	
Lymphoma	1 (1.25%)	0 (0.0%)	
Pancreatic cancer	1 (1.25%)	0 (0.0%)	
Socio-demographic characteristics, n (%)			
Age (years), median (IQR)	64 (18.2)	68 (11.2)	.028
Gender, n (%)			1.00
Male	42 (52.5%)	10 (50.0%)	
Female	38 (47.5%)	10 (50.0%)	
Habits, n (%)			
Smoking	17 (21.3%)	3 (15.0%)	.76
Coffee intake	57 (71.3%)	12 (60.0%)	.48
Alcohol consumption	35 (43.8%)	5 (25.0%)	.20
Physical activity	43 (53.8%)	10 (50.0%)	.96
Employment Status, n (%)			.21
Employed	31 (38.75%)	8 (40.0%)	
Unemployed	6 (7.5%)	4 (20.0%)	
Retired	42 (52.5%)	8 (40.0%)	
Monthly Income, n (%)			.72
<250€	7 (8.75%)	3 (15.0%)	
250-500€	22 (27.5%)	3 (15.0%)	
501-1000€	30 (37.5%)	10 (50.0%)	
1001-1500€	8 (10.0%)	2 (10.0%)	
1501-2500€	3 (3.75%)	1 (5.0%)	
Academic Qualifications, n (%)			.16
4 years	30 (37.5%)	11 (55.0%)	
6 years	10 (12.5%)	0 (0.0%)	
9 years	14 (17.5%)	6 (30.0%)	
12 years	14 (17.5%)	2 (10.0%)	
Bachelor	3 (3.75%)	0 (0.0%)	
Master or equivalent educational level	8 (10.0%)	1 (5.0%)	
PhD	1 (1.25%)	0 (0.0%)	

Abbreviation: IQR, interquartile range.