

Supplementary Table 2. Duration of infectious diseases occurring in previous 12 months in patients with hypocortisolism and in the subgroups of patients with primary adrenal insufficiency (PAI) and secondary adrenal insufficiency (SAI). Categorical variables are expressed as percentages and frequencies. p^* values for comparisons with controls. p^{\S} values for comparisons between PAI and SAI. After correction for multiple testing a $p < 0.05$ was considered as significant statistically different (bold). ^{a,b,c} significantly different duration compared with controls at post hoc analysis using multiple z-tests with Bonferroni correction.

	Controls (n=570)	Hypocortisolism (n=135)	p^*	PAI (n=32)	p^*	SAI (n=103)	p^*	p^{\S}
Upper respiratory tract infections (URTIs) - duration								
less than 1 week	65% (279/429) ^a	53% (55/103)		35% (8/23) ^a		59% (47/80)		
from 1 to 2 weeks	28% (122)	39% (40)		45.5% (10)		37.5% (30)		
from 2 to 3 weeks	4% (19)	5% (5)	0.459	13% (3)	0.036	2.5% (2)	0.748	0.058
more than 3 weeks	2% (9) ^b	3% (3)		8.5% (2) ^b		1% (1)		
Lower respiratory tract infections (LRTIs) - duration								
less than 1 week	30% (19/64)	7% (1)		33% (2/6)		9% (1/11)		
from 1 to 2 weeks	55% (35)	33% (5)		17% (1)		36% (4)		
from 2 to 3 weeks	6% (3) ^c	35% (6) ^c	0.033	33% (2)	0.291	36% (4)	0.143	0.611
more than 3 weeks	9% (6)	18% (3)		17% (1)		18% (2)		
Gastrointestinal tract infections (GITs)- duration								
less than 1 week	88% (163/185)	75% (51/68)		65% (11/17)		65% (11/17)		
from 1 to 2 weeks	11% (20)	23.5% (13)		29% (5)		29% (5)		
from 2 to 3 weeks	0.5% (1)	0% (0)	0.200	0% (0)	0.083	0% (0)	0.736	0.161
more than 3 weeks	0.5% (1)	1.5% (1)		6% (1)		6% (1)		
Skin infections 1 (SSTIs-1) - duration								
less than 1 week	56% (79/142)	50% (19/38)		38.5% (5/13)		9% (1/11)		
from 1 to 2 weeks	30% (42)	32% (12)		23% (3)		36% (4)		
from 2 to 3 weeks	6% (9)	5% (2)	1.000	15% (2)	0.469	36% (4)	0.141	0.098
more than 3 weeks	8% (12)	13% (5)		23% (3)		18% (0)		
Skin infections 2 (SSTIs-2) - duration								
less than 1 week	56% (49/88)	67% (12/18)		100% (3/3)		56% (14/25)		
from 1 to 2 weeks	33% (29)	28% (5)		0% (0)		36% (9)		
from 2 to 3 weeks	2% (2)	5% (1)	0.795	0% (0)	0.844	0% (1)	0.879	0.407
more than 3 weeks	9% (8)	0% (0)		0% (0)		8% (2)		
Mycosis (MYC) - duration								
less than 1 week	33% (21/63)	33% (10/30)		43% (3/7)		30% (7/23)		
from 1 to 2 weeks	48% (30)	10% (3)		29% (2)		4% (1)		
from 2 to 3 weeks	5% (3)	40% (12)	0.076	0% (0)	0.893	52% (12)	0.228	0.052
more than 3 weeks	14% (9)	17% (5)		29% (2)		13% (3)		
Sexually transmitted infections (STIs) - duration								
less than 1 week	22% (2/9)	33% (1/3)		0% (0/1)		50% (1/2)		
from 1 to 2 weeks	67% (6)	33% (1)		0% (0)		50% (1)		
from 2 to 3 weeks	0% (0)	33% (1)	0.547	0% (0)	0.347	0% (0)	0.897	0.223
more than 3 weeks	11% (1)	33% (1)		100% (1)		0% (0)		
Urinary tract infections (UTIs) duration								
less than 1 week	70% (59/86)	73% (30/41)		75% (9/12)		72% (21/29)		
from 1 to 2 weeks	24% (21)	22% (9)		25% (3)		21% (6)		
from 2 to 3 weeks	2% (2)	2.5% (1)	1.000	20% (0)	1.000	3% (1)	1.000	0.824
more than 3 weeks	5% (4)	2.5% (1)		0% (0)		3% (1)		