

**Supplementary Table 1.** Parameter measurements of study participants in secondary analysis, *Mean (SD)*

| <b>Age range</b> | <i>N</i> | <i>PCV</i><br>[mm/s] | <i>MDIA</i><br>[mm] | <i>MaxPD</i><br>[mm] | <i>MinPD</i><br>[mm] | <i>PD range</i><br>[mm] |
|------------------|----------|----------------------|---------------------|----------------------|----------------------|-------------------------|
| < 40             | 13       | 4.33 (0.72)          | 0.37 (0.15)         | 6.51 (1.02)          | 3.06 (0.38)          | 3.45 (0.90)             |
| 40 - 49          | 4        | 3.70 (0.75)          | 0.27 (0.12)         | 5.57 (0.96)          | 3.09 (0.55)          | 2.48 (0.62)             |
| 50 - 59          | 7        | 4.35 (0.68)          | 0.32 (0.08)         | 6.08 (0.54)          | 3.18 (0.61)          | 2.90 (0.53)             |
| 60 - 69          | 9        | 3.33 (0.57)          | 0.24 (0.06)         | 5.03 (0.47)          | 3.02 (0.49)          | 2.00 (0.33)             |
| >= 70            | 7        | 3.74 (0.65)          | 0.25 (0.10)         | 5.14 (1.25)          | 2.79 (0.58)          | 2.35 (0.69)             |

PCV: Peak Constriction Velocity; MDIA: Mid-dilated Iris Activity; MaxPD: Maximum pupil diameter; MinPD: minimum pupil diameter; PD: Pupil diameter; SD: Standard Deviation

