

Supplemental table 1. A comparison of prescribed anticoagulants among patients with AF

	≤65 years (N=43)	66–75 years (N=102)	76–85 years (N=215)	>85 years (N=230)	p-value
Any anticoagulant, n (%)	35 (81.4)	88 (86.3)	171 (79.5)	159 (69.1)	0.003
Warfarin, n (%)	15 (34.9)	48 (47.1)	90 (41.9)	78 (33.9)	0.10
DOAC, n (%)	20 (46.5)	40 (39.2)	81 (37.7)	81 (35.2)	0.55
Type of DOAC					0.044
Dabigatran, n (%)	1 (5.0)	2 (5.3)	3 (3.8)	0 (0.0)	
Rivaroxaban, n (%)	5 (25.0)	6 (15.8)	9 (11.1)	2 (2.5)	
Apixaban, n (%)	7 (35.0)	11 (28.9)	32 (41.0)	35 (43.2)	
Edoxaban, n (%)	7 (35.0)	21 (52.5)	37 (45.7)	44 (54.3)	
Low dose DOAC, n (%)	1 (5.0)	11 (27.5)	13 (16.0)	6 (7.4)	0.014

Frequency count n (%) for categorical variables.

AF, atrial fibrillation; DOAC, direct oral anticoagulant