

Supplementary table A: National Malignant Mesothelioma Incidence, stratified for sex

	MPM (M)	MPM (F)	MPM (T)	MPeM (M)	MPeM (F)	MPeM (T)
1989	5,26	0,64	2,6	0,33	0,08	0,19
1990	4,87	0,53	2,41	0,38	0,1	0,23
1991	5,15	0,46	2,45	0,26	0,1	0,17
1992	6,95	0,59	3,3	0,46	0,17	0,3
1993	5,66	0,6	2,75	0,27	-	0,12
1994	6,03	0,7	2,94	0,38	0,1	0,22
1995	6,19	0,68	3	0,31	0,07	0,19
1996	6,74	0,66	3,28	0,35	0,04	0,18
1997	6,16	0,66	3,03	0,22	0,04	0,12
1998	5,83	0,45	2,8	0,38	0,05	0,2
1999	6,79	0,74	3,38	0,37	0,09	0,21
2000	6,55	0,71	3,28	0,24	0,14	0,19
2001	6,34	0,92	3,25	0,35	0,07	0,2
2002	5,59	0,77	2,83	0,46	0,08	0,24
2003	6,6	0,72	3,25	0,23	0,07	0,14
2004	6,8	0,62	3,36	0,39	0,13	0,24
2005	7,94	0,81	3,9	0,28	0,07	0,17
2006	7,24	0,88	3,7	0,37	0,11	0,22
2007	6,77	0,85	3,47	0,31	0,04	0,16
2008	7,1	0,86	3,65	0,27	0,08	0,17
2009	6,75	0,73	3,44	0,35	0,03	0,17
2010	7,74	0,68	3,76	0,32	0,15	0,22
2011	7,31	0,8	3,65	0,32	0,1	0,18
2012	6,94	0,84	3,6	0,25	0,13	0,18
2013	7,09	0,8	3,61	0,22	0,06	0,13
2014	6,67	0,77	3,42	0,28	0,15	0,21
2015	6,44	0,87	3,37	0,28	0,07	0,16
2016	6,88	0,93	3,61	0,17	0,09	0,13
2017	5,9	0,87	3,15	0,25	0,13	0,19
2018	6,58	0,77	3,35	0,21	0,13	0,18
1989-2018	6,6	0,74	3,28	0,3	0,09	0,18

National malignant mesothelioma incidence, per 100,000 person years (RESR). MPM= malignant pleural mesothelioma; MPeM= malignant peritoneal mesothelioma; M= male; F= female; T= total.

Supplementary table B: Annual Percentage Change for malignant mesothelioma incidence

Type	Sex	Age	From (year)	To (year)	APC (%)	95%CI LL	95%CI UL	P-value
Both	M	0-54	1989	2018	-6.2	-7.2	-5.2	<0.01
Both	M	55-64	1989	2009	-1.2	-2.5	0.1	0.1
			2009	2018	-9.4	-13.1	-5.4	<0.01
Both	M	65-79	1989	2006	3.4	2.3	4.5	<0.01
			2006	2018	-1.8	-3.5	-0.1	<0.01
Both	M	≥80	1989	2018	3.3	2.5	4.1	<0.01
Both	F	0-54	1989	2018	-0.3	-2.1	1.5	0.7
Both	F	55-64	1989	2018	-0.1	-1.4	1.1	0.8
Both	F	65-79	1989	2018	1.8	0.9	2.6	<0.01
Both	F	≥80	1989	2018	4.6	2.2	6.9	<0.01
MPM	M	0-54	1989	2018	-6.1	-7.6	-4.6	<0.01
MPM	M	55-64	1989	2011	-1.6	-2.7	-0.4	<0.01
			2011	2016	-16.9	-28.9	-0.3	<0.01
			2016	2018	52.3	-6.9	149.0	0.1
MPM	M	65-79	1989	1991	22.1	5.9	40.7	<0.01
			1991	1994	-6.2	-18.7	8.1	0.3
			1994	1998	8.2	0.7	16.1	<0.01
			1998	2001	-6.4	-18.8	7.9	0.3
			2001	2004	14.0	-1.1	31.4	0.1
			2004	2018	-1.8	-2.5	-1.2	<0.01
MPM	M	≥80	1989	2018	3.2	2.3	4.1	<0.01
MPM	F	0-54	1989	2018	-0.8	-2.5	1.0	0.4
MPM	F	55-64	1989	2018	0.0	-1.4	1.3	1.0
MPM	F	65-79	1989	2018	1.6	0.7	2.5	<0.01
MPM	F	≥80	1989	1991	188.6	-11.1	837.2	0.1
			1991	2018	3.1	1.0	5.2	<0.01
Both	M+F	All ages	1889	2010	1.6	1	2.1	<0.01
			2010	2018	-1.7	-3.9	0.6	0.1
MPM	M+F	All ages	1989	2007	1.8	1.1	2.5	<0.01
			2007	2018	-1	-2.5	0.4	0.2
MPM	M	All ages	1989	2009	1.4	0.7	2.0	<0.01
			2009	2018	-1.8	-3.9	0.4	0.1
MPeM	M+F	All ages	1989	2018	-0.5	-1.4	0.4	0.2
MPeM	M	All ages	1989	2018	-1.3	-0.4	-2.9	<0.01
MPeM	F	All ages	1989	2018	13.2	-7.1	37.9	0.2

MPM=malignant pleural mesothelioma, MPeM=malignant peritoneal mesothelioma, M=male, F=female, APC=annual percentage change, CI=confidence interval, LL=lower limit, UL=upper limit.

Supplementary table C: Age specific malignant mesothelioma incidence rates

	Male				Female			
	0-54	55-64	65-79	80+	0-54	55-64	65-79	80+
1989	1.14	10.67	17.1	19.88	0.21	1.39	1.66	3.31
1990	1.05	10.44	17.17	14.79	0.11	1.52	2.34	0.62
1991	0.81	9.75	19.84	15.2	0.15	0.97	2.18	0.29
1992	1.12	12.65	27.97	20.17	0.15	1.53	2.41	2.23
1993	0.7	11.56	20.51	20.76	0.05	0.95	2.05	2.92
1994	0.54	11.67	23.12	24.53	0.12	1.1	2.72	3.82
1995	0.83	11.83	21.76	26.06	0.08	1.22	3.03	2.15
1996	0.81	15.5	23.27	24.51	0.1	1.08	2.55	2.5
1997	0.71	12.12	23.12	20.64	0.15	1.21	1.99	3.29
1998	0.71	11.39	23.78	17.3	0.08	0.53	2.28	1.09
1999	0.72	10.22	31.61	16.2	0.2	1.56	2.7	2.18
2000	0.64	12.88	27.08	16.45	0.1	1.27	3.95	1.07
2001	0.54	11.53	25.12	24.88	0.12	1.36	4.18	2.6
2002	0.53	8.13	24.35	22.86	0.14	1.7	2.86	2.54
2003	0.3	10.91	26.75	27.5	0.22	0.86	3.11	1.6
2004	0.56	10.27	30.67	21.61	0.15	2.05	1.69	2.93
2005	0.3	9.78	36.39	30.45	0.04	1.74	3.59	2.38
2006	0.26	10.21	33.33	26.61	0.19	1.93	2.72	4.41
2007	0.42	8.08	31.48	24.2	0.11	1.28	3.54	2.93
2008	0.43	9.81	31.74	24.93	0.13	1.62	3.31	3.43
2009	0.36	8.68	30	28.57	0.07	1.12	2.95	2.89
2010	0.28	7.42	34.06	39.97	0.09	1.36	3.18	2.71
2011	0.3	7.86	28.42	46.43	0.08	1.29	3.24	4.18
2012	0.34	8.5	30.36	29.74	0.14	1.29	3.29	4.78
2013	0.18	7.89	28.6	41.15	0.1	1.2	3.11	3.95
2014	0.35	5.14	29.76	35.1	0.14	0.91	4	2.63
2015	0.15	4.64	28.4	38.39	0.17	1	3.42	4.09
2016	0.22	6.04	28.39	40.63	0.1	0.89	3.62	6.36
2017	0.12	3.19	25.84	38.59	0.08	1.33	3.45	5.59
2018	0.18	4.4	25.1	49.54	0.15	1.05	3.15	4.3
1989-2018	0.49	9.02	27.22	29.79	0.12	1.28	2.97	3.12

Age specific malignant mesothelioma (pleural and peritoneal combined) incidence, per 100,000 person years (RESR).