



Supplemental Figure 1. Meta-analysis forest plots of selected associations between herbicide use and risk of all non-Hodgkin lymphoma (NHL) and NHL subtypes (individual-study estimates are odds ratios (OR) and 95% confidence intervals (CI) from logistic regression models, with adjustment for study center, age, gender, socioeconomic status (SES), race/ethnicity, and set of covariates for use of other pesticides; meta-OR (overall) estimates are from random effects meta-analysis). NHL Subtypes: DLBCL=diffuse large B-cell lymphoma; FL=follicular lymphoma; MM=multiple myeloma; OBCL=other B-cell lymphoma; TCL=T-cell lymphoma.


Any herbicide use, all NHL




Any herbicide use TCL




Phenoxy herbicide use, all NHL




Phenoxy herbicide use duration >8 years, all NHL




Phenoxy herbicide use, MM




Phenoxy herbicide use duration >8 years, MM




2,4-D use, all NHL




2,4-D use duration >8 years, all NHL (BCMM excluded due to 0% weight)




2,4-D use duration >8 years, DLBCL




2,4-D use duration >8 years, OBCL




Glyphosate use, all NHL




Glyphosate lagged-use (10-year lag), all NHL




Glyphosate use, FL




Glyphosate lagged-use (10-year lag), FL




Glyphosate lagged-use (10-year lag) duration ≤8 years, FL (NCISEER excluded due to 0% weight)



Glyphosate lagged-use (10-year lag) duration >8 years, FL



Glyphosate lagged-use (10-year lag), DLBCL



Glyphosate lagged-use (10-year lag), TCL

