

Supporting Information

Thermosensitive Liposomes Encapsulating Nedaplatin and Picoplatin Demonstrate Enhanced Cytotoxicity Against Breast Cancer Cells

Sherif Ashraf Fahmy^{1,2}, Eduard Preis³, Alice Abu Dayyih³, Mohamed Alawak³, Hassan Mohamed El-Said Azzazy¹, Udo Bakowsky^{3,*}, Tamer Shoeib^{1*}

¹ Department of Chemistry, American University in Cairo (AUC), AUC Avenue, P.O. Box 74, New Cairo 11835, Egypt

² School of Life and Medical Sciences, University of Hertfordshire Hosted by Global Academic Foundation, R5 New Garden City, New Administrative Capital, AL109AB, Cairo 11835, Egypt

³ Department of Pharmaceutics and Biopharmaceutics, University of Marburg, Robert-Koch-Str. 4, 35037 Marburg, Germany

* Corresponding authors

Udo Bakowsky: ubakowsky@aol.com

Tamer Shoeib: t.shoeib@aucegypt.edu




Figure S1. Average cell viability of MDA-MB-231 cells at 37 °C and 40 °C after treatment with void thermosensitive liposomes and *p*-SC4 for all runs for 24 h. All experiments were conducted in triplicates, the mean values of all experiments were calculated, and error bars represent the standard deviation.