

Supplemental Digital Content 2 Demographics

Parameter Category	HARD [N=107]	No treatment [N=33]	Overall [N=140]
Age (years)			
Mean (SD)	48.3 (12.39)	44.4 (15.30)	47.4 (13.18)
Median	50.0	44.0	49.5
Min, Max	20, 70	22, 73	20, 73
Sex, n (%)			
Female	95 (88.8%)	30 (90.9%)	125 (89.3%)
Male	12 (11.2%)	3 (9.1%)	15 (10.7%)
BMI (kg/m²)			
Mean (SD)	24.32 (3.952)	24.59 (3.925)	24.38 (3.933)
Median	23.62	24.74	23.75
Min, Max	17.9, 41.3	19.3, 32.5	17.9, 41.3
Race, n (%)			
White	85 (79.4%)	22 (66.7%)	107 (76.4%)
Black or African American	12 (11.2%)	5 (15.2%)	17 (12.1%)
Asian	7 (6.5%)	2 (6.1%)	9 (6.4%)
Native Hawaiian or Other Pacific Islander	1 (0.9%)	0	1 (0.7%)
Other	2 (1.9%)	4 (12.1%)	6 (4.3%)
Ethnicity, n (%)			
Hispanic or Latino	26 (24.3%)	5 (15.2%)	31 (22.1%)
Not Hispanic or Latino	81 (75.7%)	28 (84.8%)	109 (77.9%)
Fitzpatrick Skin Type, n (%)			
I	7 (6.5%)	1 (3.0%)	8 (5.7%)
II	27 (25.2%)	8 (24.2%)	35 (25.0%)
III	30 (28.0%)	10 (30.3%)	40 (28.6%)
IV	24 (22.4%)	8 (24.2%)	32 (22.9%)
V	10 (9.3%)	4 (12.1%)	14 (10.0%)
VI	9 (8.4%)	2 (6.1%)	11 (7.9%)
Blinded Evaluator Galderma Chin Retrusion Scale Scores, n (%)			
1: Mild Retrusion	40 (37.4%)	16 (48.5%)	56 (40.0%)
2: Moderate Retrusion	67 (62.6%)	17 (51.5%)	84 (60.0%)
Treating Investigator Galderma Chin Retrusion Scale Scores, n (%)			
1: Mild Retrusion	42 (39.3%)	10 (30.3%)	52 (37.1%)
2: Moderate Retrusion	65 (60.7%)	23 (69.7%)	88 (62.9%)

BMI = body mass index; max = maximum; min = minimum; n = number of subjects; SD = standard deviation