

## Appendix


### Appendix 1 - Overview of coagulation proteins and their relation with paediatric sepsis and childhood mortality in literature

	Paediatric sepsis*	Non-survivors^	References
<b>Common pathway</b>			
Fibronectin	↓	↓	[1, 2]
<b>Contact activation (intrinsic) pathway</b>			
Factor IX	No data <sup>‡</sup>	No data	[3]
<b>Protein C pathway</b>			
Protein C	↓	↓	[4-6]
Protein S	↓	↓	[4, 5]
Thrombomodulin	↑	↑	[4, 7, 8]
<b>Platelet adhesion</b>			
ADAMTS-13	↓	↓	[9, 10]
vWF	↑	↑	[10]
<b>Fibrinolysis</b>			
PAI-1	↑	↑	[11-13]

\*Compared to controls; ^Compared to survivors; <sup>‡</sup> Only literature in adults

ADAMTS-13, a disintegrin and metalloprotease with thrombospondin type-1 motives; PAI-1, plasminogen activator inhibitor type 1; vWF, von Willebrand Factor

## Appendix 2 – Flowchart of patient selection



EUCLIDS, European Union Childhood Life-threatening Infectious Disease study; GAS, group A streptococcus; NL, the Netherlands; UK, United Kingdom

### Appendix 3 – Descriptive characteristics healthy controls vs invasive bacterial infections

	Invasive bacterial infections <b>n=241</b>	Healthy controls <b>n=64</b>	P-value
Female	110 (45.6)	30 (46.9)	0.97
Age in years, median [IQR]	3.3 (1.3-9.2)	5.4 (3.1-12.3)	<0.01
Ethnicity*			0.22
European	185 (76.8)	47 (73.4)	
Asian	13 (5.4)	4 (6.3)	
African	12 (4.9)	8 (12.5)	
Other	20 (8.3)	4 (1.7)	

\*Available data: Invasive bacterial infections n=230, healthy controls n=63

IQR, interquartile range

## Appendix 4 – Levels of haemostasis proteins

**Table Appendix 4A – Invasive bacterial infections vs healthy controls**


<b>Serum</b>	<b>Invasive bacterial infections</b> <b>n=241</b>	<b>Healthy controls</b> <b>n=64</b>	<b>P-value §</b>
	median (IQR)	median (IQR)	
Fibronectin (ug/ml)	41 (15 - 103)	106 (19-228)	0.02
PAI-1 (ng/ml)	96 (30 - 197)	77 (34-97)	0.00
Thrombomodulin (pg/ml)	10 (10 - 1490)*	10 (10-10)	0.00
ADAMTS13 (ng/ml)	21 (2-61)^	7 (2-17)	0.00
<b>Citrate plasma</b>		<b>n=122</b>	<b>n=20</b>
Factor IX (%)	60 (35 - 300)	40 (34-116)	0.36
Protein C (%)	40 (25 -65)	65 (54-89)	0.00
Protein S (%)	108 (65-315)	165 (95-315)	0.08
vWF (%)	300 (131 - 300)	88 (54-190)	0.00

\*n=239, ^n=240.

§ Adjusted for multiple testing.

ADAMTS-13, a disintegrin and metalloprotease with thrombospondin type-1 motives; IQR, interquartile range; PAI-1, plasminogen activator inhibitor type 1; vWF, von Willebrand Factor

**Figure Appendix 4A – Haemostasis protein levels in controls and patients with invasive bacterial infections**



A, Fibronectin; B, PAI-1; C, Thrombomodulin; D, ADAMTS-13; E, Factor IX; F, Protein C; G, Protein S; H, vWF

\*Extreme values for Fibronectin, thrombomodulin and Factor IX were truncated to the 99<sup>th</sup> centile.

ns; not significant. \*\*p-value≤0.05, \*\*\*p-value≤0.01, \*\*\*\*p-value≤0.001

p-values are adjusted for multiple testing.


**Table Appendix 4C – No invasive ventilation/inotropes vs need for invasive ventilation or inotropes**

	No invasive ventilation/inotropes n=118	Need for invasive ventilation or inotropes n=123	P-value*
<b>Serum</b>			
Fibronectin (ug/ml)	median (IQR) 59 (26-131)	median (IQR) 28 (9-74)	0.00
PAI-1 (ng/ml)	110 (22-190)	86 (35-198)	0.65
Thrombomodulin (pg/ml)	10 (10-729)	640 (10-2641)	0.00
ADAMTS13 (ng/ml)	16 (2-60)	27 (4-64)	0.44
<b>Citrate plasma</b>	<b>n=49</b>	<b>n=73</b>	
Factor IX (%)	110 (50-300)	50 (30-115)	0.01
Protein C (%)	55 (35-80)	35 (25-55)	0.01
Protein S (%)	160 (75-315)	90 (60-135)	0.03
vWF (%)	300 (85-300)	300 (245-300)	0.06

\*Adjusted for multiple testing

ADAMTS-13, a disintegrin and metalloprotease with thrombospondin type-1 motives; IQR, interquartile range; PAI-1, plasminogen activator inhibitor type 1; vWF, von Willebrand Factor

## Appendix 5 - Haemostasis protein levels in controls and different pathogens




Legend:

A, Fibronectin; B, PAI-1; C, Thrombomodulin; D, ADAMTS-13; E, Factor IX; F, Protein C;  
G, Protein S; H, vWF

\*Extreme values for Fibronectin, thrombomodulin and Factor IX were truncated to the 99<sup>th</sup> centile. ns, not significant, \*p-value≤0.05, \*\*p-value≤0.01, \*\*\*p-value≤0.001  
p-values are adjusted for multiple testing.

## Appendix 6 - Haemostasis protein levels in controls, survivors and non-survivors



A, Fibronectin; B, PAI-1; C, Thrombomodulin; D, ADAMTS-13; E, Factor IX; F, Protein C; G, Protein S; H, vWF

\*Extreme values for Fibronectin, thrombomodulin and Factor IX were truncated to the 99<sup>th</sup> centile. ns, not significant, \*p-value≤0.05, \*\*p-value≤0.01, \*\*\*p-value≤0.001. p-values are adjusted for multiple testing.

**Appendix 7 - Correlation of coagulation proteins with PIM2 score in all invasive bacterial infections (n=241)**

	PIM2 score	
	Spearman's rho	P-value*
Fibronectin	-0.25	0.00
PAI-1	-0.01	0.94
Thrombomodulin	0.2	0.02
ADAMTS-13	0.13	0.15
Factor IX	-0.24	0.04
Protein C	-0.25	0.03
Protein S	-0.29	0.01
vWF	0.14	0.25

\*P-values adjusted for multiple testing

ADAMTS-13, a disintegrin and metalloprotease with thrombospondin type-1 motives; IQR, interquartile range; PAI-1, plasminogen activator inhibitor type 1; vWF, von Willebrand Factor

Based on the Spearman's rho, significant correlations were classified as very weak (0-0.19), weak (0.2-0.39), moderate (0.4-0.59), strong (0.6-0.79) and very strong (0.8-1).[14]

**Appendix 8 – Characteristics of patients with available citrate plasma (n=122) and patients without available citrate plasma (n=110)**

All invasive bacterial infections n=241	Citrate plasma available n=122	Citrate plasma not available n=110	P-value
<b>Severity of illness</b>			
Admitted to PICU	177 (73.4)	103 (84.8)	<0.001
Need for invasive ventilation or inotropes		73 (59.8)	<0.01
PIM2 score (%), median [IQR]		3.5 (1.0-10.9)	3.5 (0.8-11.7) p=0.67
Death	21 (8.7)	8 (6.6)	13 (10.9) 0.33
<b>Pathogen</b>			
<i>Neisseria meningitidis</i>	83 (34.4)	43 (35.2)	40 (33.6) p=0.88
<i>Streptococcus pneumoniae</i>	64 (26.6)	34 (27.9)	30 (25.2)
<i>Staphylococcus aureus</i>	50 (20.7)	25 (20.5)	25 (21.0)
<i>Group A streptococcus</i>	44 (18.3)	20 (16.4)	24 (20.2)

IQR, interquartile range; PICU, paediatric intensive care unit; PIM, paediatric index of mortality

## References

1. Blanco A, Guisasola JA, Solis P, Bachiller R, Gonzalez H: **Fibronectin in meningococcal sepsis. Correlation with antithrombin III and protein C.** *Acta Paediatr Scand* 1990, **79**(1):73-76.
2. Riordan FA, Bestwick K, Thomson AP, Sills JA, Hart CA: **Plasma fibronectin levels in meningococcal disease.** *Eur J Pediatr* 1997, **156**(6):451-453.
3. Moore JX, Zakai NA, Mahalingam M, Griffin RL, Irvin MR, Safford MM, Baddley JW, Wang HE: **Hemostasis biomarkers and risk of sepsis: the REGARDS cohort.** *J Thromb Haemost* 2016, **14**(11):2169-2176.
4. Faust SN, Levin M, Harrison OB, Goldin RD, Lockhart MS, Kondaveeti S, Laszik Z, Esmon CT, Heyderman RS: **Dysfunction of endothelial protein C activation in severe meningococcal sepsis.** *N Engl J Med* 2001, **345**(6):408-416.
5. Leclerc F, Hazelzet J, Jude B, Hofhuis W, Hue V, Martinot A, Van der Voort E: **Protein C and S deficiency in severe infectious purpura of children: a collaborative study of 40 cases.** *Intensive Care Med* 1992, **18**(4):202-205.
6. Samransamruajkit R, Hiranrat T, Prapphal N, Sritippayawan S, Deerojanawong J, Poovorawan Y: **Levels of protein C activity and clinical factors in early phase of pediatric septic shock may be associated with the risk of death.** *Shock* 2007, **28**(5):518-523.
7. Lin JJ, Hsiao HJ, Chan OW, Wang Y, Hsia SH, Chiu CH: **Increased serum thrombomodulin level is associated with disease severity and mortality in pediatric sepsis.** *PLoS One* 2017, **12**(8):e0182324.
8. Kraft-Jacobs B, Brilli R: **Increased circulating thrombomodulin in children with septic shock.** *Crit Care Med* 1998, **26**(5):933-938.
9. Lin JJ, Chan OW, Hsiao HJ, Wang Y, Hsia SH, Chiu CH: **Decreased ADAMTS 13 Activity is Associated With Disease Severity and Outcome in Pediatric Severe Sepsis.** *Medicine (Baltimore)* 2016, **95**(16):e3374.
10. Bongers TN, Emonts M, de Maat MP, de Groot R, Lisman T, Hazelzet JA, Leebeek FW: **Reduced ADAMTS13 in children with severe meningococcal sepsis is associated with severity and outcome.** *Thromb Haemost* 2010, **103**(6):1181-1187.
11. Hermans PW, Hazelzet JA: **Plasminogen activator inhibitor type 1 gene polymorphism and sepsis.** *Clin Infect Dis* 2005, **41 Suppl 7**:S453-458.
12. Kornelisse RF, Hazelzet JA, Savelkoul HF, Hop WC, Suur MH, Borsboom AN, Risseeuw-Appel IM, van der Voort E, de Groot R: **The relationship between plasminogen activator inhibitor-1 and proinflammatory and counterinflammatory mediators in children with meningococcal septic shock.** *J Infect Dis* 1996, **173**(5):1148-1156.
13. Green J, Doughty L, Kaplan SS, Sasser H, Carcillo JA: **The tissue factor and plasminogen activator inhibitor type-1 response in pediatric sepsis-induced multiple organ failure.** *Thromb Haemost* 2002, **87**(2):218-223.
14. Wechsler S: **Statistics at Square One. Ninth Edition, revised by M. J. Campbell, T. D. V. Swinscow, BMJ Publ. Group, London, 1996. No. of pages: 140. Price: £11. ISBN 0-7279-0916-9.** *Statistics in Medicine* 1997, **16**(22):2629-2630.