

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed.
We post it as supplied by the authors.

Supplement to: Chun HM, Abutu A, Milligan K, et al. Low-level viraemia among people living with HIV in Nigeria: a retrospective longitudinal cohort study. *Lancet Glob Health* 2022; **10**: e1815–24.

Table of Contents

Table S1. Analysis of Exclusions.

Table S2. Low-Level Viremia and Subsequent Virologic Non-Suppression or Virologic Failure.

Table S3. Prevalence of Virologic Suppression, Low-Level Viremia, and Virologic Non-Suppression at First Overall Viral Load ≥ 24 Weeks after ART Initiation.

Table S4. Proportion (%) of Patients in Each Low-Level Viremia Range Among All Patients with Results in the Low-Level Viremia Range 51–999 copies per mL.

Figure S1. Prevalence of Viral Load Ranges Less Than 1,000 Copies/mL as a Proportion of All Viral Loads Less Than 1,000 Copies/mL, by Calendar Year and Years on ART*.

Table S5. Unadjusted and Adjusted Relative Risks of Low-Level Viremia Among Patients on First-Line ART.

Table S6. Adjusted Relative Risks of Virologic Non-Suppression and Virologic Failure Among Patients on First-Line ART, with Low-Level Viremia and INSTI Exposure Treated as Dichotomous Variables

Table S7. Viral Load Results Preceding Virologic Non-Suppression and Virologic Failure

Table S8. Unadjusted and Adjusted Relative Risks of Virologic Suppression and Virologic Failure Following Virologic Non-Suppression.

Table S9. Nigeria Low-Level Viremia Investigation Group

Table S1. Analysis of Exclusions.

Variable	Unit	Excluded*	Included	P value
N		88,837*	402,668	
Male sex	No. (% , 95% CI)	32,839 (37.0, 36.6–37.3)	129,620 (32.2, 32.0–32.3)	0.059
Age	Median (IQR)	33 (28–40)	34 (28–41)	0.64
State	No. (% , 95% CI)			<0.0001
	Benue	11,144 (12.5, 12.3–12.8)	100,569 (25.0, 24.8–25.1)	
	Delta	3,393 (3.8, 3.7–3.9)	14,906 (3.7, 3.6–3.8)	
	Ekiti	243 (0.3, 0.2–0.3)	2,269 (0.6, 0.5–0.6)	
	Enugu	5,490 (6.2, 6.0–6.3)	15,577 (3.9, 3.8–3.9)	
	FCT	9,598 (10.8, 10.6–11.0)	32,964 (8.2, 8.1–8.3)	
	Gombe	2,754 (3.1, 3.0–3.2)	11,251 (2.8, 2.7–2.8)	
	Imo	6,562 (7.4, 7.2–7.6)	11,227 (2.8, 2.7–2.8)	
	Kaduna	1,859 (2.1, 2.0–2.2)	22,759 (5.7, 5.6–5.7)	
	Katsina	464 (0.5, 0.5–0.6)	4,686 (1.2, 1.1–1.2)	
	Kogi	1,122 (1.3, 1.2–1.3)	8,256 (2.1, 2.0–2.1)	
	Lagos	6,033 (6.8, 6.6–7.0)	30,994 (7.7, 7.6–7.8)	
	Nasarawa	6,419 (7.2, 7.1–7.4)	25,047 (6.2, 6.1–6.3)	
	Ogun	1,575 (1.8, 1.7–1.9)	10,713 (2.7, 2.6–2.7)	
	Ondo	784 (0.9, 0.8–0.9)	6,651 (1.7, 1.6–1.7)	
	Osun	508 (0.6, 0.5–0.6)	4,539 (1.1, 1.1–1.2)	
	Oyo	1,688 (1.9, 1.8–2.0)	11,473 (2.8, 2.8–2.9)	
	Plateau	2,125 (2.4, 2.3–2.5)	14,443 (3.6, 3.5–3.6)	
	Rivers	27,076 (30.5, 30.2–30.8)	74,344 (18.5, 18.3–18.6)	

Demographic differences between included and excluded patients; age was analyzed using Wilcoxon rank-sum test

for clustered data. Categorical variables were analyzed using Rao-Scott χ^2 test.

* Patients younger than 18 or older than 99 (n = 29,707) were not included due to exclusion based on age.

Table S2. Low-Level Viremia and Subsequent Virologic Non-Suppression or Virologic Failure.

	≥1 VL Result in Reference Range	≥1 Subsequent VL after Reference Result	Subsequent Virologic Suppression ≤50 cp/mL	Subsequent Virologic Non- Suppression	≥2 Subsequent VL after Reference Result	Subsequent Virologic Failure
Reference Range						
≤50 cp/mL	190,639 (96.4)	175,626 (92.1)	151,512 (86.3)	13,894 (7.9)	97,674 (51.2)	1,426 (1.5)
LLV 51–999 cp/mL	69,405 (35.1)	55,161 (79.5)	43,436 (78.7)	5,595 (10.1)	29,958 (43.2)	819 (2.7)
51–199	50,039 (25.3)	38,410 (76.8)	29,524 (76.9)	3,282 (8.5)	20,398 (40.8)	430 (2.1)
200–399	14,601 (7.4)	10,989 (75.3)	7,656 (69.7)	1,091 (9.9)	6,034 (41.3)	171 (2.8)
400–999	13,552 (6.9)	10,261 (75.7)	6,970 (67.9)	1,287 (12.5)	5,524 (40.8)	219 (4.0)
≥1,000 cp/mL	19,199 (9.7)	11,399 (59.4)	7,862 (69.0)	2,232 (19.6)	--	--
All Ranges	--	197,729	--	19,199 (9.7)	112,316 (56.8)	2,232 (2.0)

Virologic non-suppression defined as ≥1,000 cp/mL at the next VL result following the relevant VL result; VF

defined as two consecutive non-suppressed results following the relevant VL result. LLV=low-level viremia.

VF=virologic failure. cp/mL=copies/mL. VL=viral load.

Table S3. Prevalence of Virologic Suppression, Low-Level Viremia, and Virologic Non-Suppression at First Viral Load ≥24 Weeks after ART Initiation.

Age at ART initiation—yr	Virologic Suppression ≤50 cp/mL, no. (%)	LLV 51–199 cp/mL, no. (%)	LLV 200–399 cp/mL, no. (%)	LLV 400–999 cp/mL, no. (%)	LLV 51–999 cp/mL, no. (%)	VLNS ≥1,000 cp/mL, no. (%)
First- and Second-Line ART: Males and Females (n = 402,668)						
Total	292,137 (72.6)	40,830 (10.1)	11,767 (2.9)	11,883 (3.0)	64,480 (16.0)	46,051 (11.4)
First-Line: Males and Females (n = 399,684, 99.3%)						
18-19 (n=7,065)	4,953 (70.1)	704 (10.0)	205 (2.9)	227 (3.2)	1,136 (16.1)	976 (13.8)
20-29 (n=116,524)	86,279 (74.0)	10,716 (9.2)	3,144 (2.7)	3,328 (2.9)	17,188 (14.8)	13,057 (11.2)
30-39 (n=150,320)	109,391 (72.8)	14,771 (9.8)	4,332 (2.9)	4,352 (2.9)	23,455 (15.6)	17,474 (11.6)
40-49 (n=81,842)	58,547 (71.5)	9,007 (11.0)	2,520 (3.1)	2,469 (3.0)	13,996 (17.1)	9,299 (11.4)
50-999 (n=43,933)	31,271 (71.2)	5,240 (11.9)	1,445 (3.3)	1,372 (3.1)	8,057 (18.3)	4,605 (10.5)
Total (n=399,684)	290,441 (72.7)	40,438 (10.1)	11,646 (2.9)	11,748 (2.9)	63,832 (16.0)	45,411 (11.4)
First-line: Males (n = 128,633, 32.2%)						
18-19 (n=1,959)	1,416 (72.3)	225 (11.5)	52 (2.7)	52 (2.7)	329 (16.8)	214 (10.9)
20-29 (n=28,504)	21,696 (76.1)	2,741 (9.6)	709 (2.5)	766 (2.7)	4,216 (14.8)	2,592 (9.1)
30-39 (n=45,688)	32,460 (71.0)	5,202 (11.4)	1,437 (3.1)	1,381 (3.0)	8,020 (17.6)	5,208 (11.4)
40-49 (n=34,165)	23,712 (69.4)	4,155 (12.2)	1,111 (3.3)	1,037 (3.0)	6,303 (18.4)	4,150 (12.1)
50-999 (n=18,317)	12,622 (68.9)	2,368 (12.9)	650 (3.5)	573 (3.1)	3,591 (19.6)	2,104 (11.5)
Total (n=128,633)	91,906 (71.4)	14,691 (11.4)	3,959 (3.1)	3,809 (3.0)	22,459 (17.5)	14,268 (11.1)
First-line: Females (n=271,051, 67.8%)						
18-19 (n=5,106)	3,537 (69.3)	479 (9.4)	153 (3.0)	175 (3.4)	807 (15.8)	762 (14.9)
20-29 (n=88,020)	64,583 (73.4)	7,975 (9.1)	2,435 (2.8)	2,562 (2.9)	12,972 (14.7)	10,465 (11.9)
30-39 (n=104,632)	76,931 (73.5)	9,569 (9.1)	2,895 (2.8)	2,971 (2.8)	15,435 (14.8)	12,266 (11.7)
40-49 (n=47,677)	34,835 (73.1)	4,852 (10.2)	1,409 (3.0)	1,432 (3.0)	7,693 (16.1)	5,149 (10.8)
50-999 (n=25,616)	18,649 (72.8)	2,872 (11.2)	795 (3.1)	799 (3.1)	4,466 (17.4)	2,501 (9.8)
Total (n=271,051)	198,535 (73.2)	25,747 (9.5)	7,687 (2.8)	7,939 (2.9)	41,373 (15.3)	31,143 (11.5)
Second-line: Males and Females (n=2,984, 0.7%)						
18-19 (n=58)	31 (53.4)	6 (10.3)	1 (1.7)	2 (3.4)	9 (15.5)	18 (31.0)
20-29 (n=437)	242 (55.4)	57 (13.0)	10 (2.3)	21 (4.8)	88 (20.1)	107 (24.5)
30-39 (n=1,088)	597 (54.9)	137 (12.6)	51 (4.7)	46 (4.2)	234 (21.5)	257 (23.6)
40-49 (n=895)	514 (57.4)	117 (13.1)	37 (4.1)	41 (4.6)	195 (21.8)	186 (20.8)
50-999 (n=506)	312 (61.7)	75 (14.8)	22 (4.3)	25 (4.9)	122 (24.1)	72 (14.2)
Total (n=2,984)	1,696 (56.8)	392 (13.1)	121 (4.1)	135 (4.5)	648 (21.7)	640 (21.4)
Second-line: Males (n=987, 33.1%)						
18-19 (n=26)	14 (53.8)	4 (15.4)	0 (0.0)	0 (0.0)	4 (15.4)	8 (30.8)
20-29 (n=98)	48 (49.0)	15 (15.3)	2 (2.0)	4 (4.1)	21 (21.4)	29 (29.6)
30-39 (n=239)	125 (52.3)	34 (14.2)	7 (2.9)	9 (3.8)	50 (20.9)	64 (26.8)
40-49 (n=351)	187 (53.3)	47 (13.4)	17 (4.8)	15 (4.3)	79 (22.5)	85 (24.2)
50-999 (n=273)	150 (54.9)	49 (17.9)	14 (5.1)	15 (5.5)	78 (28.6)	45 (16.5)
Total (n=987)	524 (53.1)	149 (15.1)	40 (4.1)	43 (4.4)	232 (23.5)	231 (23.4)
Second-line: Females (n=1,997, 66.9%)						
18-19 (n=32)	17 (53.1)	2 (6.2)	1 (3.1)	2 (6.2)	5 (15.6)	10 (31.2)
20-29 (n=339)	194 (57.2)	42 (12.4)	8 (2.4)	17 (5.0)	67 (19.8)	78 (23.0)
30-39 (n=849)	472 (55.6)	103 (12.1)	44 (5.2)	37 (4.4)	184 (21.7)	193 (22.7)
40-49 (n=544)	327 (60.1)	70 (12.9)	20 (3.7)	26 (4.8)	116 (21.3)	101 (18.6)
50-999 (n=233)	162 (69.5)	26 (11.2)	8 (3.4)	10 (4.3)	44 (18.9)	27 (11.6)
Total (n=1,997)	1,172 (58.7)	243 (12.2)	81 (4.1)	92 (4.6)	416 (20.8)	409 (20.5)

First VL result on first- or second-line regimens ≥24 weeks after ART initiation. ART=antiretroviral therapy. cp/mL=


copies/mL. LLV=low-level viremia. VS=virologic suppression. VLNS=virologic non-suppression.

Table S4. Proportion (%) of Patients in Each Low-Level Viremia Range Among All Patients with Results in the Low-Level Viremia Range 51–999 copies per mL.

	51–199 copies/mL	200–399 copies/mL	400–999 copies/mL
2016	54.0	25.1	20.8
2017	57.9	22.6	19.5
2018	63.3	18.7	18.0
2019	63.1	18.6	18.3
2020	65.9	17.0	17.1
2021	65.9	17.3	16.8

cp/mL= copies/mL. LLV=low-level viremia.

Figure S1. Viral Load Ranges Less Than 1,000 Copies/mL as a Proportion of All Viral Loads Less Than 1,000 Copies/mL, by Calendar Year and Years on ART*.



ART=antiretroviral therapy. cp/mL= copies/mL. LLV=low-level viremia. VS=virologic suppression.

* Proportions were calculated using the first VL result of every year in order to avoid bias from repeated measurements within 1 year.

Table S5. Unadjusted and Adjusted Relative Risks of Low-Level Viremia Among Patients on First-Line ART.

	Log-binomial regression			
	Unadjusted RR (95% CI)	P value	Adjusted RR (95% CI)	P value
Sex: male	1.12 (1.06–1.19)	0.0002	1.14 (1.10–1.18)	<0.0001
Age at ART initiation				
18–19	0.89 (0.81–0.97)	0.0094	0.93 (0.86–1.00)	0.063
20–29	0.87 (0.83–0.91)	<0.0001	0.90 (0.86–0.93)	<0.0001
30–39	0.92 (0.89–0.95)	<0.0001	0.93 (0.90–0.96)	<0.0001
40–49	0.98 (0.96–1.01)	0.11	0.98 (0.96–1.00)	0.062
≥50	ref		ref	
Initiated on DTG-based ART	0.78 (0.74–0.83)	<0.0001	0.83 (0.80–0.87)	<0.0001
Time on ART, years	1.10 (1.08–1.13)	<0.0001	1.07 (1.06–1.09)	<0.0001
First VL ≥12 months after ART initiation	0.97 (0.92–1.03)	0.31	--	--
Documented dispense ratio ≥51%	0.97 (0.94–0.99)	0.022	1.03 (0.99–1.05)	0.065

Log-binomial regression analysis, unadjusted and adjusted relative risks. Confidence intervals were calculated with

cluster-robust standard errors to account for clustering at the facility level.

ART=antiretroviral therapy. cp/mL= copies/mL. CI=confidence interval. RR= relative risk. LLV=low-level viremia.

VS=virologic suppression. VLNS=virologic non-suppression.

Table S6. Adjusted Relative Risks of Virologic Non-Suppression and Virologic Failure Among Patients on First-Line ART, with LLV and INSTI Exposure Treated as Dichotomous Variables.

Variable	Virologic Non-Suppression		Virologic Failure	
	Adjusted RR (95% CI)	P value	Adjusted RR (95% CI)	P value
Male sex	1.05 (1.01–1.10)	0.041	1.07 (0.97–1.19)	0.18
Age at ART initiation - yr				
18–19	1.29 (1.14–1.46)	<0.0001	1.71 (1.23–2.38)	0.0014
20–29	1.17 (1.09–1.25)	<0.0001	1.30 (1.10–1.55)	0.0024
30–39	1.13 (1.06–1.19)	<0.0001	1.22 (1.04–1.42)	0.013
40–49	1.04 (0.98–1.10)	0.17	1.05 (0.91–1.22)	0.50
≥50	ref	--		
Time on ART - yr	1.30 (1.26–1.34)	<0.0001	1.22 (1.14–1.30)	<0.0001
First VL ≥12 months after ART initiation	0.70 (0.67–0.74)	<0.0001	0.67 (0.59–0.76)	<0.0001
Documented dispense ratio ≥51%	--	--	0.85 (0.74–0.96)	0.012
INSTI exposure (cumulative)	0.73 (0.68–0.79)	<0.0001	0.40 (0.35–0.47)	<0.0001
LLV (51–999 cp/mL) at first VL result	1.61 (1.55–1.67)	<0.0001	2.20 (1.98–2.43)	<0.0001

Log-binomial analysis and adjusted relative risks using first on-treatment viral load after at least 24 weeks of ART.

Confidence intervals were calculated using cluster-robust standard errors to account for clustering at the facility level. Only patients with 3 or more VL measurements (n=112,316) were included in analysis of virologic failure, defined as 2 consecutive VL results ≥1,000 cp/mL.

Table S7. Viral Load Results Preceding Virologic Non-Suppression and Virologic Failure

	Range of Preceding Viral Load Result*				
	Virologic Suppression ≤50 cp/mL, no. (%)	LLV 51–199 cp/mL, no. (%)	LLV 200–399 cp/mL, no. (%)	LLV 400–999 cp/mL, no. (%)	LLV 51–999 cp/mL, no. (%)
Experienced VLNS (n=19,199, 9.7%)	13,752 (71.6)	3,172 (16.5)	1,040 (5.4)	1,235 (6.4)	5,447 (28.4)
Experienced VF (n=2,232, 2.0%)	1,419 (63.6)	425 (19.0)	169 (7.6)	219 (9.8)	813 (36.4)

Analysis of virologic non-suppression included all patients who met outcomes analysis inclusion criteria

(n=197,729); analysis of virologic failure only included patients who met these criteria and who had ≥3 VL results

(n=112,316).

LLV=low-level viremia. VLNS= virologic non-suppression. VF= virologic failure. cp/mL=copies/mL.

*For patients who experienced virologic failure, the VL result shown is that preceding the first of the two non-suppressed measurements

Table S8. Unadjusted and Adjusted Relative Risks of Virologic Suppression and Virologic Failure Following Virologic Non-Suppression.

	Virologic Suppression ≤50 cp/mL (n=7,862)				Virologic Failure (n=2,232)			
	Unadjusted RR (95% CI)	P value	Adjusted RR (95% CI)	P value	Unadjusted RR (95% CI)	P value	Adjusted RR (95% CI)	P value
Sex: male	0.93 (0.90–0.97)	0.0005	0.94 (0.91–0.98)	0.0028	1.04 (0.95–1.13)	0.40	--	--
Age at ART initiation - yr								
18–19	0.99 (0.87–1.12)	0.84	--	--	1.37 (1.05–1.78)	0.019	1.37 (1.05–1.78)	0.021
20–29	1.02 (0.96–1.08)	0.56	--	--	1.08 (0.94–1.24)	0.27	1.07 (0.93–1.23)	0.34
30–39	1.00 (0.95–1.06)	0.99	--	--	1.06 (0.93–1.21)	0.40	1.05 (0.92–1.20)	0.49
40–49	1.02 (0.97–1.08)	0.46	--	--	0.98 (0.85–1.12)	0.77	0.97 (0.84–1.11)	0.66
≥50	ref	--	--	--	ref	--	ref	--
Time on ART - yr	1.00 (0.98–1.01)	0.68	--	--	1.02 (0.98–1.06)	0.35	--	--
First VL ≥12 months after ART initiation	1.05 (1.02–1.09)	0.0033	1.03 (1.00–1.07)	0.045	0.92 (0.85–1.00)	0.056	0.93 (0.85–1.01)	0.085
Documented dispense ratio ≥51%	1.02 (0.99–1.06)	0.22	--	--	0.89 (0.81–0.99)	0.028	0.92 (0.84–1.02)	0.10
On DTG-based ART at time of VLNS	1.12 (1.08–1.16)	<0.0001	1.11 (1.07–1.14)	<0.0001	0.75 (0.68–0.82)	<0.0001	0.76 (0.70–0.83)	<0.0001
LLV 51–999 at first VL	0.82 (0.79–0.85)	<0.0001	0.83 (0.80–0.86)	<0.0001	1.34 (1.22–1.47)	<0.0001	1.32 (1.21–1.44)	<0.0001

Factors associated with virologic suppression ≤50 cp/mL and virologic failure following an episode of virologic non-suppression, patients with ≥1 VL after VLNS result only (n=11,399).

VL=viral load. INSTI=integrase strand transfer inhibitor. LLV=low-level viremia. VLNS=virologic non-suppression.

cp/mL=copies/mL. RR=relative risk.

Table S9. Nigeria Low-Level Viremia Investigation Group

Co-author name, highest degree	Affiliation
Uzoma Ene, MD	Division of Global HIV/TB, Center for Global Health, Abuja, Federal Capital Territory (FCT) Nigeria
Chibuzor Onyenuobi, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Timothy Efuntoye, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Chidozie Meribe, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Obinna Ogbanufe, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Orji Bassey, MSc	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Omodele J Fagbamigbe, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Ayodele Fagbemi, MPH	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Moyosola Bamidele, PhD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Israel Audu, MSc	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Dickson Adegoke, MSc	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Jelpe Tapdiyel, MSc	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Anuri Emeh, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Christopher Obanubi, MD	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Odafrenkhoa Oke, MSc	Division of Global HIV/TB, Center for Global Health, Abuja, FCT Nigeria
Sherri Pals, PhD	Division of Global HIV/TB, Center for Global Health, Atlanta Georgia USA
Andrew T Boyd, MD	Division of Global Health Protection, Center for Global Health Atlanta, Georgia USA
Clement Zeh, PhD	Division of Global HIV/TB, Center for Global Health, Atlanta Georgia USA
Dennis Ellenberger, PhD	Division of Global HIV/TB, Center for Global Health, Atlanta Georgia USA
Michelle Williams-Sherlock, MPH	Division of Global HIV/TB, Center for Global Health, Atlanta Georgia USA
Sylvia Adebajo, MD	Center for International Health, Education, and Biosecurity, Maryland Global Initiatives Corporation - an affiliate of the University of Maryland Baltimore, Abuja, FCT, Nigeria
Uba Sabo, PhD	National AIDS/STIs Control Programme (NASCP), Federal Ministry of Health, Abuja, FCT, Nigeria
Eruona Etubi, BMLS	National AIDS/STIs Control Programme (NASCP), Federal Ministry of Health, Abuja, FCT, Nigeria
Temiwoluwa Omole, MD	Clinical services department, APIN Public Health Initiatives, Abuja, FCT, Nigeria
Prosper Okonkwo, MD	Prevention, Care and Treatment Department, Institute of Human Virology, Abuja, FCT, Nigeria
Temitope Kolade, MD	Prevention, Care and Treatment Department, Institute of Human Virology, Abuja, FCT, Nigeria
Patrick Dakum, MD	Department of Epidemiology & Public Health, University of Maryland School of Medicine, Baltimore, Maryland, USA
Olayiwola Olanrewaju, MD	Clinical Services-Care and Treatment, Catholic Caritas Foundation of Nigeria, Abuja, FCT, Nigeria
John O Oko, MD	Health & HIV Division, Catholic Caritas Foundation of Nigeria, Abuja, FCT, Nigeria
Inyang Ayo, MD	Clinical Service Unit, Centre for Integrated Health Programs, Abuja, FCT, Nigeria

Bolanle Oyeledun, MD	Program Management, Centre for Integrated Health Programs, Abuja, FCT, Nigeria
Deborah Odoh, MD	National AIDS/STIs Control Programme (NASCP), Federal Ministry of Health, Abuja, FCT, Nigeria
Oluwasanmi Adeokun, MD	Center for International Health, Education, and Biosecurity, Maryland Global Initiatives Corporation - an affiliate of the University of Maryland Baltimore, Abuja, FCT, Nigeria
Grace Bassey, MSc	National AIDS/STIs Control Programme (NASCP), Federal Ministry of Health, Abuja, FCT, Nigeria