

Supplementary Table S3. Changes from study baseline in mUFC, LNSC, and random serum cortisol levels (extended evaluation population)

Parameter	Study Baseline		Change From Study Baseline to Month 6		<i>P</i> value*	Change From Study Baseline to Month 9		<i>P</i> value*	Change From Study Baseline to Month 12		<i>P</i> value*
	Mean (SD)	<i>n</i>	Mean (SD)	<i>n</i>		Mean (SD)	<i>n</i>		Mean (SD)	<i>n</i>	
mUFC, nmol/24h	528.1 (438.8)	60	−397.9 (461.4)	54	<0.0001	−383.9 (421.7)	49	<0.0001	−328.1 (475.6)	44	<0.0001
LNSC, nmol/L	11.7 (17.1)	58	−5.7 (18.5)	54	0.028	−4.9 (21.7)	47	0.131	−2.9 (24.2)	37	0.475
Random serum cortisol, nmol/L [†]	525.2 (150.0)	27	−71.5 (189.3)	22	0.091	−39.8 (165.8)	23	0.262	−74.3 (225.3)	16	0.207

*Two-sided *P* value from the paired t-test performed on the change from study baseline to Months 6, 9, and 12.

[†]No restriction on the clock time samples were collected.

LNSC, late-night salivary cortisol; mUFC, mean urinary free cortisol.