Supplementary Table S5. mUFC concentrations in patients who had prior radiation therapy at Months 6, 9, and 12 (extended evaluation population)

| | mUFC | | | | | |
|----------|-----------|------|-----------|------|-----------|------|
| | Month 6 | | Month 9 | | Month 12 | |
| Patient* | nmol/24 h | ×ULN | nmol/24 h | ×ULN | nmol/24 h | ×ULN |
| 1 | 210.5 | 1.5 | 96.0 | 0.7 | 277.4 | 2.0 |
| 2 | 161.3 | 1.2 | 215.4 | 1.6 | 195.1 | 1.4 |
| 3 | 160.2 | 1.2 | 129.4 | 0.9 | 102.3 | 0.7 |
| 4 | 129.8 | 0.9 | 181.1 | 1.3 | 164.7 | 1.2 |
| 5 | 109.7 | 0.8 | 134.3 | 1.0 | 216.6 | 1.6 |
| 6 | 46.6 | 0.3 | 48.4 | 0.4 | 37 | 0.3 |

ULN for UFC = 138 nmol/24 h.

mUFC, mean urinary free cortisol; UFC, urinary free cortisol; ULN, upper limit of normal.

^{*}One patient did not have mUFC values at Months 6, 9, and 12.