

Supplementary table 2. Treatment for sexually transmitted infections among patients attending a sexual assault centre in Oslo, Norway

	First treatment		Test after initial treatment	Second treatment		Test after second treatment
Chlamydia trachomatis						
58	Azithromycin	25	Negative 6 Positive 7	Doxycycline	7	Negative 3
	Doxycycline	23	Negative 14 Positive 2	Doxycycline	1	
	Moxifloxacin	1 ^a	Positive 1	Azithromycin	1	Positive 1
	Unspecified ^b	7	Negative 4	Doxycycline	1	Negative 1
	No information	2				
Mycoplasma genitalium						
24	Azithromycin	13	Negative 8			
	Moxifloxacin	1	Negative 1			
	Doxycycline	3	Positive 2 Negative 1	Azithromycin	1	
	Unspecified ^b	4	Negative 3			
	No information	3	Negative 2			
Mycoplasma genitalium macrolide resistant						
21	Azithromycin	4		Moxifloxacin	1	Negative 1
				Unspecified ^b	1	
	Moxifloxacin	10	Negative 9			
	Unspecified ^b	2	Negative 2			
	No information	5	Negative 2			
Neisseria gonorrhoeae						
5	Treated at specialist venereal clinic ^b	4				
	No information	1				

Total numbers at the primary examination and during follow-up: genital *Chlamydia trachomatis* in 58 patients, amongst whom 25 (43.1%) had symptoms; *Mycoplasma genitalium* in 45 patients, amongst whom 19 (42.2%) had symptoms; and *Neisseria gonorrhoeae* in 5 patients, amongst whom 2 (40.0%) had symptoms.

Missing data not shown for second treatment and tests after treatment.

^aCo-infection with macrolide resistant *Mycoplasma genitalium*.

^bAntibiotic treatment, drug not specified.

Among patients with no information about treatment, several may have received treatment elsewhere.