

**Supplemental Table 1.** Condition-specific ICD-9 and ICD-10 diagnostic codes used in this study

Ischemic stroke or transient ischemic attack  
**ICD-9-CM:** 362.3x, 433.x1, 434.x1, 436  
**ICD-10 CM:** H34.1, I63

Systemic embolism  
**ICD-9-CM:** 434.1, 444.xx, 445.xx  
**ICD-10 CM:** I74, I75

Myocardial infarction  
**ICD-9-CM:** 410.xx  
**ICD-10 CM:** I21.x(x), I22.x(x)

Hemorrhagic stroke  
**ICD-9-CM:** 430, 431, 432  
**ICD-10 CM:** I60, I61, I62

**Major bleeding**

1. Resulting in death  
**DEATHFM:** 38, 39, 40, 41, 42, 43, 72

2. Intracranial  
**ICD-9-CM:** 430, 431, 432.x

**ICD-10 CM:** I60, I61, I62

**Supplemental Table 2.** Crude number of events

Outcome	P2Y12 status at baseline	Events (apixaban arm)	Events (no anticoagulation arm)
Stroke (ischemic or hemorrhagic), transient ischemic attack, or systemic thromboembolism	Yes	<11	29
	No	22	70
Fatal or intracranial bleeding	Yes	<11	<11
	No	<11	37
Ischemic stroke or myocardial infarction	Yes	<11	119
	No	34	254