

Supplementary Material

Cultivable Endophytic Fungal Community

Associated with The Karst Endemic Plant Nervilia fordii

and Their Antimicrobial Activity

Ya-Qin Zhou^{1,2†}, Shao-Chang Yao^{1†}, Jie Wang^{1,3}, Xin-Yi Xie¹, Xiao-Ming Tan^{1*}, Rong-Shao Huang¹, Xin-Feng Yang¹, Yong Tan¹, Li-Ying Yu² and Peng Fu^{1,3}

¹ College of Pharmacy, Guangxi University of Chinese Medicine, Nanning, China

² Guangxi Key Laboratory of Medicinal Resources Conservation and Genetic Improvement, Guangxi Botanical Garden of Medicinal Plant, Nanning, China


³ Guangxi Zhuang Yao Key Laboratory of Medicine, Guangxi University of Chinese Medicine, Nanning, China

*Correspondence:


Xiaoming Tan (tanxm@gxtcmu.edu.cn)

[†]These authors have contributed equally to this work.


1 Supplementary Figures (A total of 3)



SUPPLEMENTARY FIGURE S1 ¹H (A) and ¹³C (B) NMR chromatogram of the monomer component collected from the EA crude extracts of *P. macrosclerotiorum*



SUPPLEMENTARY FIGURE S2 ¹H-¹H COSY (A) and HMBC (B) chromatogram of the monomer component collected from the EA crude extracts of *P. macrosclerotiorum*



SUPPLEMENTARY FIGURE S3 HSQC (A) and NOESY (B) chromatogram of the monomer component collected from the EA crude extracts of *P. macrosclerotiorum*