



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 16, 2022 – 11:43 am GMT

PDB ID : 7Z73
Title : Crystal structure of p63 tetramerization domain in complex with darpin 8F1
Deposited on : 2022-03-14
Resolution : 2.27 Å (reported)

This wwPDB validation report is for manuscript review

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtrriage (Phenix)	:	1.13
EDS	:	2.27
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.27


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	 13% 62% 15% 21%
1	B	61	 7% 75% 21%
1	C	61	 20% 64% 8% 28%
1	D	61	 2% 66% 5% 30%
2	E	126	 2% 87% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	126	 A horizontal bar chart representing the quality of the chain. The bar is divided into three segments: a small red segment on the left labeled '2%', a large green segment in the middle labeled '90%', and a small yellow segment on the right labeled '10%'. The bar ends with a small black dot.

For Manuscript Review

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 3565 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 2 of Tumor protein 63.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	48	420	265	74	79	2	0	0	0
1	B	48	420	267	74	77	2	0	0	0
1	C	44	382	243	64	73	2	0	0	0
1	D	43	374	239	63	70	2	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	GLY	-	expression tag	UNP Q9H3D4
A	357	SER	-	expression tag	UNP Q9H3D4
B	356	GLY	-	expression tag	UNP Q9H3D4
B	357	SER	-	expression tag	UNP Q9H3D4
C	356	GLY	-	expression tag	UNP Q9H3D4
C	357	SER	-	expression tag	UNP Q9H3D4
D	356	GLY	-	expression tag	UNP Q9H3D4
D	357	SER	-	expression tag	UNP Q9H3D4

- Molecule 2 is a protein called Darpin 8F1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	126	964	612	163	188	1	0	1	0
2	F	125	954	606	162	185	1	0	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total O 3 3	0	0
3	B	2	Total O 2 2	0	0
3	D	3	Total O 3 3	0	0
3	E	20	Total O 20 20	0	0
3	F	23	Total O 23 23	0	0

For Manuscript Review



3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoform 2 of Tumor protein 63




- Molecule 1: Isoform 2 of Tumor protein 63





- Molecule 1: Isoform 2 of Tumor protein 63




- Molecule 1: Isoform 2 of Tumor protein 63



- Molecule 2: Darpin 8F1



- Molecule 2: Darpin 8F1



For Manuscript Review

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.56Å 50.76Å 65.86Å 80.93° 79.92° 68.75°	Depositor
Resolution (Å)	47.06 – 2.27 47.06 – 2.27	Depositor EDS
% Data completeness (in resolution range)	92.6 (47.06-2.27) 92.6 (47.06-2.27)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.27Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.187 , 0.235 0.197 , 0.241	Depositor DCC
R_{free} test set	1065 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtrriage
Anisotropy	0.130	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3565	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/428	0.69	0/576
1	B	0.75	0/428	0.70	0/576
1	C	0.68	0/388	0.66	0/522
1	D	0.71	0/380	0.68	0/511
2	E	0.84	0/985	0.73	0/1338
2	F	0.79	0/972	0.70	0/1321
All	All	0.77	0/3581	0.70	0/4844

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	420	0	413	8	0
1	B	420	0	420	2	0
1	C	382	0	383	6	0
1	D	374	0	379	4	1
2	E	964	0	945	11	0
2	F	954	0	933	6	1
3	A	3	0	0	0	0
3	B	2	0	0	0	0
3	D	3	0	0	1	0
3	E	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	23	0	0	0	0
All	All	3565	0	3473	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:379:LYS:HE3	1:D:365:PRO:O	1.89	0.71
1:A:378:ILE:HD11	1:D:378:ILE:HD11	1.77	0.65
1:B:398:GLN:NE2	2:E:112:ASP:OD2	2.36	0.59
2:E:96:VAL:HG11	2:E:126:ALA:O	2.04	0.57
1:C:398:GLN:O	1:C:401:GLN:HG2	2.05	0.57
2:E:59:THR:O	2:E:59:THR:HG22	2.05	0.56
1:A:374:MET:CE	1:D:378:ILE:HG13	2.36	0.55
1:C:368:GLY:HA3	3:D:503:HOH:O	2.07	0.55
1:A:379:LYS:HE3	1:A:383:GLU:OE2	2.07	0.53
2:F:32:ALA:O	2:F:40:PRO:HD3	2.10	0.52
2:E:21:ARG:NH1	2:E:55:VAL:HG11	2.24	0.52
2:E:6:LYS:O	2:E:10:GLU:HG3	2.11	0.51
1:C:379:LYS:CE	1:D:365:PRO:O	2.60	0.50
2:F:66:ARG:HD3	2:F:72:THR:HG22	1.93	0.50
1:A:394:GLU:HA	1:A:394:GLU:OE1	2.12	0.49
1:A:401:GLN:O	1:A:405:GLN:HG2	2.12	0.49
2:F:59:THR:HG22	2:F:59:THR:O	2.12	0.49
1:B:394:GLU:HA	1:B:394:GLU:OE1	2.11	0.49
2:E:21:ARG:NH1	2:E:55:VAL:CG1	2.77	0.48
1:A:397:ARG:NE	1:C:380:GLU:OE2	2.46	0.47
2:E:32:ALA:O	2:E:40:PRO:HD3	2.14	0.47
1:A:389:PRO:HB3	2:F:46:TRP:CH2	2.50	0.46
1:A:385:MET:CE	1:C:380:GLU:HG2	2.47	0.44
2:F:98:ALA:O	2:F:105:THR:HA	2.18	0.43
2:E:48:GLY:HA2	2:E:85:ILE:HD12	1.99	0.43
2:E:48:GLY:HA2	2:E:85:ILE:CD1	2.50	0.41
2:E:98:ALA:O	2:E:105:THR:HA	2.21	0.41
2:E:52:ILE:HA	2:E:55:VAL:HG22	2.02	0.41
2:F:48:GLY:HA2	2:F:85:ILE:HD12	2.03	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:398:GLN:NE2	2:F:58:LYS:O[1_456]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	46/61 (75%)	46 (100%)	0	0	100	100
1	B	46/61 (75%)	46 (100%)	0	0	100	100
1	C	42/61 (69%)	41 (98%)	1 (2%)	0	100	100
1	D	41/61 (67%)	41 (100%)	0	0	100	100
2	E	125/126 (99%)	125 (100%)	0	0	100	100
2	F	123/126 (98%)	123 (100%)	0	0	100	100
All	All	423/496 (85%)	422 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	47/59 (80%)	46 (98%)	1 (2%)	53	68
1	B	47/59 (80%)	47 (100%)	0	100	100
1	C	43/59 (73%)	43 (100%)	0	100	100
1	D	42/59 (71%)	42 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	95/94 (101%)	94 (99%)	1 (1%)	73	84
2	F	94/94 (100%)	93 (99%)	1 (1%)	73	84
All	All	368/424 (87%)	365 (99%)	3 (1%)	81	90

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	401	GLN
2	E	67	ASP
2	F	67	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	48/61 (78%)	0.95	8 (16%) 1 2	49, 66, 120, 134	0
1	B	48/61 (78%)	0.70	4 (8%) 11 14	43, 59, 113, 133	0
1	C	44/61 (72%)	1.07	12 (27%) 0 0	48, 71, 154, 162	0
1	D	43/61 (70%)	0.61	1 (2%) 60 66	41, 59, 105, 117	0
2	E	126/126 (100%)	0.37	2 (1%) 72 77	39, 51, 70, 95	0
2	F	125/126 (99%)	0.46	3 (2%) 59 65	36, 51, 78, 99	0
All	All	434/496 (87%)	0.59	30 (6%) 16 21	36, 56, 108, 162	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	398	GLN	4.3
2	E	126	ALA	4.2
1	A	403	GLN	4.2
1	A	398	GLN	3.7
1	A	369	ARG	3.6
1	A	402	GLN	3.6
1	C	391	HIS	3.5
1	C	396	TYR	3.4
1	B	406	HIS	3.4
1	C	402	GLN	3.2
1	D	396	TYR	3.2
1	B	403	GLN	3.2
1	C	359	ASP	3.0
2	E	1	GLY	3.0
1	C	399	GLN	2.8
1	C	400	GLN	2.8
1	A	405	GLN	2.8
1	B	407	LEU	2.8
1	C	397	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	390	GLN	2.6
1	C	363	TYR	2.5
2	F	126	ALA	2.4
1	A	406	HIS	2.3
1	A	401	GLN	2.3
1	C	401	GLN	2.2
1	C	361	LEU	2.2
1	A	384	LEU	2.1
2	F	26	ASN	2.1
2	F	120	GLU	2.1
1	C	390	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.