

Immunological Pathogenesis of Bovine *E. coli* Infection in a Model of *C. elegans*

Hao Peng^{1†}, Huili Bai^{1†}, Yan Pan^{2†}, Jun Li^{1*}, Zhe Pei³, Yuying Liao¹, Cuilan Wu¹,
Changting Li¹, Li Tao¹, Shuhong Zhong¹, Chunxia Ma¹, Zhongwei Chen¹, Xiaoning Li¹,
Yu Gong⁴, Leping Wang¹, Fengsheng Li¹

*Correspondence:

Prof. Jun Li, Guangxi Key Laboratory of Veterinary Biotechnology, Guangxi

Veterinary Research Institute, Nanning 530001, China

Tel: 86-0771-3186349

E-mail: jlee9981@163.com

† These authors contribute equally to this paper.

¹Guangxi Key Laboratory of Veterinary Biotechnology, Guangxi Veterinary Research Institute, Nanning China; ²Guangxi Agricultural Vocational University, Nanning, China; ³The City College of New York, New York, USA and ⁴Animal Science and Technology Station of Guizhou, Guiyang, China.

Table S4. Longest survival time and LT50 of *Caenorhabditis elegans* infected with 13 strains of bovine *Escherichia coli*

strain	Longest survival time of <i>Caenorhabditis elegans</i>	LT50
<i>E.coli1</i>	7d	3d
<i>E.coli2</i>	7d	3.5d
<i>E.coli3</i>	7d	3d
<i>E.coli4</i>	8d	3.8d
<i>E.coli5</i>	11d	6d
<i>E.coli6</i>	8d	3.2d
<i>E.coli7</i>	9d	4.5d
<i>E.coli8</i>	7d	3.75d
<i>E.coli9</i>	10d	4d
<i>E.coli10</i>	9d	3d
<i>E.coli11</i>	10d	4.3d
<i>E.coli12</i>	8d	3.3d
<i>E.coli13</i>	10d	5d