

**Table S2: Plasmids used in this study**

<b>Plasmid number</b>	<b>Description</b>	<b>Source</b>
pLC3	ampR, HYGB	1,2
pLC136	ampR, <i>GPD1p</i> , <i>CYC1t</i> , <i>LEU2</i> (p415GPD)	3
pLC1543	<i>GPD1p-HAL9-CYCt</i> (ampR, <i>LEU2</i> )	This study
pLC1544	<i>GPD1p-HAL9<sup>C2214A</sup>-CYCt</i> (ampR, <i>LEU2</i> )	This study
pLC1545	<i>GPD1p-HAL9<sup>A1543T</sup>-CYCt</i> (ampR, <i>LEU2</i> )	This study
pLC1546	<i>GPD1p-HAL9<sup>A2479C</sup>-CYCt</i> (ampR, <i>LEU2</i> )	This study
pLC1609	<i>GPD1p-HSP12-CYCt</i> (ampR, <i>LEU2</i> )	This study

**References**

1. Wach A, Brachat A, Pöhlmann R, Philippsen P. New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast*. 1994;10(13):1793-1808. doi:10.1002/yea.320101310
2. Goldstein AL, McCusker JH. Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. *Yeast*. 1999;15(14):1541-1553. doi:10.1002/(SICI)1097-0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K
3. Mumberg D, Müller R, Funk M. Yeast vectors for the controlled expression of heterologous proteins in different genetic backgrounds. *Gene*. 1995;156(1):119-122. doi:https://doi.org/10.1016/0378-1119(95)00037-7

