

Supplementary Table 1. Frequency of local and systemic immediate adverse events by dose and age group.

Frequency of local immediate adverse events by dose and age group.			
	Dose 1 (n=963)		Dose 2 (n=624)
		Pain	
3-11 years	33 (3.8)		9 (1.7)
12-17 years	2 (2.2)		7 (8.2)
		Redness	
3-11 years	20 (2.3)		3 (0.6)
12-17 years	0		0
		Induration	
3-11 years	5 (0.6)		0
12-17 years	1 (1.1)		0
		Pruritus	
3-11 years	2 (0.2)		0
12-17 years	0		1 (1.2)
		Swelling	
3-11 years	0		0
12-17 years	1 (1.1)		0
		Other	
3-11 years	1 (0.1)		1 (0.2)
12-17 years	0		0
Frequency of systemic immediate adverse events by dose and age group.			
		Headache	
3-11 years	6 (0.7)		1 (0.2)
12-17 years	2 (2.2)		1 (1.2)
		Fatigue	
3-11 years	4 (0.5)		1 (0.2)
12-17 years	0		0

Muscle pain		
3-11 years	1 (0.1)	0
12-17 years	0	1 (1.2)
Nausea		
3-11 years	1 (0.1)	0
12-17 years	0	0
Skin or mucosa abnormality		
3-11 years	0	0
12-17 years	0	1* (1.2)

* **Generalized pruritus, without rash or other symptoms, mild and self-limited**

- *Total volunteers for first dose (n=963): 3-11 years (n=873) and 12-17 years (n=90)*
- *Total volunteers for second dose (n=624): 3-11 years (n=539) and 12-17 years (n=85)*
-