

All Panelists

The following document presents the group's median and range of ratings. Each cell is color coded. Cells in yellow are ones where the group disagreed (≥2 panelists gave that cell a rating of 1-3 and ≥2 panelists gave that cell a rating of 7-9); cells in blue are ones where the group agreed, with the darker blues representing higher medians, as shown in the key on the right.

Median	(Range)
Yellow: Disagreement (≥2 ratings of 1-3 and ≥2 ratings of 7-9)	
Blue 1: Median ≥7-9 without disagreement	
Blue 2: Median ≥4-<7, without disagreement	
Blue 3: Median 1-<4 without disagreement	

SECTION 1

TABLE 1. SOLID ORGAN CANCERS

Please enter a rating in each yellow cell, where each cell represents a cancer or cancer stage. If there is heterogeneity within a category, do your best to imagine a typical patient.		Q1: What is the likelihood that adults diagnosed with this cancer at this stage today will be cured [1] with typical treatment [2]?			Q2: What is the likelihood that untreated adults with stage I or II of this cancer will progress to stage III or IV disease over their lifetime?			Q3: How long does it take this cancer to progress from the beginning of this stage [3] to the beginning of the next stage in undiagnosed adults (natural history of disease)?			Imagine an annual hypothetical screening blood test for patients ≥50 years of age that is 100% sensitive and 100% specific for the stage of cancer you are being asked to rate. Assume this test cannot differentiate between higher and lower risk cancers. If all cancers of this type were diagnosed no later than the stage [4] indicated by the row heading, to what extent would... Q4: ... life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [6]? Assume patients are receiving typical treatment cure rates increase in all patients with this cancer, relative to cure rates without the screening test? Q5: Assume patients are receiving typical treatment .			Q6: Assume patients are receiving best available treatment [7].			Now consider what medical practice might be like in 2030 [5] with cure rates higher than today's. Q7: If all cancers of this type were diagnosed no later than this stage, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?		
		Lower risk [8]	Higher risk [9]	Overall	Lower risk	Higher risk	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall			
		A	B	C	D	E	F	G	H	I	J	K										
Prostate		1			3.5 (1 - 5)	7.0 (5 - 9)	6.0 (3 - 7)															
	Stage I	2	9.0 (9 - 9)	8.0 (5 - 9)	8.5 (7 - 9)			7.0 (5 - 8)	3.5 (1 - 6)	3.0 (1 - 5)	3.5 (1 - 5)	3.0 (1 - 7)										
	Stage II	3	9.0 (8 - 9)	7.0 (5 - 9)	8.0 (7 - 9)			5.0 (4 - 6)	4.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	4.5 (3 - 5)										
	Stage III	4	7.0 (5 - 9)	4.0 (3 - 6)	5.5 (4 - 8)			3.0 (2 - 5)	3.0 (2 - 5)	2.0 (1 - 5)	3.0 (2 - 5)	4.0 (3 - 6)										
	Stage IV	5	1.5 (1 - 7)	1.0 (1 - 4)	1.0 (1 - 5)																	
Breast		6			6.0 (5 - 7)	8.0 (5 - 9)	7.0 (7 - 8)															
	Stage I	7	9.0 (8 - 9)	8.0 (6 - 9)	9.0 (7 - 9)			3.0 (2 - 4)	7.0 (6 - 8)	6.0 (3 - 8)	6.5 (4 - 8)	7.5 (3 - 9)										
	Stage II	8	8.0 (8 - 9)	6.5 (4 - 8)	7.5 (6 - 8)			2.0 (1 - 3)	5.0 (4 - 6)	5.5 (2 - 7)	5.5 (3 - 7)	6.0 (4 - 7)										
	Stage III	9	6.5 (4 - 7)	4.0 (3 - 5)	5.5 (4 - 6)			1.5 (1 - 2)	3.0 (2 - 5)	3.0 (2 - 6)	3.0 (2 - 6)	5.0 (3 - 7)										
	Stage IV	10	1.5 (1 - 3)	1.0 (1 - 4)	1.0 (1 - 2)																	
Lung		11			7.0 (4 - 9)	9.0 (8 - 9)	8.0 (7 - 9)															
	Stage I	12	7.5 (7 - 9)	6.0 (4 - 7)	7.0 (6 - 9)			2.0 (2 - 3)	8.0 (7 - 9)	8.0 (6 - 9)	8.0 (6 - 9)	8.0 (3 - 9)										
	Stage II	13	5.0 (4 - 7)	4.5 (3 - 6)	5.0 (3 - 8)			1.0 (1 - 2)	6.0 (5 - 7)	6.0 (4 - 6)	7.0 (5 - 7)	7.0 (4 - 8)										
	Stage III	14	3.0 (1 - 5)	2.5 (1 - 4)	3.0 (1 - 5)			1.0 (1 - 1)	3.5 (3 - 5)	3.0 (2 - 5)	4.0 (2 - 6)	5.0 (3 - 6)										
	Stage IV	15	1.0 (1 - 2)	1.0 (1 - 1)	1.0 (1 - 2)																	

[1] Curability is the receipt of effective treatment such that a population of individuals who are "cured" would have the same life expectancy as a population that never had the cancer being considered. Some members of each of those populations (cured and never had cancer) might be expected to die prematurely (e.g., from trauma), but many more would be expected to die of more typical age-related causes.

[2] Includes active treatment or surveillance.

[3] For example, to answer 2G, consider the time from the beginning of stage I to the beginning of stage II. For 3G, consider the time from the beginning of stage II to the beginning of stage III. Similarly, for 4G, consider the time from the beginning of stage III to the beginning of stage IV.

[4] For stage I, assume all stage II-IV cancers are detected during stage I. For stage II, assume that all stage III and IV cancers are detected by stage II; any cancers that had been detected in stage I or II would still be detected during those stages. For the stage III, assume all stage IV cancers were diagnosed during stage III; any cancers that had been detected in stages I-III would still be detected during those stages.

[5] If there are many treatment advancements, cure rates may increase, and if treatment remains unchanged, cure rates are unlikely to change.

[6] A patient who is not cured could have increased life expectancy (e.g. still die of this cancer).

[7] Treatment consistent with NCCN guidelines.

[8] Please refer to the table above for examples of lower risk cancers.

[9] Please refer to the table above for examples of higher risk cancers.

Please enter a rating in each yellow cell, where each cell represents a cancer or cancer stage. If there is heterogeneity within a category, do your best to imagine a typical patient.		Q1: What is the likelihood that adults diagnosed with this cancer at this stage today will be cured [1] with typical treatment [2]?			Q2: What is the likelihood that untreated adults with stage I or II of this cancer will progress to stage III or IV disease over their lifetime?			Q3: How long does it take this cancer to progress from the beginning of this stage [3] to the beginning of the next stage in undiagnosed adults (natural history of disease)?			Q4: ... life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [6]?			Q5: Assume patients are receiving typical treatment ... cure rates increase in all patients with this cancer, relative to cure rates without the screening test?			Q6: Assume patients are receiving typical treatment Q6: Assume patients are receiving best available treatment [7].			Q7: If all cancers of this type were diagnosed no later than this stage, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?		
		For columns A through F , enter a rating of 1 through 9, where: 1=Extremely unlikely 3=Somewhat unlikely 5=Neutral, neither likely nor unlikely 7=Somewhat likely 9=Extremely likely						For column G , enter a rating of 1 through 9, where: 1=Less than 1 year 3=Slightly 5=5 years 9=9 or more years						For columns H through K , enter a rating of 1 through 9, where your answer reflects your judgement about how much the outcome of interest (e.g., life expectancy in Q4 and cure rates in Q5, Q6, Q7) would increase: 1=Not at all 3=Slightly 5=Moderately 7=Considerably 9=A great deal, doubling the outcome of interest								
		Lower risk [8]	Higher risk [9]	Overall	Lower risk	Higher risk	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall		
A	B	C	D	E	F	G	H	I	J	K												
Colon/Rectum	16																					
Stage I	17	9.0 (8 - 9)	9.0 (8 - 9)	9.0 (8 - 9)	7.0 (6 - 8)	8.0 (8 - 9)	8.0 (7 - 9)	3.5 (2 - 5)	7.0 (6 - 9)	7.0 (6 - 9)	7.5 (6 - 9)	8.0 (3 - 9)										
Stage II	18	8.0 (7 - 9)	7.0 (7 - 8)	8.0 (7 - 8)				3.0 (2 - 5)	6.0 (5 - 8)	6.0 (5 - 7)	6.0 (5 - 7)	7.0 (4 - 8)										
Stage III	19	6.0 (4 - 8)	5.0 (3 - 6)	5.0 (3 - 6)				1.0 (1 - 2)	5.0 (1 - 6)	4.0 (3 - 5)	5.0 (3 - 6)	6.0 (4 - 7)										
Stage IV	20	1.5 (1 - 4)	1.0 (1 - 2)	1.5 (1 - 3)																		
Melanoma	21				6.5 (4 - 8)	8.0 (6 - 9)	7.0 (6 - 9)															
Stage I	22	9.0 (8 - 9)	9.0 (7 - 9)	9.0 (8 - 9)				3.0 (1 - 5)	7.0 (3 - 8)	7.0 (6 - 8)	7.0 (6 - 9)	7.0 (3 - 8)										
Stage II	23	8.0 (7 - 8)	7.0 (6 - 8)	7.0 (6 - 8)				2.0 (1 - 4)	5.0 (2 - 6)	5.0 (5 - 6)	5.0 (5 - 8)	6.0 (5 - 7)										
Stage III	24	5.0 (3 - 6)	3.5 (2 - 6)	4.0 (3 - 6)				1.0 (1 - 2)	3.0 (1 - 5)	3.0 (2 - 5)	4.0 (3 - 5)	6.0 (3 - 7)										
Stage IV	25	2.0 (1 - 3)	1.0 (1 - 2)	1.5 (1 - 8)																		
Kidney	26				5.5 (4 - 8)	8.0 (7 - 9)	7.0 (6 - 8)															
Stage I	27	9.0 (8 - 9)	9.0 (4 - 9)	9.0 (7 - 9)				5.0 (1 - 7)	7.0 (3 - 8)	7.0 (5 - 8)	7.0 (5 - 9)	7.0 (3 - 8)										
Stage II	28	8.0 (7 - 9)	7.0 (4 - 8)	8.0 (6 - 8)				3.0 (1 - 5)	5.5 (4 - 7)	5.5 (4 - 6)	6.0 (5 - 7)	6.0 (5 - 7)										
Stage III	29	6.0 (5 - 7)	4.0 (3 - 6)	5.0 (4 - 6)				2.0 (1 - 2)	4.5 (3 - 5)	5.0 (3 - 6)	5.0 (3 - 6)	6.0 (3 - 7)										
Stage IV	30	2.0 (1 - 3)	1.0 (1 - 2)	2.0 (1 - 2)																		
Head and Neck	31				7.0 (5 - 8)	8.0 (6 - 9)	8.0 (6 - 9)															
Stage I	32	8.5 (7 - 9)	9.0 (7 - 9)	9.0 (7 - 9)				3.0 (2 - 6)	7.0 (4 - 9)	6.0 (4 - 9)	7.0 (5 - 8)	7.0 (3 - 8)										
Stage II	33	8.0 (6 - 9)	7.0 (5 - 8)	7.5 (6 - 8)				2.0 (1 - 4)	5.0 (3 - 7)	5.0 (3 - 7)	6.0 (4 - 7)	6.0 (4 - 7)										
Stage III	34	6.0 (3 - 7)	4.5 (3 - 6)	5.0 (3 - 6)				1.0 (1 - 2)	3.0 (2 - 6)	3.0 (2 - 6)	4.0 (3 - 7)	5.0 (4 - 7)										
Stage IV	35	4.0 (2 - 6)	2.0 (1 - 4)	3.0 (1 - 5)																		
Uterus	36				6.0 (3 - 9)	8.0 (6 - 9)	7.0 (5 - 9)															
Stage I	37	9.0 (8 - 9)	8.0 (7 - 9)	9.0 (8 - 9)				4.0 (3 - 5)	6.0 (2 - 7)	6.0 (2 - 7)	6.0 (2 - 8)	7.0 (2 - 8)										
Stage II	38	8.0 (7 - 9)	6.5 (5 - 8)	8.0 (7 - 8)				3.0 (1 - 5)	5.0 (3 - 6)	5.0 (3 - 6)	5.0 (4 - 7)	6.0 (4 - 7)										
Stage III	39	6.0 (5 - 9)	4.5 (2 - 5)	5.0 (5 - 7)				1.5 (1 - 3)	4.0 (3 - 6)	4.0 (2 - 6)	4.0 (3 - 6)	5.0 (3 - 8)										
Stage IV	40	2.0 (1 - 4)	1.0 (1 - 2)	1.0 (1 - 3)																		
Pancreas	41						9.0 (8 - 9)															
Stage I	42			4.0 (3 - 7)				1.0 (1 - 2)	5.0 (5 - 9)	5.0 (3 - 9)	6.0 (4 - 9)	6.5 (4 - 9)										
Stage II	43			2.5 (1 - 5)				1.0 (1 - 2)	3.0 (2 - 7)	3.0 (1 - 6)	4.0 (2 - 9)	5.0 (2 - 8)										
Stage III	44			1.0 (1 - 3)				1.0 (1 - 1)	2.0 (1 - 4)	1.0 (1 - 3)	2.0 (1 - 4)	4.0 (1 - 6)										
Stage IV	45			1.0 (1 - 1)																		

Please enter a rating in each yellow cell, where each cell represents a cancer or cancer stage. If there is heterogeneity within a category, do your best to imagine a typical patient.		Q1: What is the likelihood that adults diagnosed with this cancer at this stage today will be cured [1] with typical treatment [2]?			Q2: What is the likelihood that untreated adults with stage I or II of this cancer will progress to stage III or IV disease over their lifetime?			Q3: How long does it take this cancer to progress from the beginning of this stage [3] to the beginning of the next stage in undiagnosed adults (natural history of disease)?			Q4: ... life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [6]?			Q5: Assume patients are receiving typical treatment ... cure rates increase in all patients with this cancer, relative to cure rates without the screening test?			Q6: Assume patients are receiving typical treatment Q6: Assume patients are receiving best available treatment [7].			Q7: If all cancers of this type were diagnosed no later than this stage, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?		
		For columns A through F , enter a rating of 1 through 9, where: 1=Extremely unlikely 3=Somewhat unlikely 5=Neutral, neither likely nor unlikely 7=Somewhat likely 9=Extremely likely						For column G , enter a rating of 1 through 9, where: 1=Less than 1 year 5=5 years 9=9 or more years			For columns H through K , enter a rating of 1 through 9, where your answer reflects your judgement about how much the outcome of interest (e.g., life expectancy in Q4 and cure rates in Q5, Q6, Q7) would increase: 1=Not at all 3=Slightly 5=Moderately 7=Considerably 9=A great deal, doubling the outcome of interest											
		Lower risk [8]	Higher risk [9]	Overall	Lower risk	Higher risk	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall		
A	B	C	D	E	F	G	H	I	J	K												
Thyroid	46																					
Stage I	47	9.0 (9 - 9)	6.0 (1 - 7)	9.0 (8 - 9)	4.0 (1 - 7)	7.5 (6 - 9)	5.0 (2 - 7)	5.5 (4 - 8)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)	3.0 (1 - 7)			
Stage II	48	9.0 (8 - 9)	5.0 (1 - 6)	8.0 (7 - 9)				5.0 (3 - 7)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)	3.0 (1 - 5)			
Stage III	49	8.0 (7 - 9)	3.0 (1 - 4)	7.0 (6 - 9)				4.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)			
Stage IV	50	6.0 (1 - 7)	1.0 (1 - 2)	5.0 (1 - 7)																		
Bladder	51						7.0 (5 - 9)															
Stage I	52			8.5 (7 - 9)				3.0 (2 - 5)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	6.0 (5 - 7)	7.0 (5 - 8)			
Stage II	53			6.5 (6 - 8)				2.0 (1 - 5)	5.0 (4 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	4.0 (3 - 6)	6.0 (4 - 6)			
Stage III	54			4.0 (3 - 5)				1.0 (1 - 2)	3.0 (2 - 4)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	4.0 (3 - 7)			
Stage IV	55			1.0 (1 - 2)																		
Liver/Intrahepatic Bile-duct	56				7.0 (5 - 8)	8.0 (5 - 9)	7.5 (6 - 9)															
Stage I	57	5.0 (3 - 7)	3.5 (1 - 6)	4.0 (2 - 7)				2.0 (1 - 3)	6.5 (4 - 7)	6.0 (4 - 7)	6.0 (4 - 7)	6.5 (4 - 8)	6.5 (4 - 8)	6.5 (4 - 8)	6.5 (4 - 8)	6.5 (4 - 8)	6.5 (4 - 8)	6.5 (4 - 8)	7.0 (5 - 8)			
Stage II	58	3.0 (2 - 7)	2.0 (1 - 7)	3.0 (2 - 7)				1.0 (1 - 2)	5.0 (3 - 6)	4.5 (2 - 5)	4.5 (2 - 5)	5.0 (4 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	6.0 (4 - 6)			
Stage III	59	2.0 (1 - 5)	1.0 (1 - 5)	1.5 (1 - 5)				1.0 (1 - 1)	2.5 (2 - 5)	2.0 (1 - 5)	2.0 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	2.5 (1 - 5)	4.0 (3 - 6)			
Stage IV	60	1.0 (1 - 1)	1.0 (1 - 1)	1.0 (1 - 1)																		
Ovary	61				7.0 (5 - 9)	8.0 (8 - 9)	8.0 (7 - 9)															
Stage I	62	9.0 (6 - 9)	8.0 (3 - 8)	8.0 (7 - 9)				3.0 (1 - 3)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	8.0 (6 - 9)			
Stage II	63	8.0 (4 - 9)	6.0 (3 - 7)	7.0 (5 - 8)				2.0 (1 - 2)	6.0 (5 - 8)	5.0 (5 - 8)	5.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	6.0 (5 - 8)	7.0 (6 - 8)			
Stage III	64	7.0 (5 - 9)	2.0 (1 - 4)	3.0 (2 - 5)				1.0 (1 - 1)	3.0 (1 - 5)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	3.0 (1 - 6)	5.0 (3 - 7)			
Stage IV	65	5.5 (1 - 9)	1.0 (1 - 1)	1.0 (1 - 3)																		
Stomach	66				7.0 (6 - 8)	9.0 (7 - 9)	8.0 (7 - 9)															
Stage I	67	8.0 (6 - 9)	7.0 (5 - 7)	7.0 (6 - 8)				3.0 (2 - 5)	7.0 (6 - 8)	7.0 (5 - 8)	7.0 (5 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	7.0 (6 - 8)	8.0 (6 - 9)			
Stage II	68	5.0 (4 - 7)	4.0 (3 - 6)	4.5 (2 - 7)				2.0 (1 - 2)	5.0 (4 - 8)	5.0 (3 - 7)	5.0 (3 - 7)	5.5 (4 - 8)	5.5 (4 - 8)	5.5 (4 - 8)	5.5 (4 - 8)	5.5 (4 - 8)	5.5 (4 - 8)	5.5 (4 - 8)	6.0 (5 - 9)			
Stage III	69	3.0 (1 - 6)	2.0 (1 - 4)	2.0 (1 - 5)				1.0 (1 - 2)	3.0 (2 - 5)	3.0 (1 - 4)	3.0 (1 - 4)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	4.0 (2 - 6)			
Stage IV	70	1.0 (1 - 2)	1.0 (1 - 1)	1.0 (1 - 1)																		
Esophagus	71				7.5 (7 - 8)	9.0 (8 - 9)	8.0 (8 - 9)															
Stage I	72	7.0 (4 - 9)	6.0 (5 - 7)	7.0 (5 - 8)				2.5 (2 - 5)	7.0 (6 - 9)	7.0 (5 - 7)	7.0 (5 - 7)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)	7.0 (6 - 9)			
Stage II	73	4.0 (2 - 6)	3.5 (3 - 6)	4.0 (3 - 7)				1.0 (1 - 2)	4.5 (3 - 8)	4.5 (3 - 7)	4.5 (3 - 7)	5.0 (4 - 8)	5.0 (4 - 8)	5.0 (4 - 8)	5.0 (4 - 8)	5.0 (4 - 8)	5.0 (4 - 8)	5.0 (4 - 8)	6.0 (5 - 9)			
Stage III	74	2.5 (1 - 4)	2.0 (1 - 4)	2.0 (1 - 5)				1.0 (1 - 2)	2.0 (2 - 5)	2.0 (1 - 4)	2.0 (1 - 4)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	3.0 (2 - 5)	4.0 (3 - 6)			
Stage IV	75	1.0 (1 - 2)	1.0 (1 - 1)	1.0 (1 - 1)																		

Please enter a rating in each yellow cell, where each cell represents a cancer or cancer stage. If there is heterogeneity within a category, do your best to imagine a typical patient.		Q1: What is the likelihood that adults diagnosed with this cancer at this stage today will be cured [1] with typical treatment [2]?			Q2: What is the likelihood that untreated adults with stage I or II of this cancer will progress to stage III or IV disease over their lifetime?			Q3: How long does it take this cancer to progress from the beginning of this stage [3] to the beginning of the next stage in undiagnosed adults (natural history of disease)?			Q4: ... life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [6]?			Q5: Assume patients are receiving typical treatment ... cure rates increase in all patients with this cancer, relative to cure rates without the screening test?			Q6: Assume patients are receiving typical treatment Q6: Assume patients are receiving best available treatment [7].			Q7: If all cancers of this type were diagnosed no later than this stage, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?		
		For columns A through F , enter a rating of 1 through 9, where: 1=Extremely unlikely 3=Somewhat unlikely 5=Neutral, neither likely nor unlikely 7=Somewhat likely 9=Extremely likely						For column G , enter a rating of 1 through 9, where: 1=Less than 1 year 5=5 years 9=9 or more years						For columns H through K , enter a rating of 1 through 9, where your answer reflects your judgement about how much the outcome of interest (e.g., life expectancy in Q4 and cure rates in Q5, Q6, Q7) would increase: 1=Not at all 3=Slightly 5=Moderately 7=Considerably 9=A great deal, doubling the outcome of interest								
		Lower risk [8]	Higher risk [9]	Overall	Lower risk	Higher risk	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall	Overall		
A	B	C	D	E	F	G	H	I	J	K												
Gallbladder	76					9.0 (4 - 9)																
Stage I	77					5.0 (4 - 6)					2.0 (1 - 3)	7.0 (4 - 8)	6.0 (5 - 8)	6.5 (5 - 8)	7.0 (5 - 8)							
Stage II	78					3.0 (2 - 5)					1.0 (1 - 1)	4.5 (3 - 5)	4.0 (3 - 5)	5.0 (3 - 6)	5.0 (4 - 6)							
Stage III	79					2.0 (1 - 3)					1.0 (1 - 1)	2.0 (1 - 4)	2.0 (1 - 4)	2.0 (1 - 4)	3.0 (2 - 5)							
Stage IV	80					1.0 (1 - 1)																
Cervix	81					8.0 (7 - 9)																
Stage I	82					9.0 (8 - 9)					4.0 (1 - 5)	7.0 (5 - 8)	7.0 (5 - 8)	7.0 (6 - 8)	7.0 (5 - 8)							
Stage II	83					7.0 (6 - 8)					2.5 (1 - 4)	5.5 (4 - 7)	5.5 (4 - 7)	6.0 (5 - 7)	6.0 (5 - 7)							
Stage III	84					5.0 (4 - 6)					1.0 (1 - 2)	3.0 (3 - 6)	3.0 (2 - 6)	4.0 (2 - 6)	4.5 (3 - 7)							
Stage IV	85					1.0 (1 - 3)																
Sarcoma	86						5.5 (4 - 9)	8.0 (7 - 9)	7.5 (6 - 9)													
Stage I	87	8.5 (8 - 9)	7.0 (5 - 8)	8.0 (6 - 8)							3.5 (1 - 6)	7.0 (5 - 8)	7.0 (6 - 8)	7.0 (7 - 8)	8.0 (5 - 9)							
Stage II	88	7.0 (6 - 9)	6.0 (4 - 7)	7.0 (4 - 7)							2.0 (1 - 4)	5.0 (4 - 7)	5.0 (4 - 7)	6.0 (5 - 7)	6.0 (5 - 7)							
Stage III	89	4.5 (3 - 6)	3.0 (1 - 4)	4.0 (2 - 5)							1.0 (1 - 2)	3.0 (2 - 5)	3.0 (2 - 5)	3.5 (2 - 5)	4.5 (3 - 5)							
Stage IV	90	1.0 (1 - 3)	1.0 (1 - 1)	1.0 (1 - 2)																		
Urothelial Tract	91					8.5 (6 - 9)																
Stage I	92					8.0 (6 - 9)					3.0 (2 - 7)	7.0 (5 - 7)	7.0 (5 - 7)	7.0 (5 - 7)	7.0 (5 - 7)							
Stage II	93					5.5 (5 - 7)					2.0 (2 - 5)	5.0 (5 - 6)	5.0 (4 - 6)	5.0 (4 - 6)	5.5 (5 - 7)							
Stage III	94					4.0 (3 - 5)					1.0 (1 - 4)	3.0 (2 - 5)	3.0 (2 - 4)	3.0 (2 - 5)	4.0 (3 - 6)							
Stage IV	95					1.5 (1 - 3)																
Anus	96					8.0 (6 - 9)																
Stage I	97					9.0 (8 - 9)					3.0 (2 - 7)	5.5 (4 - 8)	6.0 (4 - 8)	6.0 (4 - 8)	7.0 (5 - 8)							
Stage II	98					8.0 (6 - 8)					2.0 (2 - 5)	5.0 (4 - 6)	5.0 (4 - 7)	5.0 (4 - 7)	5.0 (5 - 7)							
Stage III	99					5.0 (5 - 7)					1.0 (1 - 3)	3.5 (2 - 5)	3.0 (2 - 5)	3.5 (2 - 5)	4.0 (3 - 6)							
Stage IV	100					1.0 (1 - 3)																

SECTION 2

TABLE 2. HEMATOLOGIC CANCERS: LYMPHOMA

<p>Please enter a rating in each yellow cell, where each cell represents a cancer or cancer stage.</p> <p>If there is heterogeneity within a category, do your best to imagine a typical patient.</p>	<p>Q1: What is the likelihood that adults diagnosed with this cancer at this stage today will be cured [10] with typical treatment [11]?</p>	<p>Q2: What is the likelihood that untreated adults with stage I or II of this cancer will progress to stage III or IV disease over their lifetime?</p>	<p>Q3: How long does it take this cancer to progress from the beginning of this stage [12] to the beginning of the next stage in undiagnosed adults (natural history of disease)?</p>	<p>Imagine an annual hypothetical screening blood test for patients ≥50 years of age that is 100% sensitive and 100% specific for the stage of cancer you are being asked to rate. Assume this test cannot differentiate between higher and lower risk cancers.</p> <p>If all cancers of this type were diagnosed no later than the stage [13] indicated by the row heading, to what extent would...</p> <p>Q4: ...life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [15]?</p> <p>Assume patients are receiving typical treatment.</p>	<p>...cure rates increase in all patients with this cancer, relative to cure rates without the screening test?</p> <p>Q5: Assume patients are receiving typical treatment.</p>	<p>Q6: Assume patients are receiving best available treatment [16].</p> <p>Q7: If all cancers of this type were diagnosed no later than this stage, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?</p>	<p>For columns A through F, enter a rating of 1 through 9, where:</p> <p>1=Extremely unlikely 3=Somewhat unlikely 5=Neutral, neither likely nor unlikely 7=Somewhat likely 9=Extremely likely</p>						<p>For column G, enter a rating of 1 through 9, where:</p> <p>1=Less than 1 year 5=5 years 9=9 or more years</p>						<p>For columns H through K, enter a rating of 1 through 9, where your answer reflects your judgement about how much the outcome of interest (e.g., life expectancy in Q4 and cure rates in Q5, Q6, Q7) would increase:</p> <p>1=Not at all 3=Slightly 5=Moderately 7=Considerably 9=A great deal, doubling the outcome of interest</p>					
							Lower risk [17]	Higher risk [18]	Overall	Lower risk	Higher risk	Overall	Overall	Overall	Overall	Overall	Overall	Overall						
							A	B	C	D	E	F	G	H	I	J	K	K						
Lymphoma	101																							
Stage I	102	9.0 (8 – 9)	8.0 (5 – 8)	8.0 (7 – 9)			5.0 (3 – 6)	9.0 (7 – 9)	7.0 (6 – 8)															
Stage II	103	8.0 (7 – 9)	7.0 (3 – 8)	8.0 (5 – 8)						4.0 (2 – 6)	4.5 (2 – 8)	4.0 (3 – 8)	5.0 (3 – 8)	7.0 (5 – 7)										
Stage III	104	6.5 (3 – 8)	5.5 (1 – 7)	6.0 (3 – 7)						2.5 (1 – 5)	4.0 (3 – 6)	4.0 (3 – 6)	4.0 (3 – 7)	6.0 (3 – 7)										
Stage IV	105	5.0 (2 – 7)	5.0 (1 – 6)	5.0 (2 – 6)						2.0 (1 – 3)	3.0 (2 – 5)	3.0 (2 – 4)	3.5 (2 – 4)	5.0 (3 – 7)										

[10] Curability is the receipt of effective treatment such that a population of individuals who are “cured” would have the same life expectancy as a population that never had the cancer being considered. Some members of each of those populations (cured and never had cancer) might be expected to die prematurely (e.g., from trauma), but many more would be expected to die of more typical age-related causes.

[11] Includes active treatment or surveillance.

[12] For example, to answer 2G, consider the time from the beginning of stage I to the beginning of stage II. For 3G, consider the time from the beginning of stage II to the beginning of stage III. Similarly, for 4G, consider the time from the beginning of stage III to the beginning of stage IV.

[13] For stage I, assume all stage II-IV cancers are detected during stage I. For stage II, assume that all stage III and IV cancers are detected by stage II; any cancers that had been detected in stage I or II would still be detected during those stages. For the stage III, assume all stage IV cancers were diagnosed during stage III; any cancers that had been detected in stages I-III would still be detected during those stages.

[14] If there are many treatment advancements, cure rates may increase, and if treatment remains unchanged, cure rates are unlikely to change.

[15] A patient who is not cured could have increased life expectancy (e.g. still die of this cancer).

[16] Treatment consistent with NCCN guidelines.

[17] Please refer to the table above for examples of lower risk cancers.

[18] Please refer to the table above for examples of higher risk cancers.

TABLE 3. OTHER HEMATOLOGIC CANCERS

Please enter a rating in each yellow cell, where each cell represents a cancer type. If there is heterogeneity within a category, do your best to imagine a typical patient.	Q1: What is the likelihood that adults diagnosed with this cancer today will be cured [19] with typical treatment [20]?			Q2: What is the likelihood that untreated adults early [21] in their disease course with this cancer will progress and require treatment?			Imagine an annual hypothetical screening blood test for patients ≥50 years of age that is 100% sensitive and 100% specific for the cancer you are being asked to rate. Assume this test cannot differentiate between higher and lower risk cancers. If all cancers of this type were diagnosed earlier [21] in their disease course, to what extent would... Q3: ... life expectancy increase in all patients with this cancer, relative to life expectancy without the screening test [23]? Assume patients are receiving typical treatment cure rates increase in all patients with this cancer, relative to cure rates without the screening test? Q4: Assume patients are receiving typical treatment . Q5: Assume patients are receiving best available treatment [24].			Now consider what medical practice might be like in 2030 [22] with cure rates higher than today's. Q6: If all cancers of this type were diagnosed earlier in their disease course, to what extent would cure rates increase with typical treatment in all patients with this cancer, relative to 2030 cure rates without the screening test?		
	For columns A through F , enter a rating of 1 through 9, where: 1=Extremely unlikely 3=Somewhat unlikely 5=Neutral, neither likely nor unlikely 7=Somewhat likely 9=Extremely likely						For columns G through J , enter a rating of 1 through 9, where your answer reflects your judgement about how much the outcome of interest (e.g., life expectancy in Q3 and cure rates in Q4, Q5, Q6) would increase: 1=Not at all 3=Slightly 5=Moderately 7=Considerably 9=A great deal, doubling the outcome of interest								
		Lower risk [25] A	Higher risk [26] B	Overall C	Lower risk D	Higher risk E	Overall F	Overall G	Overall H	Overall I	Overall J				
ALL [27]	106		4.0 (3 – 5)			9.0 (1 – 9)	3.0 (1 – 5)	3.0 (1 – 5)	3.0 (1 – 5)	5.0 (4 – 5)					
CLL [28]	107	4.5 (1 – 7)	4.0 (1 – 7)	3.5 (1 – 7)	7.0 (1 – 9)	5.0 (1 – 8)	2.5 (1 – 5)	2.5 (1 – 5)	3.0 (1 – 5)	4.0 (1 – 5)					
Lymphoid Leukemia	108		5 (3 – 6)			7.0 (4 – 8)	3 (1 – 5)	3 (1 – 5)	3 (1 – 5)	5 (1 – 5)					
Plasma Cell Neoplasm (Multiple Myeloma)	109	7.0 (5 – 8)	2.0 (1 – 3)	3.5 (2 – 5)	6.0 (2 – 7)	8.0 (6 – 9)	7.0 (7 – 8)	4.5 (3 – 5)	3.5 (2 – 5)	4.5 (2 – 5)	5.0 (4 – 7)				
AML	110	6.0 (5 – 7)	2.0 (2 – 4)	3.5 (3 – 5)	8.5 (7 – 9)	9.0 (7 – 9)	9.0 (7 – 9)	3.5 (1 – 6)	3.5 (1 – 5)	4.0 (1 – 5)	5.0 (1 – 5)				
CML	111		7.0 (5 – 8)			8.0 (6 – 9)	8.0 (6 – 9)	3.0 (1 – 5)	2.0 (1 – 5)	3.0 (1 – 5)	4.0 (1 – 5)				
Myeloid Neoplasm	112		4.0 (3 – 7)			8.0 (7 – 9)	8.0 (7 – 9)	3.0 (1 – 5)	3.0 (1 – 4)	3.0 (1 – 5)	5.0 (1 – 5)				

[19] Curability is the receipt of effective treatment such that a population of individuals who are "cured" would have the same life expectancy as a population that never had the cancer being considered. Some members of each of those populations (cured and never had cancer) might be expected to die prematurely (e.g., from trauma), but many more would be expected to die of more typical age-related causes.

[20] Includes active treatment or surveillance.

[21] We define "early" as asymptomatic but already meeting the definition of cancer (e.g., not having a precursor condition).

[22] If there are many treatment advancements, cure rates may increase, and if treatment remains unchanged, cure rates are unlikely to change.

[23] A patient who is not cured could have increased life expectancy (e.g. still die of this cancer).

[24] Treatment consistent with NCCN guidelines.

[25] Please refer to the table above for examples of lower risk cancers.

[26] Please refer to the table above for examples of higher risk cancers.

[27] Acute lymphocytic (or lymphoblastic) leukemia.

[28] Chronic lymphocytic (or lymphoblastic) leukemia.

[29] Acute myeloid leukemia.

[30] Chronic myeloid leukemia.