

Supplementary Information

Effects of rifampin on the pharmacokinetics and pharmacodynamics of milvexian, a potent, selective, oral small molecule factor XIa inhibitor

Vidya Perera,^{1,*} Zhaoqing Wang,¹ Susan Lubin,¹ Lisa J. Christopher,¹ Wei Chen,¹ Sophia Xu,¹
Dietmar Seiffert,¹ Mary DeSouza,¹ Bindu Murthy¹

¹Bristol Myers Squibb, Princeton, NJ, USA.

Supplemental Table S1. PK and PD sampling schedule

Study day	Time (relative to milvexian dose); hour:min	PK blood sample	PD blood sample
1	00:00	X	X
1	00:30	X	X
1	1:00	X	
1	2:00	X	X
1	3:00	X	X
1	4:00	X	X
1	6:00	X	
1	8:00	X	X
1	10:00	X	
1	12:00	X	X
2	24:00	X	X
2	36:00	X	
3	48:00	X	X
4	00:00	X	X
4	00:30	X	
4	1:00	X	
4	2:00	X	
4	3:00	X	
4	4:00	X	
4	6:00	X	
4	8:00	X	
4	10:00	X	
4	12:00	X	

5	24:00	X	
5	36:00	X	
6	48:00	X	
7	72:00	X	
12	00:00	X	X
12	00:30	X	X
12	1:00	X	
12	2:00	X	X
12	3:00	X	X
12	4:00	X	X
12	6:00	X	
12	8:00	X	X
12	10:00	X	
12	12:00	X	X
13	24:00	X	X
13	36:00	X	
14	48:00	X	X
15	72:00	X	X

PK, pharmacokinetic; PD, pharmacodynamic.