

Supplemental material, table 1. Treatment and outcomes between patients with and without infarction.

	All (496)	Without infarction (353)	With infarction (143)	P-value
Anticoagulation alone	357 (72%)	259 (73%)	98 (69%)	0.545
CDT	70 (14%)	53 (15%)	17 (12%)	0.365
Thrombolysis				0.212
Half dose	25 (5%)	15 (4%)	10 (7%)	
Full dose	14 (3%)	8 (2%)	6 (4%)	
Mechanical thrombectomy	19 (4%)	10 (3%)	9 (6%)	0.069
Surgical thrombectomy	9 (2%)	6 (2%)	3 (2%)	0.763
ECMO	11 (2%)	6 (2%)	5 (4%)	0.218
Mechanical ventilation	48 (10%)	30 (9%)	18 (13%)	0.163
Antimicrobials	39 (8%)	19 (5%)	22 (15%)	<0.001
In-hospital death	37 (8%)	27 (8%)	10 (7%)	0.801
Disposition				0.759
Home	349 (76%)	245 (75%)	104 (78%)	
ECF	100 (22%)	74 (23%)	26 (20%)	
Hospice	10 (2%)	7 (2%)	3 (2%)	
Length of stay, days	10±13.0	9.5±12.3	10.7±14.7	0.698
Bleeding complication				0.089
Minor bleeding	31 (6%)	25 (7%)	6 (4%)	
Major bleeding	32 (7%)	18 (5%)	14 (10%)	
Need for transfusion	63 (13%)	41 (12%)	22 (16%)	0.253
Readmission within 30 days	76 (16.6%)	52 (16%)	24 (18%)	0.584
New O2 on discharge	75 (16%)	61 (19%)	14 (11%)	0.031
Pulmonary follow-up	199 (42%)	144 (44%)	55 (40%)	0.610

CDT= catheter-directed thrombolysis; ECMO= extracorporeal membrane oxygenation;  
ECF= extended care facility

Supplemental material, table 2. Outcomes and complication rates in CDT.

	All (70)	Without infarction (53)	With infarction (17)	P-value
In-hospital death	4 (6%)	3 (6%)	1 (6%)	0.973
Disposition				0.436
Home	57 (86%)	42 (84%)	15 (94%)	
ECF	9 (14%)	8 (16%)	1 (7%)	
Length of Stay	9±14.2	7±8.4	15±24.4	0.044
Readmission within 30 days	4 (6%)	1 (2%)	3 (18%)	0.047
New O2 on discharge	10 (16%)	8 (17%)	2 (12%)	0.609
Pulmonary follow-up	47 (72%)	38 (76%)	9 (56%)	0.166
<b>Complications</b>				
Bleeding				0.676
Minor	10 (14%)	8 (15%)	2 (12%)	
Major	5 (7%)	3 (6%)	2 (12%)	
RBC transfusion	10 (14%)	7 (13%)	3 (18%)	0.649
Access site hematoma	3 (5%)	1 (2%)	2 (13%)	0.134

ECF= extended care facility; RBC= red blood cell

