

Study or Subgroup	CRPS			Control			Weight	Std. Mean Difference		Year
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	IV, Random, 95% CI	
Sieweke 1999	44.2	4.12	23	44	2	15	7.2%	0.06	[-0.59, 0.71]	1999
Kemler 2000	40.3	24.18	33	46.5	13.71	50	11.2%	-0.33	[-0.77, 0.11]	2000
Kemler 2000	41.6	19.39	20	46.7	12.45	50	9.4%	-0.34	[-0.86, 0.18]	2000
Huge 2008	44.2	4.3	34	45.2	3	56	11.5%	-0.28	[-0.71, 0.15]	2008
Munts 2011	42.3	10.5	44	44.6	3	35	11.1%	-0.28	[-0.73, 0.16]	2011
Gierthmühlen 2012	41.4	4.7	298	44.7	2.7	32	13.1%	-0.72	[-1.09, -0.35]	2012
Kolb 2012	41.4	7.15	20	43.8	3.8	20	7.5%	-0.41	[-1.04, 0.22]	2012
Rooijen 2013	43.9	7.4	17	44	5.2	42	8.6%	-0.02	[-0.58, 0.55]	2013
Rooijen 2013	41.77	9.64	31	44	5.2	42	10.6%	-0.30	[-0.76, 0.17]	2013
Terkelsen 2014	38.4	4.1	20	44.3	3.3	20	6.3%	-1.55	[-2.27, -0.84]	2014
Rasmussen 2018	35.6	3.8	8	38.4	2.8	8	3.6%	-0.79	[-1.82, 0.24]	2018
<b>Total (95% CI)</b>			<b>548</b>			<b>370</b>	<b>100.0%</b>	<b>-0.41</b>	<b>[-0.63, -0.20]</b>	

Heterogeneity: Tau<sup>2</sup> = 0.06; Chi<sup>2</sup> = 18.05, df = 10 (P = 0.05); I<sup>2</sup> = 45%

Test for overall effect: Z = 3.78 (P = 0.0002)

