

Study or Subgroup	CRPS			Control			Weight	Std. Mean Difference		Year
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	IV, Random, 95% CI	
Kemler 2000	31.3	1.76	33	31.3	1.44	50	11.9%	0.00	[-0.44, 0.44]	2000
Kemler 2000	30	12.2	20	30.9	5.1	50	10.6%	-0.11	[-0.63, 0.40]	2000
Huge 2008	29.2	1.6	34	30.5	0.9	56	11.7%	-1.06	[-1.52, -0.61]	2008
Munts 2011	31.1	1	7	31.5	0.6	35	6.6%	-0.58	[-1.40, 0.24]	2011
Kolb 2012	29.5	1.7	20	29.5	1.4	20	9.1%	0.00	[-0.62, 0.62]	2012
Rooijen 2013	30.4	1.35	21	30.6	0.7	42	10.5%	-0.21	[-0.73, 0.32]	2013
Rooijen 2013	28.8	2.4	30	28.2	2.4	42	11.4%	0.25	[-0.22, 0.72]	2013
Rooijen 2013	30.6	0.5	48	31	0.8	42	12.2%	-0.60	[-1.03, -0.18]	2013
Rooijen 2013	27.4	4.6	24	28.9	2.5	42	10.8%	-0.44	[-0.94, 0.07]	2013
Rasmussen 2018	28.5	3.8	8	30	1.1	8	5.1%	-0.51	[-1.51, 0.49]	2018
<b>Total (95% CI)</b>			<b>245</b>			<b>387</b>	<b>100.0%</b>	<b>-0.32</b>	<b>[-0.58, -0.05]</b>	

Heterogeneity: Tau<sup>2</sup> = 0.11; Chi<sup>2</sup> = 22.16, df = 9 (P = 0.008); I<sup>2</sup> = 59%

Test for overall effect: Z = 2.30 (P = 0.02)

