


Study or Subgroup	CRPS			Control			Weight	Std. Mean Difference		Year
	Mean	SD	Total	Mean	SD	Total		IV, Random, 95% CI	95% CI	
Kemler 2000	33.1	5.7	33	32.5	2.89	50	12.8%	0.14	[-0.30, 0.58]	2000
Kemler 2000	36.5	19	20	35.4	20.2	50	11.2%	0.05	[-0.46, 0.57]	2000
Huge 2008	37.2	4.1	34	34.3	1.5	56	12.5%	1.04	[0.58, 1.49]	2008
Munts 2011	34.7	3.7	44	32.5	0.3	35	12.4%	0.79	[0.32, 1.25]	2011
Kolb 2012	35.7	4	20	35.5	2.6	20	9.4%	0.06	[-0.56, 0.68]	2012
Gierthmühlen 2012	32.49	0.32	298	32.35	0.2	32	14.5%	0.45	[0.08, 0.82]	2012
Rooijen 2013 (1)	38.3	6.4	17	35	2	42	10.0%	0.86	[0.27, 1.44]	2013
Rooijen 2013	38.2	6.84	31	35	2	42	12.0%	0.67	[0.20, 1.15]	2013
Rasmussen 2018	36.3	1.7	8	37	3.4	8	5.1%	-0.25	[-1.23, 0.74]	2018
<b>Total (95% CI)</b>			<b>505</b>			<b>335</b>	<b>100.0%</b>	<b>0.48</b>	<b>[0.22, 0.73]</b>	

Heterogeneity: Tau<sup>2</sup> = 0.08; Chi<sup>2</sup> = 18.41, df = 8 (P = 0.02); I<sup>2</sup> = 57%  
Test for overall effect: Z = 3.62 (P = 0.0003)



Footnotes

(1) 6