

Supplementary Table. Changes in Symptom Score of Korean Gastrointestinal Symptom Rating Scale in Each Group

KGSRS scores	Tegoprazan 50 mg (n = 22)				Esomeprazole 40 mg (n = 24)			
	Baseline	Week 2	Change	<i>P</i> -value ^a	Baseline	Week 2	Change	<i>P</i> -value ^a
Reflux	33.8 ± 27.0	14.5 ± 19.3	-19.3 ± 24.5	0.0004	31.5 ± 21.5	19.8 ± 18.9	-11.7 ± 19.7	0.0107
Abdominal pain	27.5 ± 23.7	12.7 ± 21.2	-14.8 ± 14.9	0.0001	18.9 ± 19.4	13.2 ± 14.3	-5.7 ± 17.0	0.1129
Indigestion	32.2 ± 27.6	23.3 ± 21.3	-8.8 ± 14.8	0.0078	31.1 ± 18.8	24.2 ± 20.5	-6.9 ± 15.2	0.0033
Diarrhea	17.2 ± 20.4	10.2 ± 15.9	-7.0 ± 11.4	0.0073	9.4 ± 14.5	8.5 ± 12.3	-0.9 ± 15.2	0.9683
Constipation	14.6 ± 23.2	12.7 ± 25.8	-1.9 ± 14.2	0.1367	15.5 ± 19.9	13.4 ± 18.5	-2.2 ± 16.6	0.2095

^aPaired *t* test or Wilcoxon signed rank test.

KGSRS, Korean gastrointestinal symptom rating scale; Change = Week 2 – Baseline.

Data are presented as mean ± SD.