

Breastfeeding is not a risk factor for clinical severity in Autism Spectrum Disorder in children from the ELENA cohort

Scientific Report

Marianne Peries^{1,2,3}, Fanny Duhr¹, Marie-Christine Picot^{4,3,2}, Barbara Heude⁵, Jonathan Y. Bernard^{5,6}, Amaria Baghdadli^{1,2,3,7}

¹ Centre de Ressource Autisme Languedoc-Roussillon, CHU Montpellier, Montpellier, France

² Centre d'Excellence sur l'Autisme et les Troubles Neuro-développementaux, CHU Montpellier, France

³ Université Paris-Saclay, UVSQ, Inserm, CESP, Team DevPsy, 94807, Villejuif, France

⁴ Clinical research and epidemiology unit, Department of Medical Information, University Hospital, CHU Montpellier, France

⁵ Université Paris Cité, Inserm, INRAE, Centre de Recherche en Épidémiologie et Statistiques, F-7005 Paris, France

⁶ Singapore Institute for Clinical Sciences (SICS), Agency for Science, Technology and Research (A*STAR), Singapore, Singapore

⁷ Faculté de Médecine, Université de Montpellier, France

Corresponding author: Amaria BAGHDADLI, Centre de Ressource Autisme Languedoc-Roussillon et Centre d'Excellence sur l'Autisme et les Troubles Neuro-développementaux (CeAND), 39 Avenue Charles Flahaut, 34295 Montpellier cedex 05, France. Tel: +33467330986 ; Fax: +33467330832 ; E-mail address: rech-clinique-autisme@chu-montpellier.fr

Supplementary Table S1. Description of missing data for each confounders included in the multiple imputation models

Confounders	Missing value n (%)
Mothers' age (years)	0
Mother's tobacco smoking during pregnancy	3 (1.2)
Mother's obesity before pregnancy	8 (3.3)
Child's sex	0
Gestational age (weeks)	16 (6.6)
Parental education level	1 (0.41)
Number of older siblings	4 (1.64)
Parental socioeconomic status (SES)	2 (0.8)

Supplementary Table S2. Comparison of clinical characteristics at inclusion between participants considered in the analyses and the other participants involved in the ELENA Cohort

	Participants included in our study (n = 243)	Other participants included in ELENA Cohort (n = 633)	p value
Children characteristics at assessment			
Age (years)	5.8 (± 3.4)	6.0 (± 3.3)	0.34
Sex			
Boy	199 (81.9%)	525 (82.9%)	0.72
Girl	44 (18.1%)	108 (17.1%)	
Intellectual quotient	77.1 (± 28.0)	71.3 (± 25.7)	0.01
VABS-II standard scores			
Communication SS	72.6 (± 15.5)	68.9 (± 15.2) ^a	0.001
Daily living skills SS	75.6 (± 12.8)	72.5 (± 12.7)	0.001
Social skills SS	70.9 (± 10.6)	69.5 (± 10.8) ^a	0.20
Level of autism spectrum (ADOS-2 CSS)			
No evidence to low related symptoms	29 (13.7%) ^b	61 (10.8%) ^c	0.47
Moderate related symptoms	100 (47.1%)	263 (46.6%)	
High related symptoms	83 (39.2%)	240 (42.6%)	
ABC			
Irritability / aggressiveness	31.1 (17.8 - 46.7) ^d	31.1 (17.8 - 47.8) ^g	0.55
Lethargy / social withdrawal	25.0 (12.5 - 35.4) ^e	25.0 (14.6 - 37.5) ^h	0.56
Stereotyped behaviors / self-harm	23.8 (9.5 - 47.6) ^e	28.6 (14.3 - 52.4) ^h	0.10
Hyperactivity / lack of cooperation	43.8 (20.8 - 64.6) ^e	43.6 (27.1 - 60.4) ⁱ	0.88
SRS-2 T-score	94.9 (± 19.0) ^f	93.9 (± 18.3) ^j	0.62
Parents characteristics			
Age at birth (years)			
Mother	31.8 (± 5.4)	31.2 (± 5.4)	0.24
Father	35.3 (± 7.3)	34.2 (± 6.4)	0.16
Mother's education level			
High school or lower	66 (34.4%)	145 (41.4%)	0.11
University	126 (65.5%)	205 (58.6%)	
Father's education level			
High school or lower	91 (47.6%)	168 (49.1%)	0.74
University	100 (52.4%)	174 (50.9%)	

Data are presented as the mean ($\pm SD$), median (IQR), or n (%).

^a 1 missing value, ^b 31 missing values, ^c 69 missing values, ^d 59 missing values, ^e 58 missing values, ^f 95 missing values, ^g 321 missing values, ^h 323 missing values, ⁱ 320 missing values, ^j 410 missing values

ADOS-2 CSS, autism diagnostic observation schedule second version calibrate severity scale; VABS-II, Vineland second version; ABC, Aberrant Behavior Checklist; SRS-2, Social-Responsiveness Scale, second version.

Supplementary Table S3. Association between predominant and any breastfeeding in tertiles and ASD clinical severity at inclusion

Predominant Breastfeeding				Any Breastfeeding				
	Low tertile (<2 months)	Middle tertile (2-4.5 months)	High tertile (≥4.5 months)	P value	Low tertile (<3 months)	Middle tertile (3-8 months)	High tertile (≥8months)	P value
Intellectual quotient	(N = 60)	(N = 76)	(N = 41)		(N = 62)	(N = 61)	(N = 54)	
<i>IQ</i>	74.0 (±24.4)	82.2 (±30.7)	77.2 (±27.1)	0.23	77.5 (±24.4)	76.2 (±30.0)	81.4 (±29.6)	0.60
VABS-II standard scores	(N = 60)	(N = 76)	(N = 41)		(N = 62)	(N = 61)	(N = 54)	
<i>Communication</i>	73.8 (±14.0)	74.8 (±14.3)	71.5 (±18.4)	0.53	74.5 (±14.1)	71.8 (±14.9)	74.9 (±16.8)	0.47
<i>Daily living skills</i>	77.9 (±11.7)	76.9 (±13.0)	76.4 (±13.9)	0.82	76.9 (±11.8)	77.4 (±13.2)	77.1 (±13.5)	0.98
<i>Socialization</i>	73.0 (±10.4)	70.1 (±11.3)	71.4 (±12.1)	0.33	71.3 (±11.3)	71.23 (±11.4)	71.54 (±11.1)	0.99
ADOS-2 CSS	(N = 52)	(N = 70)	(N = 35)		(N = 53)	(N = 56)	(N = 48)	
	7.0 (±1.5)	6.6 (±2.3)	6.9 (±1.8)	0.75	6.8 (±1.9)	7.2 (±1.9)	6.4 (±2.0)	0.09
ABC	(N = 43)	(N = 63)	(N = 30)		(N = 47)	(N = 48)	(N = 41)	
<i>Irritability/ aggressiveness</i>	26.7 (11.1 - 40.0)	33.3 (20.0 - 51.1) ^a	37.8 (20.0 - 44.4)	0.14	26.7 (13.3 - 46.7)	31.1 (13.3 - 42.2) ^a	37.8 (20.0 - 48.9)	0.30
<i>Lethargy/ social withdrawal</i>	22.9 (12.5 - 35.4)	22.9 (10.4 - 41.7)	27.1 (12.5 - 35.4)	0.85	22.9 (12.5 - 43.8)	22.9 (10.4 - 38.5)	25.0 (12.5 - 31.3)	0.91
<i>Stereotyped behaviors/ self-harm</i>	19.1 (4.8 - 42.9)	28.6 (9.5 - 47.6)	23.8 (9.5 - 52.4)	0.28	28.6 (9.5 - 47.6)	26.2 (9.5 - 47.6)	23.8 (9.5 - 42.9)	0.99
<i>Hyperactivity/ lack of cooperation</i>	35.4 (20.8 - 62.5)	45.8 (20.8 - 68.8)	50.0 (37.5 - 64.6)	0.23	37.5 (22.9 - 64.6)	43.8 (22.9 - 65.6)	50.0 (31.3 - 66.7)	0.63
SRS-2 T-score	(N = 36)	(N = 51)	(N = 23)		(N = 40)	(N = 39)	(N = 31)	
	93.3 (±18.1)	93.8 (±17.9)	99.7 (±21.7)	0.38	93.3 (±18.6)	98.7 (±18.7)	91.9 (±19.1)	0.26

Data are presented as the mean (±SD) or median (IQR).

^a 1 missing value

VABS-II, Vineland second version; ADOS-2 CSS, autism diagnostic observation schedule second version calibrate severity scale; ABC, Aberrant Behavior Checklist; SRS-2, Social-Responsiveness Scale, second version.