

Supporting Information

Laser-induced Combinatorial Chemotherapeutic, Chemodynamic, and Photothermal Therapy for Hepatocellular Carcinoma based on Oxaliplatin-loaded Metal-Organic Frameworks

Run Huang^{1,2}, Weiren Liu^{1,2}, Qinghao Zhang³, Guiqi Zhu^{1,2}, Weifeng Qu^{1,2}, Chenyang Tao^{1,2}, Jun Gao^{1,2}, Yuan Fang^{1,2}, Xiutao Fu^{1,2}, Jian Zhou^{1,2}, Yinghong Shi^{1,2}, Jia Fan^{1,2*}, Zheng Tang^{1,2*}*




Figure S1, Representative TEM images of degradation of MOF after incubation for 3h in PBS with pH 6.5 (A) and 5.6 (B).




Figure S2, Heating and cooling curves of MIL-PDA-PEGTK under laser irradiation (808 nm, 1 W/cm⁻²).

•




Figure S3, *In vitro* catalytic performance of nanoparticles. Chromogenic reaction of TMB under different nanoparticles concentrations (1 mM H₂O₂, pH = 5.6).




Figure S4, Histopathological images of main organs at the end of treatment.




Figure S5, Mouse blood serum analysis testing the biochemical markers of liver function, kidney function, and heart function.