

Supplementary information

Structural basis of lipopolysaccharide maturation by the O-antigen ligase

In the format provided by the authors and unedited

Structural basis of Lipopolysaccharide Maturation by O-Antigen Ligase WaaL

Khuram U. Ashraf¹, Rie Nygaard¹, Owen N. Vickery^{2,3,#}, Satchal K. Erramilli^{4,#}, Carmen M. Herrera^{5,#}, Thomas H. McConville^{6,#}, Vasileios I. Petrou^{7,8}, Sabrina I. Giacometti¹, Meagan Belcher Dufresne¹, Kamil Nosol⁴, Allen P. Zinkle¹, Chris L.B. Graham², Michael Loukeris⁹, Brian Kloss⁹, Karolina Skorupinska-Tudek¹⁰, Ewa Swiezewska¹⁰, David I. Roper^{2,1}, Oliver B. Clarke^{1,11}, Anne-Catrin Uhlemann⁶, Anthony A. Kossiakoff⁴, M. Stephen Trent^{5*}, Phillip J. Stansfeld^{2,3*} and Filippo Mancia^{1*}

¹Department of Physiology and Cellular Biophysics, Columbia University Irving Medical Center, New York, NY 10032, USA.

²School of Life Sciences, University of Warwick, Coventry, CV4 7AL, UK.

³Department of Chemistry, University of Warwick, Coventry, CV4 7AL, UK.

⁴Department of Biochemistry and Molecular Biology, University of Chicago, Chicago, IL, 60637, USA.

⁵Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia, Athens, GA, 30602, USA.

⁶Department of Medicine, Division of Infectious Diseases, Columbia University Medical Center, New York, New York, 10032, USA.

⁷Department of Microbiology, Biochemistry, and Molecular Genetics, New Jersey Medical School, Rutgers Biomedical Health Sciences, Newark, NJ, 07103, USA.

⁸Center for Immunity and Inflammation, New Jersey Medical School, Rutgers Biomedical Health Sciences, Newark, NJ, 07103, USA.

⁹New York Consortium on Membrane Protein Structure, New York Structural Biology Center, 89 Convent Avenue, New York, NY, 10027, USA.

¹⁰Institute of Biochemistry and Biophysics, Polish Academy of Sciences, 02-106 Warsaw, Poland.

¹¹Department of Anesthesiology, Columbia University Irving Medical Center, New York, NY, 10032, USA.

#These authors contributed equally.

*Correspondence to be addressed to: strent@uga.edu (M.S.T.),

Phillip.Stansfeld@warwick.ac.uk (P.J.S.), fm123@cumc.columbia.edu (F.M.)

Table of Content:

***Red, dashed, boxes indicate how gels where cropped.**


SI Figure 1 | Uncropped LPS assay gels from Main Figures. a, LPS assay gel from [Fig. 2c](#) showing analysis of *CmWaaL* ligase activity in whole cells. LPS gel showing O-antigen extension in *C. metallidurans* expressing *CmWaaL* variants. With and empty plasmid as a control. **b,** Functional analysis gel from [Fig. 4c](#) of *EcWaaL* ligase variants in whole cells by LPS gel analysis. WaaL proteins were expressed from plasmid pWSK29 in the W3110 Δ waaL strain. With and empty plasmid as a control.

SI Figure 2 | Uncropped LPS assay gels from Extended Data Figures. a, LPS assay gel from [Extended Data Fig. 1c](#) showing functional analysis of *CmWaaL* ligase activity in whole cells, showing O-antigen ligase activity is abolished when the gene encoding WaaL is deleted and activity is restored by plasmid complementation. **b,** Functional analysis of *EcWaaL* ligase activity from [Extended Data Fig. 7a](#) carried out in whole cells by LPS gel analysis. W3110 Δ waaL containing either empty vector pWSK29 (\emptyset), pWSK29::*EcWaaL* (WT) as controls and pWSK19::*EcWaaL*- mutant variants to test for O-antigen ligase activity. **c,** Functional analysis gel from [Extended Data Fig. 7e](#) showing *CmWaaL* ligase activity in whole cells. *C. metallidurans* Δ waaL containing either empty vector pBBR1(\emptyset), pBBR1:*CmWaaL* (WT) as controls for evaluating *CmWaaL*-variants for O-antigen extension.


SI Figure 3 | Uncropped SDS-page gels from Extended Data Figures. a, SDS-PAGE gel from [Extended Data Fig. 1e](#) depicting *CmWaaL* purified in detergent, *CmWaaL* reconstituted into nanodiscs (MSP1E3D1 and POPG), and *CmWaaL* reconstituted into nanodiscs (MSP1E3D1 and POPG) with Fab (WaB10) bound. Molecular weight markers can also be observed and are labelled. **b,** SDS-PAGE gels from [Extended Data Fig. 7c](#) of all *EcWaaL* mutants used for LPS ligase analysis, that were purified to verify expression.

SI Figure 4 | Uncropped TLC and Western Blot analysis from Extended Data Figures. a, TLC analysis from Extended Data Fig. 5d of detergent purified CmWaaL along with suitable controls: POPG, Und-P, Und-PP and an unrelated control protein expressed in *E. coli* were run in separate lanes. **b,** Western blot analysis, from Extended Data Fig. 7d, using a mouse monoclonal anti-Flag antibody, of Flag purified WT *CmWaaL* and mutants, grown in *C. metallidurans*, for testing in the O-antigen ligase assay. Red markers indicate molecular weight markers.


Supplementary Table 1 | Raw Single-point ELISA and Multi-point sAB ELISA data




Supplementary Figure 1a | Uncropped gel from Figure 2c.




Supplementary Figure 1b | Uncropped gel from Figure 4c.




Supplementary Figure 2a | Uncropped gel from Extended Data Fig. 1c





Supplementary Figure 2b | Uncropped gels from Extended Data Fig. 7a



Supplementary Figure 2c | Uncropped gels from Extended Data Fig. 7e




Supplementary Figure 3a | Uncropped gel from Extended Data Fig. 1e



Supplementary Figure 3b | Uncropped gel from Extended Data Fig. 7c



Supplementary Figure 4a | Uncropped gel from Extended Data Fig. 5d



Supplementary Figure 4b | Uncropped gel from Extended Data Fig. 7d