

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

BPDE, the Migration and Invasion of Human Trophoblast Cells, and Occurrence of Miscarriage in Humans: Roles of a Novel *lncRNA-HZ09*

Mengyuan Dai, Wenxin Huang, Xinying Huang, Chenglong Ma, Rong Wang, Peng Tian, Weina Chen, Ying Zhang, Chenyang Mi, and Huidong Zhang

Table of Contents

Table S1. DNA sequences used for construction of pcDNA3.1 overexpression plasmids.

Table S2. RNA sequences used for RNA transfection.

Table S3. PCR protocols for RACE assays.

Table S4. Primer sequences used for RACE assays.

Table S5. The full-length sequence of *lnc-HZ09*.

Table S6. RT-qPCR programs.

Table S7. Primer sequences used for RT-qPCR analysis (Sangon Biotech, Shanghai, China).

Table S8. Primer sequences used in various RT-qPCR assays (Sangon Biotech, Shanghai, China).

Table S9. DNA sequence in pGEM-T used for transcription of antisense sequence of *lnc-HZ09* or PLD1 mRNA.

Table S10. Protein-coding potential of *lnc-HZ09* predicted by CPAT software.

Table S11. m6A modification site on *lnc-HZ09* identified by SRAMP software.

Table S12. Univariate analysis of the parameters of the RM and HC groups (each n = 15).

Table S13. Sequence conservation among various species.

Figure S1. Identification of *lnc-HZ09* and overexpression or knockdown of *lnc-HZ09* in human trophoblast cells.

Figure S2. Expression levels of nine mRNAs with knockdown of *lnc-HZ09* and the functions of PLD1 in human trophoblast cells.

Figure S3. Expression levels of members of the PLD1/RAC1/CDC42 pathway regulated by *lnc-HZ09* in human trophoblast cells.

Figure S4. Transcription factors of PLD1 predicted by PROMO software.

Figure S5. PLD1 mRNA stability regulated by *lnc-HZ09* in human trophoblast cells.

Figure S6. Transcription factors of *lnc-HZ09* predicted by PROMO software.

Figure S7. METTL3-mediated m6A RNA methylation on *lnc-HZ09* regulated PLD1 expression level and migration and invasion of human trophoblast cells.

Figure S8. Expression levels of members of PLD1/RAC1/CDC42 pathway regulated by BPDE or *lnc-HZ09* in BPDE-exposed human trophoblast cells.

Figure S9. PLD1 and *lnc-HZ09* RNA stability in BPDE-exposed human trophoblast cells.

Figure S10. The BPDE-DNA adduct levels and the correlation among various RNAs and proteins in HC and RM villous tissues.

Figure S11. Sequence alignment of various genes in different species and the protein levels of Sp1, Pld1, Rac1, and Cdc42 in BaP-treated mice.

References

Additional File- Excel Document