

Supplementary Material

Development of a Nomogram for Predicting Asymptomatic Coronary Artery Disease in Patients with Ischemic Stroke

Jie Yang^{1,2}, Xinguang Yang^{1,2}, Jun Wen³, Jiayi Huang^{1,2,4}, Lihong Jiang^{1,2}, Sha Liao^{1,2}, Chun Lian^{1,2}, Haiyan Yao^{1,2}, Li Huang^{1,2}, Youming Long^{1,2,*}.

¹Department of Neurology, The Second Affiliated Hospital of GuangZhou Medical University, 250# Changgang East Road, GuangZhou, 510260, Guangdong Province, China; ²Key Laboratory of Neurogenetics and Channelopathies of Guangdong Province and The Ministry of Education of China, Institute of Neuroscience and the Second Affiliated Hospital of GuangZhou Medical University, 250# Changgang east Road, GuangZhou, 510260, Guangdong Province, China; ³Department of Neurology, Jiangmen Central Hospital, 23# Haibang street, North Street, Jiangmen, 529000, Guangdong Province, China; ⁴Department of Neurology, Dongguan Dongcheng Hospital, 56# Nancheng Road, DongGuan, 523000, Guangdong Province, China

Supplementary Table 1

Stepwise regression model for variables selection.

Variable	Estimate	Std Error	z-value	p-value	AIC
(Intercept)	-3.91	1.378	-2.837	0.005	322.2
Male	0.682	0.366	1.861	0.063	
Age	0.029	0.015	1.885	0.059	
Hypertension	1.07	0.404	2.647	0.008	
Blood glucose level	0.134	0.059	2.251	0.024	
HDL-C	-1.146	0.653	-1.756	0.079	
Presence of CVA \geq 50% stenosis	0.513	0.286	1.795	0.073	

HDL-C, high-density lipoprotein cholesterol; CVA, cervicocephalic artery; AIC, Akaike Information Criterion.