

Supplemental Tables

Supplemental Table 1. Comparison of Rates of cpRNFLT Thinning between DH and Non-DH Eyes in Glaucoma Suspects and Early Stage of POAG

DH Group	Non-DH Group	Difference	P value
Mean (95% CI)	Mean (95% CI)	Mean (95% CI)	(adjusted)
cpRNFL Change Rate ($\mu\text{m/year}$)			
Mean Global	-0.48 (-0.63, -0.34)	-0.01 (-0.33, 0.30)	0.935 (0.904)
Nasal	-0.10 (-0.48, 0.27)	0.29 (-0.12, 0.71)	0.165 (0.283)
Temporal	-0.12 (-0.40, 0.16)	-0.16 (-0.78, 0.45)	0.601 (0.565)
Inferonasal	-0.98 (-1.27, -0.70)	0.31 (-0.32, 0.94)	0.333 (0.243)
Inferotemporal	-0.78 (-1.09, -0.47)	-1.00 (-1.70, -0.31)	0.005 (0.013)
Superonasal	-0.55 (-0.84, -0.26)	0.24 (-0.41, 0.88)	0.472 (0.726)
Superotemporal	-0.58 (-0.86, -0.29)	0.29 (-0.35, 0.93)	0.370 (0.303)

cp = circumpapillary; DH = disc hemorrhage; RNFL = retinal nerve fiber layer. Values are shown in mean (95% confidence interval), unless otherwise indicated. Statistically significant P values are shown in bold.

*Adjusted to age, gender, baseline 24-2 visual field mean deviation, and mean IOP.