

Supplemental Table 2. Comparison of Rates of CD Loss between DH and Non-DH Eyes in Glaucoma Suspects and Early Stage of POAG

	DH Group Mean (95% CI)	Non-DH Group Mean (95% CI)	Difference Mean (95% CI)	P value (adjusted)
CD Change Rate (%/year)				
wiCD	-0.91 (-1.19, -0.63)	-0.58 (-0.73, -0.44)	-0.32 (-0.64, -0.01)	0.042 (0.038)
cpCD	-0.83 (-1.15, -0.51)	-0.50 (-0.66, -0.34)	-0.33 (-0.69, 0.02)	0.066 (0.078)
Nasal	-1.06 (-1.47, -0.65)	-0.58 (-0.78, -0.37)	-0.48 (-0.94, -0.03)	0.038 (0.015)
Temporal	-0.55 (-0.94, -0.16)	-0.61 (-0.80, -0.41)	0.06 (-0.38, 0.50)	0.799 (0.709)
Inferonasal	-0.86 (-1.36, -0.35)	-0.26 (-0.51, 0.00)	-0.60 (-1.17, -0.03)	0.038 (0.067)
Inferotemporal	-1.36 (-1.90, -0.83)	-0.43 (-0.70, -0.17)	-0.93 (-1.53, -0.34)	0.002 (0.006)
Superonasal	-0.56 (-1.06, -0.07)	-0.33 (-0.57, -0.08)	-0.24 (-0.79, 0.31)	0.401 (0.707)
Superotemporal	-0.73 (-1.28, -0.18)	-0.51 (-0.79, -0.23)	-0.22 (-0.84, 0.40)	0.488 (0.525)

CD = capillary density; cp = circumpapillary; DH = disc hemorrhage; IOP = intraocular pressure; M = male; wi = whole image. Values are shown in mean (95% confidence interval), unless otherwise indicated. Statistically significant P values are shown in bold.

*Adjusted to age, gender, baseline 24-2 visual field mean deviation, and mean IOP.