

# **QEEG Signatures are Associated with Nonmotor Dysfunctions in Parkinson's Disease and Atypical Parkinsonism: An Integrative Analysis**

**Hailing Liu<sup>1, 2#</sup>, Zifeng Huang<sup>1#</sup>, Bin Deng<sup>1#</sup>, Zihan Chang<sup>1#</sup>, Xiaohua Yang<sup>1</sup>, Xingfang Guo<sup>1</sup>, Feilan Yuan<sup>1</sup>, Qin Yang<sup>1</sup>, Liming Wang<sup>3</sup>, Haiqiang Zou<sup>4</sup>, Mengyan Li<sup>5</sup>, Zhaohua Zhu<sup>6</sup>, Kunlin Jin<sup>7\*</sup>, Qing Wang<sup>1\*</sup>**

# SUPPLEMENTARY DATA

**Supplementary Table 1.** Correlation analysis of EEG indices and all variables in PD, PSP and MSA patients.

Variable	Frontal slow-to-fast ratio				Occipital alpha/theta ratio			
	PD		PSP		PD		MSA	
	r	p	r	p	r	p	r	p
H&Y stage	-0.034	0.812	0.154	0.493	0.111	0.438	0.116	0.533
UPDRS-III	-0.039	0.785	0.242	0.278	0.172	0.229	0.183	0.324
MMSE	-0.308	0.026*	-0.451	0.035*	0.005	0.973	-0.001	0.997
NMSS (total)	-0.077	0.588	0.299	0.176	0.156	0.268	-0.094	0.615
Cardiovascular	-0.189	0.180	-0.169	0.452	0.221	0.115	-0.216	0.243
Sleep/fatigue	-0.233	0.096	0.267	0.229	0.090	0.527	0.007	0.969
Mood/apathy	-0.049	0.732	0.434	0.043*	0.141	0.319	0.220	0.235
Perception/illusion	-0.060	0.670	0.465	0.029*	0.087	0.541	0.250	0.175
Attention/memory	-0.189	0.179	0.107	0.634	0.173	0.221	0.078	0.675
Gastrointestinal	0.033	0.816	-0.127	0.572	0.118	0.403	0.117	0.532
Urinary	-0.175	0.214	0.077	0.734	0.051	0.718	-0.161	0.386
Sexual function	-0.187	0.183	0.010	0.966	0.093	0.511	-0.082	0.662
Miscellaneous	-0.217	0.123	0.083	0.081	0.039	0.783	0.225	0.225
HAMA scores	-0.111	0.435	0.108	0.631	0.246	0.082	0.396	0.027*
HAMD scores	0.122	0.391	0.325	0.140	0.070	0.627	0.267	0.147

Abbreviations: r: Spearman's rank correlation coefficient, \*P < 0.05