Variable		Not Seen	Seen	P Value
Race	White	5 (33.3%)	20 (29.9%)	0.766
	Black	10 (66.7%)	47 (70.1%)	
Age	Q1	2 (13%)	17 (25.3%)	0.408
	Q2	2 (13%)	16 (23.8%)	
	Q3	4 (27%)	17 (25.3%)	
	Q4	7 (47%)	17 (25.3%)	
Language	English	10 (66.7%)	51 (76.1%)	0.516
	Non-English	5 (33.3%)	16 (23.94%)	
Organ Site	Ovary	10 (66.7%)	21 (31.3%)	0.033
	Uterus	1 (6.7%)	21 (31.3%)	
	Cervix / Vulva	4 (26.7%)	25 (37.3%)	
Country of Birth	USA Born	0 (0.0%)	15 (22.4%)	0.104
	Foreign	4 (26.7%)	12 (17.9%)	
	Unknown	11 (73.3%)	40 (59.7%)	
Insurance	Private	7 (46.7%)	47 (70.1%)	0.114
	Other	8 (53.3%)	20 (29.9%)	
Stage at Diagnosis	1	1 (6.7%)	6 (9.0%)	0.545
	II	1 (6.7%)	12 (17.9%)	
		9 (60.0%)	26 (38.8%)	
	IV	4 (26.7%)	23 (34.3%)	
leason for Referral	Symptom Management	11 (73.3%)	49 (73.1%)	0.222
	Other	4 (26.7%)	18 (26.9%)	

Supplementary Table. Variations in patient demographics by completion of palliative care consult.