

Supplemental Table 3: Adverse maternal outcomes, stratified by MRP and antenatal risk categories

	Midwife MRP	Family Physician MPR	Obstetrician MPR
Low risk (0-2)			
Blood transfusion	123/46,632 (0.3%)	603/132,309 (0.5%)	481/76,694 (0.6%)
ICU admissions	7/46,632 (0%)	37/132,309 (0%)	39/76,694 (0.1%)
Uterine Rupture*	5/43,276 (0%)	10/116,138 (0%)	<5 cases/44,218
Postpartum wound infection	21/46,632 (0%)	148/132,309 (0.1%)	143/76,694 (0.2%)
Moderate risk (3-6)			
Blood transfusion	94/15,437 (0.6%)	491/51,519 (1.0%)	797/77,142 (1.0%)
ICU admissions	<5 cases/15,437	53/51,519 (0.1%)	93/77,142 (0.1%)
Uterine Rupture*	<5 cases/13,041	12/38,181 (0%)	6/24,235 (0%)
Postpartum wound infection	14/15,437 (0.1%)	176/51,519 (0.3%)	246/77,142 (0.3%)
High risk (≥ 7)			
Blood transfusion	19/1082 (1.8%)	90/5851 (1.5%)	501/18,390 (2.7%)
ICU admissions	<5 cases/1082	15/5851 (0.3%)	87/18,390 (0.5%)
Uterine Rupture*	<5 cases/671	<5 cases/3003	<5 cases/3607
Postpartum wound infection	61,082 (0.6%)	69/5851 (1.2%)	100/18,390 (0.5%)

*Analysis restricted to those with vaginal birth