

Appendices to “Oral iron supplementation and anemia in children according to schedule, duration, dose, and co-supplementation: a systematic review and meta-analysis of 129 randomized trials”**Table of Contents**

Appendix 1: Search terms.	2
Appendix 2: Variables extracted from eligible studies.	3
Appendix 3. Elemental iron supplementation dose tertile ranges.	7
Appendix 4. References to included studies.	8
Appendix 5. Characteristics of included studies.	16
Appendix 6. Modification of iron effects after control for baseline anemia.	36
Appendix 7. Effect modification of iron by baseline anemia, child age, child sex, WHO region, and iron formulation.	37
Appendix 8. Comparison of iron supplementation effects within factorial trials, using double placebo as a common referent group.	40
Appendix 9. Effect heterogeneity p-values for anthropometric, infectious, and development outcomes.*†	41
Appendix 10. Risk of bias within included studies.	42
Appendix 11. Cochrane risk of bias assessment of included studies (n=129).	45
Appendix 12. Effect of oral iron supplementation versus placebo among children and adolescents aged <20 years among trials judged to not be at “high” risk of bias.	46
Appendix 13. Assessment of small study bias for outcomes reported by ≥10 groups randomized to iron.	47

Appendix 1: Search terms.

Title and abstract searches of the terms below were performed in PubMed, Scopus, Web of Science, and the Cochrane Central Register of Controlled Trials from inception through November 2020.

Infant OR Infan* OR newborn* OR new-born* OR perinat* OR neonat* OR child OR child OR child* OR children* OR schoolchild* OR schoolchild OR "school child" OR adolescent OR adolescen* OR youth* OR teen* OR pubescen* OR pediatrics OR pediatric* OR paediatric* OR peadiatric* OR school OR school* OR prematur* OR preterm* OR "Pregnant Women" OR Pregnancy OR pregnan* OR gravid OR obstetric OR antenatal OR antepartum OR gestation* OR lactation OR "breast feeding" OR lactation OR "breast feeding"[Title/Abstract] OR breast-fe* OR breastfe* OR breast-milk OR breastmilk OR "lactating mother*" OR "lactating woman" OR "lactating women"

AND

iron OR hematinics OR "ferrous" OR "ferric" OR "hematinic" OR "haematinic" OR "haematinics" OR "iron compounds"

AND

"randomized controlled trial" OR "controlled clinical trial" OR "clinical trial" OR randomized OR randomised OR placebo OR random* OR trial

Appendix 2: Variables extracted from eligible studies.**Study characteristics**

Endnote Record Number
 First author (last name)
 Publication year
 Corresponding author (full name)
 Corresponding author (email address)
 Study name
 Country
 Rural/Urban
 Altitude for the study site (meters)
 According to study report, is malaria endemic?
 According to study report, are helminths endemic?
 Is this a subgroup analysis of a priorly extracted study?
 Which subgroup is being reported here?
 Does this paper reference other publications using the same trial or data? If so, indicate the Endnote record numbers for those studies.
 Notes on this paper

Intervention characteristics

Mode of supplementation
 Dose of intervention (mg of elemental iron)
 Dose of intervention (mg) per kg bodyweight
 Mean body weight for the intervention group (kg).
 Frequency of intervention
 Unit of intervention frequency
 Formulation of Iron
 Total dose of supplement (used for conversion to elemental iron if elemental unavailable)
 Unit for dose
 Duration of the intervention (weeks) (there are 4.3 weeks per month)
 Is outcome data post intervention endline available?
 Year at baseline (if range pick median)
 Were all children in the study dewormed as part of the study protocol?
 What other compounds were given along with iron (to both intervention & control groups)?
 Is this a factorial trial?
 Daily dose consumed of other compound given.
 Unit for daily dose of other compound given.
 Did the control group receive a placebo?
 Is the control group being compared to multiple intervention groups?
 Population group targeted by intervention
 List study inclusion and exclusion criteria
 Is compliance reported in the paper?
 Indicate the compliance data reported
 Other notes on intervention

Population characteristics

Is population characteristic data given for the full sample or for the control group only? (Use the full sample data whenever possible.)

Proportion of female children at baseline (%)
 Child age at supplementation start/baseline (mean/median months)
 Child age at baseline (SD)
 Child age at baseline (min)
 Child age at baseline (max)
 Child age at outcome assessment (mean/median months)
 Child age at outcome assessment (SD)
 Age reported if other than age at supplementation start
 Mean HAZ at baseline
 Mean WHZ at baseline
 Mean WAZ at baseline
 Mean BAZ at baseline
 Proportion of iron deficient children if reported (%)
 Cutoff used to define iron deficiency
 Proportion of anemic children if reported (%)
 Cutoff used to define anemia (g/dl)
 Proportion of children with iron deficiency anemia (%)
 Cutoff used to define iron deficiency anemia
 Proportion of children stunted at baseline (%)
 Proportion of children low birth weight at baseline (%)
 Cutoff used to define low birth weight (g)
 Proportion of children born premature (%)
 Cutoff used to define prematurity (wks)
 Maternal age at baseline (mean/median years)
 Maternal age at baseline (SD)
 If pregnant, gestational age at baseline (mean/median weeks)
 If pregnant, gestational age (sd)
 If any family socioeconomic status measures are reported, please describe
 Proportion of mothers who are iron deficient, if reported (%)
 Proportion of mothers anemic at baseline, if reported (%)

Outcome characteristics

Specify outcome
 Outcome unit
 Method of Assessment
 Extra information pertaining to outcome
 Comparator group
 How many weeks after baseline was the outcome measured?
 How many participants were initially recruited for treatment?
 How many participants were initially recruited for control?
 Intervention total group sample size (total analyzed)
 Was the difference-in-differences measure calculated in an adjusted regression model?
 Difference in the differences between baseline & follow-up between intervention & control group (mean)
 Difference in the differences between baseline & follow-up between intervention & control group (SD)
 Difference in the differences between baseline & follow-up between intervention & control group (SE)

Difference in the differences between baseline & follow-up between intervention & control group (LCI)
Difference in the differences between baseline & follow-up between intervention & control group (UCI)
Difference in the differences between baseline & follow-up between intervention & control group (p-value)
Intervention group baseline sample size
Intervention group baseline estimate (mean)
Intervention group baseline estimate (SD)
Intervention group baseline estimate (SE)
Intervention group baseline estimate (LCI)
Intervention group baseline estimate (UCI)
Intervention group baseline estimate (LIQR)
Intervention group baseline estimate (UIQR)
Intervention group follow-up sample size
Intervention group follow-up estimate (mean)
Intervention group follow-up estimate (SD)
Intervention group follow-up estimate (SE)
Intervention group follow-up estimate (LCI)
Intervention group follow-up estimate (UCI)
Intervention group follow-up estimate (LIQR)
Intervention group follow-up estimate (UIQR)
Was the intervention group difference measure calculated in an adjusted regression model?
Intervention group difference sample size
Intervention group difference in outcome between baseline & follow-up (mean)
Intervention group difference in outcome between baseline & follow-up (SD)
Intervention group difference in outcome between baseline & follow-up (SE)
Intervention group difference in outcome between baseline & follow-up (LCI)
Intervention group difference in outcome between baseline & follow-up (UCI)
Intervention group difference in outcome between baseline & follow-up (LIQR)
Intervention group difference in outcome between baseline & follow-up (UIQR)
Intervention group difference in outcome between baseline & follow-up (p-value)
Control group baseline sample size
Control group baseline estimate (mean)
Control group baseline estimate (SD)
Control group baseline estimate (SE)
Control group baseline estimate (LCI)
Control group baseline estimate (UCI)
Control group baseline estimate (LIQR)
Control group baseline estimate (UIQR)
Control group follow-up sample size
Control group follow-up estimate (mean)
Control group follow-up estimate (SD)
Control group follow-up estimate (SE)
Control group follow-up estimate (LCI)
Control group follow-up estimate (UCI)
Control group follow-up estimate (LIQR)

Control group follow-up estimate (UIQR)
 Was the control group difference measure calculated in an adjusted regression model?
 Control group difference sample size
 Control group difference in outcome between baseline & follow-up (mean)
 Control group difference in outcome between baseline & follow-up (SD)
 Control group difference in outcome between baseline & follow-up (SE)
 Control group difference in outcome between baseline & follow-up (LCI)
 Control group difference in outcome between baseline & follow-up (UCI)
 Control group difference in outcome between baseline & follow-up (LIQR)
 Control group difference in outcome between baseline & follow-up (UIQR)
 Control group difference in outcome between baseline & follow-up (p-value)
 Flag indicating whether SD/SE was imputed during data entry
 Note other outcomes reported in the study but not extracted

Subgroup data

Was sub-group data on outcome presented by gender?
 Was sub-group data on outcome presented by age group?
 Was sub-group data on outcome presented by SES group?
 Was sub-group data on outcome presented by baseline anemia status?
 Was sub-group data on outcome presented by baseline iron status?
 Was sub-group data on outcome presented by baseline stunting/birthweight status?
 Other subgroup reported? (specify)

Cochrane risk of bias assessment

Method for generating randomization sequence
 How were individuals randomly assigned to treatment?
 Number of clusters randomized (total)
 Risk of bias due to randomization method?
 Method for concealing treatment allocation
 Risk of bias due to allocation concealment?
 Were participants blinded?
 Were supplementation providers blinded?
 Bias due to lack of participant or supplement provider blinding?
 Were outcome assessors blinded?
 Bias due to lack of outcome assessor blinding?
 Percent drop-out in intervention group
 Percent drop-out in control group
 Were reasons for withdrawals similar across treatment groups?
 Were reasons for withdrawals associated with other important covariates?
 If withdrawals were related to the outcome, describe the reason for withdrawal.
 Were appropriate methods used to impute missing data?
 Was an intention-to-treat analyses conducted (i.e. analyzed as randomized)?
 Was a per protocol analysis conducted?
 Risk of bias due to incomplete outcome data?
 Were data reported for all pre-specified outcomes?
 Were data reported for all pre-specified sub-group analyses
 Risk of bias due to selective outcome reporting?

Appendix 3. Elemental iron supplementation dose tertile ranges.

	Tertile		
	Low	Middle	High
mg/week			
0 - 5 mo	18.2 – 50.0	51.5 – 70.0	79.8 - 99.4
6 - 23 mo	20.0 – 70.0	87.5 – 87.5	105.0 – 462.0
24 - 59 mo	20.0 – 100.0	114.0 – 207.9	210.0 – 420.0
5 - 11 y	18.2 – 120.0	143.6 – 380.8	403.1 – 650.0
12 - 19 y	50.0 – 100.0	120.0 – 250.0	300.0 – 1820.0
mg/day			
0 - 5 mo	2.6 – 7.1	7.4 – 10.0	11.4 – 14.2
6 - 23 mo	2.9 – 10.0	12.5 – 12.5	15.0 – 66.0
24 - 59 mo	2.9 – 14.3	16.3 – 29.7	30.0 – 60.0
5 - 11 y	2.6 – 17.1	20.5 – 54.4	57.6 – 92.9
12 - 19 y	7.1 – 14.3	17.1 – 35.7	42.9 – 260.0

Note 1: The recommended dose of iron increases as children age. To account for the strong correlation between age and dose, we grouped children into five age categories and then categorized them into low, moderate, or high tertiles based on the total amount of iron they received each week.

Note 2: Tertiles reported here are for studies which reported hematologic outcomes and had sufficient data to calculate dose (n=94 iron treatment groups).

Appendix 4. References to included studies.

1. Adish AA, Esrey SA, Gyorkos TW, et al. Effect of consumption of food cooked in iron pots on iron status and growth of young children: a randomised trial. *Lancet (London, England)* 1999;353(9154):712-6. doi: 10.1016/s0140-6736(98)04450-x [published Online First: 1999/03/12]
2. Aggarwal D, Sachdev HP, Nagpal J, et al. Haematological effect of iron supplementation in breast fed term low birth weight infants. *Archives of disease in childhood* 2005;90(1):26-9. doi: 10.1136/adc.2003.040410 [published Online First: 2004/12/23]
3. Aguayo VM. School-administered weekly iron supplementation--effect on the growth and hemoglobin status of non-anemic Bolivian school-age children. A randomized placebo-controlled trial. *European journal of nutrition* 2000;39(6):263-9. [published Online First: 2001/06/09]
4. Akman M, Cebeci D, Okur V, et al. The effects of iron deficiency on infants' developmental test performance. *Acta paediatrica (Oslo, Norway : 1992)* 2004;93(10):1391-6. [published Online First: 2004/10/27]
5. Angeles IT, Schultink WJ, Matulesi P, et al. Decreased rate of stunting among anemic Indonesian preschool children through iron supplementation. *The American journal of clinical nutrition* 1993;58(3):339-42. [published Online First: 1993/09/01]
6. Angulo-Barroso RML, M.: Santos, D. C.: Bian, Y.: Sturza, J.: Jiang, Y.: Kaciroti, N.: Richards, B.: Lozoff, B. Iron Supplementation in Pregnancy or Infancy and Motor Development: A Randomized Controlled Trial. *Pediatrics* 2016;137(4) doi: 10.1542/peds.2015-3547 [published Online First: 2016/03/05]
7. Arcanjo FP, Arcanjo CC, Amancio OM, et al. Weekly iron supplementation for the prevention of anemia in pre-school children: a randomized, double-blind, placebo-controlled trial. *Journal of tropical pediatrics* 2011;57(6):433-8. doi: 10.1093/tropej/fmq119 [published Online First: 2011/02/03]
8. Arcanjo FPN, Santos PR, Arcanjo CPC, et al. Daily and Weekly Iron Supplementations are Effective in Increasing Hemoglobin and Reducing Anemia in Infants. *Journal of tropical pediatrics* 2013;59(3):175-79. doi: 10.1093/tropej/fms071
9. Aukett MA, Parks YA, Scott PH, et al. Treatment with iron increases weight gain and psychomotor development. *Archives of disease in childhood* 1986;61(9):849-57. [published Online First: 1986/09/01]
10. Ayoya MA, Spiekermann-Brouwer GM, Traore AK, et al. Multiple micronutrients including iron are not more effective than iron alone for improving hemoglobin and iron status of Malian school children. *The Journal of nutrition* 2009;139(10):1972-9. doi: 10.3945/jn.109.106625 [published Online First: 2009/08/14]
11. Ayoya MA, Spiekermann-Brouwer GM, Traore AK, et al. Effect on school attendance and performance of iron and multiple micronutrients as adjunct to drug treatment of Schistosoma-infected anemic schoolchildren. *Food and nutrition bulletin* 2012;33(4):235-41. [published Online First: 2013/02/22]
12. Ballin A, Berar M, Rubinstein U, et al. Iron state in female adolescents. *American journal of diseases of children (1960)* 1992;146(7):803-5. [published Online First: 1992/07/01]
13. Baqui AH, Zaman K, Persson LA, et al. Simultaneous weekly supplementation of iron and zinc is associated with lower morbidity due to diarrhea and acute lower respiratory infection in Bangladeshi infants. *The Journal of nutrition* 2003;133(12):4150-7. [published Online First: 2003/12/04]
14. Barclay SM, Aggett PJ, Lloyd DJ, et al. Reduced erythrocyte superoxide dismutase activity in low birth weight infants given iron supplements. *Pediatric research* 1991;29(3):297-301. doi: 10.1203/00006450-199103000-00015 [published Online First: 1991/03/01]
15. Baumgartner JS, C. M.: Malan, L.: Kvalsvig, J.: van Stuijvenberg, M. E.: Hurrell, R. F.: Zimmermann, M. B. Effects of iron and n-3 fatty acid supplementation, alone and in combination, on cognition in school children: a randomized, double-blind, placebo-controlled intervention in South Africa. *The American journal of clinical nutrition* 2012;96(6):1327-38. doi: 10.3945/ajcn.112.041004 [published Online First: 2012/10/26]
16. Berger J, Aguayo VM, Tellez W, et al. Weekly iron supplementation is as effective as 5 day per week iron supplementation in Bolivian school children living at high altitude. *European journal of clinical nutrition* 1997;51(6):381-6. [published Online First: 1997/06/01]
17. Berger J, Dyck JL, Galan P, et al. Effect of daily iron supplementation on iron status, cell-mediated immunity, and incidence of infections in 6-36 month old Togolese children. *European journal of clinical nutrition* 2000;54(1):29-35. [published Online First: 2000/03/01]

18. Berger J, Ninh NX, Khan NC, et al. Efficacy of combined iron and zinc supplementation on micronutrient status and growth in Vietnamese infants. *European journal of clinical nutrition* 2006;60(4):443-54. doi: 10.1038/sj.ejcn.1602336 [published Online First: 2005/11/25]
19. Berglund S, Westrup B, Domellof M. Iron supplements reduce the risk of iron deficiency anemia in marginally low birth weight infants. *Pediatrics* 2010;126(4):e874-83. doi: 10.1542/peds.2009-3624 [published Online First: 2010/09/08]
20. Bhatia D, Seshadri S. Growth performance in anemia and following iron supplementation. *Indian pediatrics* 1993;30(2):195-200. [published Online First: 1993/02/01]
21. Black MM, Baqui AH, Zaman K, et al. Iron and zinc supplementation promote motor development and exploratory behavior among Bangladeshi infants. *American Journal of Clinical Nutrition* 2004;80(4):903-10.
22. Bora R, Ramasamy S, Brown B, et al. Effect of iron supplementation from neonatal period on the iron status of 6-month-old infants at-risk for early iron deficiency: a randomized interventional trial. *J Matern-Fetal Neonatal Med* 2019 doi: 10.1080/14767058.2019.1638358
23. Bruner AB, Joffe A, Duggan AK, et al. Randomised study of cognitive effects of iron supplementation in non-anaemic iron-deficient adolescent girls. *Lancet (London, England)* 1996;348(9033):992-96. doi: 10.1016/S0140-6736(96)02341-0
24. Burman D. Haemoglobin levels in normal infants aged 3 to 24 months, and the effect of iron. *Archives of disease in childhood* 1972;47(252):261-71. doi: 10.1136/adc.47.252.261
25. Buzina-Suboticanec KB, R.: Stavljenic, A.: Tadinac-Babic, M.: Juhovic-Markus, V. Effects of iron supplementation on iron nutrition status and cognitive functions in children. *Food and nutrition bulletin* 1998;19(4):298-306.
26. Charoenlarp P, Pholpothi T, Chatpunyaporn P, et al. The effect of riboflavin on the hematologic changes in iron supplementation of schoolchildren. *The Southeast Asian journal of tropical medicine and public health* 1980;11(1):97-103. [published Online First: 1980/03/01]
27. Chen KL, Y. F.: Chen, L.: Zhang, X.: Liu, Y. X.: Chen, J.: Li, T. Y. [Effects of vitamin A, vitamin A plus iron and multiple micronutrient-fortified seasoning powder on iron metabolic homeostasis]. *Zhonghua er ke za zhi Chinese journal of pediatrics* 2011; 49(12). <http://onlinelibrary.wiley.com/doi/10.1111/j.1532-2039.2011.00881.x>
28. Chen K, Chen XR, Zhang L, et al. Effect of simultaneous supplementation of vitamin A and iron on diarrheal and respiratory tract infection in preschool children in Chengdu City, China. *Nutrition (Burbank, Los Angeles County, Calif)* 2013;29(10):1197-203. doi: 10.1016/j.nut.2013.03.025
29. Chen K, Zhang L, Luo HY, et al. No enhancing effect of vitamin A administration on iron absorption or body total iron content in preschool children from Chengdu, China. *Journal of nutritional science and vitaminology* 2014;60(4):223-30. [published Online First: 2014/10/10]
30. Cheng XHP, H.: Wang, Y. B. [The Effect of Low Molecular Protein Iron on Iron Deficiency Anemia]. *Zhongguo shi yan xue ye xue za zhi / Zhongguo bing li sheng li xue hui = Journal of experimental hematology / Chinese Association of Pathophysiology* 2001;9(3):240-42. [published Online First: 2003/02/13]
31. Choe YH, Kim SK, Son BK, et al. Randomized placebo-controlled trial of Helicobacter pylori eradication for iron-deficiency anemia in preadolescent children and adolescents. *Helicobacter* 1999;4(2):135-9. [published Online First: 1999/06/26]
32. Chwang LC, Soemantri AG, Pollitt E. Iron supplementation and physical growth of rural Indonesian children. *The American journal of clinical nutrition* 1988;47(3):496-501. [published Online First: 1988/03/01]
33. Das BK, Bal MS, Tripathi AM, et al. Evaluation of frequency and dose of iron and other hematinics--an alternative strategy for anemia prophylaxis in rural preschoolers. *Indian pediatrics* 1984;21(12):933-8. [published Online First: 1984/12/01]
34. de Silva AA, S.: Weerasinghe, I.: Ahluwalia, N. Iron supplementation improves iron status and reduces morbidity in children with or without upper respiratory tract infections: a randomized controlled study in Colombo, Sri Lanka. *The American journal of clinical nutrition* 2003;77(1):234-41. [published Online First: 2002/12/25]
35. Desai MR, Mei JV, Kariuki SK, et al. Randomized, controlled trial of daily iron supplementation and intermittent sulfadoxine-pyrimethamine for the treatment of mild childhood anemia in western Kenya. *Journal of Infectious Diseases* 2003;187(4):658-66. doi: 10.1086/367986

36. Devaki PBC, R. K.: Geisser, P. Effects of oral supplementation with iron(III) hydroxide polymaltose complex on the hematological profile of adolescents with varying iron status. *Arzneimittel-Forschung* 2008;58(8):389-97. doi: 10.1055/s-0031-1296526 [published Online First: 2008/09/24]
37. Dewey KG, Domellof M, Cohen RJ, et al. Iron supplementation affects growth and morbidity of breast-fed infants: Results of a randomized trial in Sweden and Honduras. *Journal of Nutrition* 2002;132(11):3249-55.
38. Dijkhuizen MA, Wieringa FT, West CE, et al. Effects of iron and zinc supplementation in Indonesian infants on micronutrient status and growth. *The Journal of nutrition* 2001;131(11):2860-5. [published Online First: 2001/11/06]
39. Dijkhuizen MA, Winichagoon P, Wieringa FT, et al. Zinc supplementation improved length growth only in anemic infants in a multi-country trial of iron and zinc supplementation in South-East Asia. *The Journal of nutrition* 2008;138(10):1969-75. [published Online First: 2008/09/23]
40. Domellof M, Cohen RJ, Dewey KG, et al. Iron supplementation of breast-fed Honduran and Swedish infants from 4 to 9 months of age. *The Journal of pediatrics* 2001;138(5):679-87. doi: 10.1067/mpd.2001.112895 [published Online First: 2001/05/09]
41. Dossa RA, Ategbro EA, de Koning FL, et al. Impact of iron supplementation and deworming on growth performance in preschool Beninese children. *European journal of clinical nutrition* 2001;55(4):223-8. doi: 10.1038/sj.ejcn.1601126 [published Online First: 2001/05/22]
42. Dossa RAA, E. A.: Van Raaij, J. M.: de Graaf, C.: Hautvast, J. G. Multivitamin-multimineral and iron supplementation did not improve appetite of young stunted and anemic Beninese children. *The Journal of nutrition* 2001;131(11):2874-9. [published Online First: 2001/11/06]
43. Eftekhari MH, Simondon KB, Jalali M, et al. Effects of administration of iron, iodine and simultaneous iron-plus-iodine on the thyroid hormone profile in iron-deficient adolescent Iranian girls. *European journal of clinical nutrition* 2006;60(4):545-52. doi: 10.1038/sj.ejcn.1602349 [published Online First: 2005/12/13]
44. Elwood PCW, W. E.: Greene, W. J. W. EVALUATION OF IRON SUPPLEMENTS IN PREVENTION OF IRON-DEFICIENCY ANÆMIA. *The Lancet* 1970;296(7665):175-77. doi: 10.1016/S0140-6736(70)92534-1
45. Engstrom EM, Castro IR, Portela M, et al. Effectiveness of daily and weekly iron supplementation in the prevention of anemia in infants. *Revista de saude publica* 2008;42(5):786-95. [published Online First: 2008/08/19]
46. Ermis B, Demirel F, Demircan N, et al. Effects of three different iron supplementations in term healthy infants after 5 months of life. *Journal of tropical pediatrics* 2002;48(5):280-4. [published Online First: 2002/10/31]
47. Fahmida U, Rumawas JS, Utomo B, et al. Zinc-iron, but not zinc-alone supplementation, increased linear growth of stunted infants with low haemoglobin. *Asia Pacific journal of clinical nutrition* 2007;16(2):301-9. [published Online First: 2007/05/01]
48. Fallahi E, Kimiagar M, Nazari A, et al. Effect of zinc and iron supplementation on indicators of iron, zinc and vitamin A status of primary school children. *Pakistan journal of biological sciences : PJBS* 2007;10(7):1088-92. [published Online First: 2007/04/01]
49. Franz AR, Mihatsch WA, Sander S, et al. Prospective randomized trial of early versus late enteral iron supplementation in infants with a birth weight of less than 1301 grams. *Pediatrics* 2000;106(4):700-6. [published Online First: 2000/10/04]
50. Friel JK, Aziz K, Andrews WL, et al. A double-masked, randomized control trial of iron supplementation in early infancy in healthy term breast-fed infants. *The Journal of pediatrics* 2003;143(5):582-6. doi: 10.1067/s0022-3476(03)00301-9 [published Online First: 2003/11/15]
51. Fujii T, Maruyama K, Koizumi T. Oral iron supplementation in preterm infants treated with erythropoietin. *Pediatrics international : official journal of the Japan Pediatric Society* 2004;46(6):635-9. doi: 10.1111/j.1442-200x.2004.01979.x [published Online First: 2005/01/22]
52. Gebreselassie H. Iron supplementation and malaria infection: results of a randomized controlled field trial. McGill University, 1996.
53. Geltman PL, Meyers AF, Bauchner H. Daily multivitamins with iron to prevent anemia in infancy: a randomized clinical trial. *Clinical pediatrics* 2001;40(10):549-54. [published Online First: 2001/10/30]
54. Geltman PL, Meyers AF, Mehta SD, et al. Daily multivitamins with iron to prevent anemia in high-risk infants: a randomized clinical trial. *Pediatrics* 2004;114(1):86-93. [published Online First: 2004/07/03]

55. Gokcay G, Ozden T, Karakas Z, et al. Effect of iron supplementation on development of iron deficiency anemia in breastfed infants. *Journal of tropical pediatrics* 2012;58(6):481-5. doi: 10.1093/tropej/fms028 [published Online First: 2012/07/04]
56. Gopaldas T, Kale M. Prophylactic iron supplementation for underprivileged school boys. I. Two levels of dosing and efficacy of teacher-distributions. *Indian pediatrics* 1985;22(10):731-6. [published Online First: 1985/10/01]
57. Gopaldas T, Kale M, Bhardwaj P. Prophylactic iron supplementation for underprivileged school boys. II. Impact on selected tests of cognitive function. *Indian pediatrics* 1985;22(10):737-43. [published Online First: 1985/10/01]
58. Greisen G. Mild anaemia in African school children: effect on running performance and an intervention trial. *Acta paediatrica Scandinavica* 1986;75(4):662-7. [published Online First: 1986/07/01]
59. Hacıhamdioglu DOK, A. E.: Gursel, O.: Atay, A. A.: Balamtekin, N.: Aydin, A.: Hasimi, A.: Ozcan, O. Evaluation of lipid peroxidation and antioxidant system in healthy iron-replete infants receiving iron prophylaxis. *Nutrition (Burbank, Los Angeles County, Calif)* 2013;29(1):138-42. doi: 10.1016/j.nut.2012.05.009 [published Online First: 2012/11/03]
60. Harvey PWH, P. F.: Nesheim, M. C.: Galme, K.: Zegans, M.: Habicht, J. P.: Stephenson, L. S.: Radimer, K. L.: Brabin, B.: Forsyth, K.: et al.,. The effect of iron therapy on malarial infection in Papua New Guinean schoolchildren. *The American journal of tropical medicine and hygiene* 1989;40(1):12-8. [published Online First: 1989/01/01]
61. Hathirat P, Valyasevi A, Kotchabhakdi NJ, et al. Effects of an iron supplementation trial on the Fe status of Thai schoolchildren. *The British journal of nutrition* 1992;68(1):245-52. [published Online First: 1992/07/01]
62. Hess SY, Zimmermann MB, Adou P, et al. Treatment of iron deficiency in goitrous children improves the efficacy of iodized salt in Cote d'Ivoire. *The American journal of clinical nutrition* 2002; 75(4). <http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/467/CN-00593467/frame.html>.
63. Hettiarachchi M, Liyanage C, Wickremasinghe R, et al. The efficacy of micronutrient supplementation in reducing the prevalence of anaemia and deficiencies of zinc and iron among adolescents in Sri Lanka. *European journal of clinical nutrition* 2008;62(7):856-65. doi: 10.1038/sj.ejcn.1602791
64. Hieu NT, Sandalinas F, de Sesmaisons A, et al. Multi-micronutrient-fortified biscuits decreased the prevalence of anaemia and improved iron status, whereas weekly iron supplementation only improved iron status in Vietnamese school children. *The British journal of nutrition* 2012;108(8):1419-27. doi: 10.1017/S0007114511006945 [published Online First: 2012/01/17]
65. Hop LT, Berger J. Multiple micronutrient supplementation improves anemia, micronutrient nutrient status, and growth of Vietnamese infants: Double-blind, randomized, placebo-controlled trial. *Journal of Nutrition* 2005;135(3):660S-65S.
66. Htet MK, Fahmida U, Dillon D, et al. Is Iron Supplementation Influenced by Sub-Clinical Inflammation?: A Randomized Controlled Trial Among Adolescent Schoolgirls in Myanmar. *Nutrients* 2019;11(4) doi: 10.3390/nu11040918 [published Online First: 2019/04/27]
67. Idjradinata P, Pollitt E. Reversal of developmental delays in iron-deficient anaemic infants treated with iron. *Lancet (London, England)* 1993;341(8836):1-4. [published Online First: 1993/01/02]
68. Idjradinata P, Watkins WE, Pollitt E. Adverse effect of iron supplementation on weight gain of iron-replete young children. *Lancet (London, England)* 1994;343(8908):1252-4. [published Online First: 1994/05/21]
69. Irigoyen M, Davidson LL, Carriero D, et al. Randomized, placebo-controlled trial of iron supplementation in infants with low hemoglobin levels fed iron-fortified formula. *Pediatrics* 1991;88(2):320-6. [published Online First: 1991/08/01]
70. Jalambo M, Karim N, Naser I, et al. Effects of iron supplementation and nutrition education on haemoglobin, ferritin and oxidative stress in iron-deficient female adolescents in Palestine: randomized control trial. *East Mediterr Health J* 2018;24(6):560-68. doi: 10.26719/2018.24.6.560 [published Online First: 2018/08/07]
71. Kapur D, Sharma S, Agarwal KN. Effectiveness of nutrition education, iron supplementation or both on iron status in children. *Indian pediatrics* 2003;40(12):1131-44. [published Online First: 2004/01/15]
72. Kashyap PG, T. Hematinic supplementation and hematological status of underprivileged school girls (8-15 YRS of age). *Nutrition Research* 1987;7(11):1127-38. doi: 10.1016/S0271-5317(87)80038-6
73. Kashyap P, Gopaldas T. Impact of hematinic supplementation on cognitive function in underprivileged school girls (8-15 Yrs of age). *Nutrition Research* 1987;7(11):1117-26. doi: 10.1016/S0271-5317(87)80037-4
74. Kianfar H, Kimiagar M, Ghaffarpour M. Effect of daily and intermittent iron supplementation on iron status of high school girls. *International journal for vitamin and nutrition research Internationale Zeitschrift fur Vitamin- und*

- Ernahrungsforschung Journal international de vitaminologie et de nutrition* 2000;70(4):172-7. doi: 10.1024/0300-9831.70.4.172 [published Online First: 2000/09/16]
75. Kordas K, Stoltzfus RJ, Lopez P, et al. Iron and zinc supplementation does not improve parent or teacher ratings of behavior in first grade Mexican children exposed to lead. *The Journal of pediatrics* 2005;147(5):632-9. doi: 10.1016/j.jpeds.2005.06.037 [published Online First: 2005/11/18]
76. Kusumastuti AC, Ardiaria M, Hendrianingtyas M. Effect of Zinc and Iron Supplementation on Appetite, Nutritional Status and Intelligence Quotient in Young Children. *Indonesian Biomedical Journal* 2018;10(2):133-39. doi: 10.18585/inabj.v10i2.365
77. Lambert AK, K.: Scragg, R.: Metcalf, P.: Schaaf, D. Effects of iron treatment on cognitive performance and working memory in non-anaemic, iron-deficient girls. *New Zealand Journal of Psychology* 2002;31(1):19-28.
78. Latham MC, Stephenson LS, Kinoti SN, et al. Improvements in growth following iron supplementation in young Kenyan school children. *Nutrition (Burbank, Los Angeles County, Calif)* 1990;6(2):159-65. [published Online First: 1990/03/01]
79. Lawless JW, Latham MC, Stephenson LS, et al. Iron supplementation improves appetite and growth in anemic Kenyan primary school children. *The Journal of nutrition* 1994;124(5):645-54. [published Online First: 1994/05/01]
80. Lee JWP, C. K. Effect of Vitamin C Supplementation on Iron Status of Adolescent Girls with Low Hemoglobin Levels. *Korean Journal of Community Nutrition* 1997; 2(5).
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/827/CN-01045827/frame.html>.
81. Leenstra T, Kariuki SK, Kurtis JD, et al. The effect of weekly iron and vitamin A supplementation on hemoglobin levels and iron status in adolescent schoolgirls in western Kenya. *European journal of clinical nutrition* 2009;63(2):173-82. doi: 10.1038/sj.ejcn.1602919 [published Online First: 2007/10/12]
82. Lind T, Lonnerdal B, Stenlund H, et al. A community-based randomized controlled trial of iron and zinc supplementation in Indonesian infants: interactions between iron and zinc. *American Journal of Clinical Nutrition* 2003;77(4):883-90.
83. Lind TL, B.: Stenlund, H.: Gamayanti, I. L.: Ismail, D.: Seswandhana, R.: Persson, L. A. A community-based randomized controlled trial of iron and zinc supplementation in Indonesian infants: effects on growth and development. *The American journal of clinical nutrition* 2004;80(3):729-36. [published Online First: 2004/08/24]
84. Loria ASM, L.: Arroyo, P. Nutritional anemia. VIII. Hemoglobin and plasma iron in infants treated prenatally with iron. *Nutrition Reports International* 1979;19(4):451-61.
85. Lozoff BJ, Y.: Li, X.: Zhou, M.: Richards, B.: Xu, G.: Clark, K. M.: Liang, F.: Kaciroti, N.: Zhao, G.: Santos, D. C.: Zhang, Z.: Tardif, T.: Li, M. Low-Dose Iron Supplementation in Infancy Modestly Increases Infant Iron Status at 9 Mo without Decreasing Growth or Increasing Illness in a Randomized Clinical Trial in Rural China. *The Journal of nutrition* 2016;146(3):612-21. doi: 10.3945/jn.115.223917 [published Online First: 2016/01/23]
86. Lopez de Romana GC, S.: Lopez de Romana, D.: Gross, R. Efficacy of multiple micronutrient supplementation for improving anemia, micronutrient status, growth, and morbidity of Peruvian infants. *The Journal of nutrition* 2005;135(3):646s-52s. [published Online First: 2005/03/01]
87. Majumdar I, Paul P, Talib VH, et al. The effect of iron therapy on the growth of iron-replete and iron-deplete children. *Journal of tropical pediatrics* 2003;49(2):84-8. [published Online First: 2003/05/06]
88. Malan L, Baumgartner J, Calder PC, et al. n-3 Long-chain PUFAs reduce respiratory morbidity caused by iron supplementation in iron-deficient South African schoolchildren: a randomized, double-blind, placebo-controlled intervention. *American Journal of Clinical Nutrition* 2015;101(3):668-79. doi: 10.3945/ajcn.113.081208
89. Massaga JJ, Kitua AY, Lemnge MM, et al. Effect of intermittent treatment with amodiaquine on anaemia and malarial fevers in infants in Tanzania: a randomised placebo-controlled trial. *Lancet (London, England)* 2003; 361(9372).
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/073/CN-00438073/frame.html>.
90. Mebrahtu T, Stoltzfus RJ, Chwaya HM, et al. Low-dose daily iron supplementation for 12 months does not increase the prevalence of malarial infection or density of parasites in young Zanzibari children. *Journal of Nutrition* 2004;134(11):3037-41.
91. Mejia LA, Chew F. Hematological effect of supplementing anemic children with vitamin A alone and in combination with iron. *The American journal of clinical nutrition* 1988;48(3):595-600. [published Online First: 1988/09/01]
92. Menendez C, Kahigwa E, Hirt R, et al. Randomised placebo-controlled trial of iron supplementation and malaria chemoprophylaxis for prevention of severe anaemia and malaria in Tanzanian infants. *Lancet (London, England)* 1997;350(9081):844-50. doi: 10.1016/s0140-6736(97)04229-3 [published Online First: 1997/10/06]

93. Menendez C, Schellenberg D, Quintero L, et al. The effects of short-term iron supplementation on iron status in infants in malaria-endemic areas. *American Journal of Tropical Medicine and Hygiene* 2004;71(4):434-40.
94. Metallinos-Katsaras E, Valassi-Adam E, Dewey KG, et al. Effect of iron supplementation on cognition in Greek preschoolers. *European journal of clinical nutrition* 2004; 58(11).
<http://onlinelibrary.wiley.com/doi/10.1046/j.1365-3010.2004.01361.x>
95. Mitra AK, Akramuzzaman SM, Fuchs GJ, et al. Long-term oral supplementation with iron is not harmful for young children in a poor community of Bangladesh. *Journal of Nutrition* 1997;127(8):1451-55.
96. Mozaffari-Khosravi H, Noori-Shadkam M, Fatehi F, et al. Once weekly low-dose iron supplementation effectively improved iron status in adolescent girls. *Biological trace element research* 2010;135(1-3):22-30. doi: 10.1007/s12011-009-8480-0 [published Online First: 2009/08/05]
97. Mwanri L, Worsley A, Ryan P, et al. Supplemental vitamin A improves anemia and growth in anemic school children in Tanzania. *The Journal of nutrition* 2000;130(11):2691-6. [published Online First: 2000/10/29]
98. Nagpal J, Sachdev HP, Singh T, et al. A randomized placebo-controlled trial of iron supplementation in breastfed young infants initiated on complementary feeding: effect on haematological status. *Journal of health, population, and nutrition* 2004;22(2):203-11. [published Online First: 2004/10/12]
99. Nair VKMB, A. S. Effect of Iron Prophylaxis in Preventing Iron Deficiency Anaemia in Term Infants. *Journal of Evolution of Medical and Dental Sciences-Jemds* 2017;6(7):559-61. doi: 10.14260/jemds/2017/119
100. Nchito M, Geissler PW, Mubila L, et al. The effect of iron and multi-micronutrient supplementation on *Ascaris lumbricoides* reinfection among Zambian schoolchildren. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2009;103(3):229-36. doi: 10.1016/j.trstmh.2008.08.005 [published Online First: 2008/10/22]
101. Northrop-Clewes CA, Paracha PI, McLoone UJ, et al. Effect of improved vitamin A status on response to iron supplementation in Pakistani infants. *The American journal of clinical nutrition* 1996;64(5):694-9. [published Online First: 1996/11/01]
102. Olsen A, Nawiri J, Friis H. The impact of iron supplementation on reinfection with intestinal helminths and *Schistosoma mansoni* in western Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2000;94(5):493-9. [published Online First: 2001/01/02]
103. Olsen A, Nawiri J, Magnussen P, et al. Failure of twice-weekly iron supplementation to increase blood haemoglobin and serum ferritin concentrations: results of a randomized controlled trial. *Annals of tropical medicine and parasitology* 2006;100(3):251-63. doi: 10.1179/136485906x91486 [published Online First: 2006/04/25]
104. Paganini D, Uyoga MA, Kortman GAM, et al. Prebiotic galacto-oligosaccharides mitigate the adverse effects of iron fortification on the gut microbiome: a randomised controlled study in Kenyan infants. *Gut* 2017;66(11):1956-67. doi: 10.1136/gutjnl-2017-314418 [published Online First: 2017/08/05]
105. Palupi L, Schultink W, Achadi E, et al. Effective community intervention to improve hemoglobin status in preschoolers receiving once-weekly iron supplementation. *The American journal of clinical nutrition* 1997;65(4):1057-61. [published Online First: 1997/04/01]
106. Paracha PIA, I.: Nawab, G. Effect of iron supplementation on biochemical indices of iron status in selected pre-adolescent schoolgirls in North West Frontier Province, Pakistan. *Asia Pacific journal of clinical nutrition* 1993;2(4):177-81. [published Online First: 1993/12/01]
107. Perrone L, Salerno M, Gialanella G, et al. Long-term zinc and iron supplementation in children of short stature: effect of growth and on trace element content in tissues. *Journal of trace elements in medicine and biology : organ of the Society for Minerals and Trace Elements (GMS)* 1999;13(1-2):51-6. doi: 10.1016/s0946-672x(99)80023-6 [published Online First: 1999/08/13]
108. Prasetyani MAdM, Q.: Afeworki, R.: Kaiser, M. M.: Stefanie, D.: Sartono, E.: Supali, T.: van der Ven, A. J. Effect of a short course of iron polymaltose on acquisition of malarial parasitaemia in anaemic Indonesian schoolchildren: a randomized trial. *Malaria journal* 2017;16(1):50. doi: 10.1186/s12936-017-1691-5 [published Online First: 2017/01/29]
109. Reeves JD, Yip R. Lack of adverse side effects of oral ferrous sulfate therapy in 1-year-old infants. *Pediatrics* 1985;75(2):352-5. [published Online First: 1985/02/01]
110. Rezaeian A, Ghayour-Mobarhan M, Mazloum SR, et al. Effects of iron supplementation twice a week on attention score and haematologic measures in female high school students. *Singapore medical journal* 2014;55(11):587-92. [published Online First: 2015/01/30]

111. Richard SA, Zavaleta N, Caulfield LE, et al. Zinc and iron supplementation and malaria, diarrhea, and respiratory infections in children in the Peruvian Amazon. *The American journal of tropical medicine and hygiene* 2006;75(1):126-32. [published Online First: 2006/07/14]
112. Rosado JL, Lopez P, Munoz E, et al. Zinc supplementation reduced morbidity, but neither zinc nor iron supplementation affected growth or body composition of Mexican preschoolers. *American Journal of Clinical Nutrition* 1997;65(1):13-19.
113. Rosado JL, Lopez P, Kordas K, et al. Iron and/or zinc supplementation did not reduce blood lead concentrations in children in a randomized, placebo-controlled trial. *The Journal of nutrition* 2006;136(9):2378-83. [published Online First: 2006/08/22]
114. Roschnik N, Parawan A, Baylon MA, et al. Weekly iron supplements given by teachers sustain the haemoglobin concentration of schoolchildren in the Philippines. *Tropical medicine & international health : TM & IH* 2004;9(8):904-9. doi: 10.1111/j.1365-3156.2004.01279.x [published Online First: 2004/08/12]
115. Sarker SA, Mahmud H, Davidsson L, et al. Causal relationship of Helicobacter pylori with iron-deficiency anemia or failure of iron supplementation in children. *Gastroenterology* 2008;135(5):1534-42. doi: 10.1053/j.gastro.2008.07.030 [published Online First: 2008/09/09]
116. Seshadri SG, T. Impact of iron supplementation on cognitive functions in preschool and school-aged children: the Indian experience. *The American journal of clinical nutrition* 1989;50(3 Suppl):675-84; discussion 85-6. [published Online First: 1989/09/01]
117. Smith AW, Hendrickse RG, Harrison C, et al. The effects on malaria of treatment of iron-deficiency anaemia with oral iron in Gambian children. *Annals of tropical paediatrics* 1989;9(1):17-23. [published Online First: 1989/03/01]
118. Smuts CM, Dhansay MA, Faber M, et al. Efficacy of multiple micronutrient supplementation for improving anemia, micronutrient status, and growth in south African infants. *Journal of Nutrition* 2005;135(3):653S-59S.
119. Smuts CMG, J.: Kvalsvig, J.: Zimmermann, M. B.: Baumgartner, J. Long-chain n-3 PUFA supplementation decreases physical activity during class time in iron-deficient South African school children. *The British journal of nutrition* 2014:1-13. doi: 10.1017/s0007114514003493 [published Online First: 2014/12/05]
120. Soemantri AG. Preliminary findings on iron supplementation and learning achievement of rural Indonesian children. *The American journal of clinical nutrition* 1989;50(3 Suppl):698-701; discussion 01-2. [published Online First: 1989/09/01]
121. Soewondo S, Husaini M, Pollitt E. Effects of iron deficiency on attention and learning processes in preschool children: Bandung, Indonesia. *The American journal of clinical nutrition* 1989;50(3 Suppl):667-73; discussion 73-4. [published Online First: 1989/09/01]
122. Stoltzfus RJ, Kvalsvig JD, Chwaya HM, et al. Effects of iron supplementation and anthelmintic treatment on motor and language development of preschool children in Zanzibar: double blind, placebo controlled study. *BMJ (Clinical research ed)* 2001;323(7326):1389-93. [published Online First: 2001/12/18]
123. Stoltzfus RJ, Chway HM, Montresor A, et al. Low dose daily iron supplementation improves iron status and appetite but not anemia, whereas quarterly anthelmintic treatment improves growth, appetite and anemia in Zanzibari preschool children. *The Journal of nutrition* 2004;134(2):348-56. [published Online First: 2004/01/30]
124. Sungthong R, Mo-Suwan L, Chongsuvivatwong V, et al. Once weekly is superior to daily iron supplementation on height gain but not on hematological improvement among schoolchildren in Thailand. *The Journal of nutrition* 2002;132(3):418-22. [published Online First: 2002/03/07]
125. Sungthong R, Mo-suwan L, Chongsuvivatwong V, et al. Once-weekly and 5-days a week iron supplementation differentially affect cognitive function but not school performance in Thai children. *The Journal of nutrition* 2004;134(9):2349-54. [published Online First: 2004/08/31]
126. Teshome EM, Andang'o PEA, Osoti V, et al. Daily home fortification with iron as ferrous fumarate versus NaFeEDTA: a randomised, placebo-controlled, non-inferiority trial in Kenyan children. *Bmc Medicine* 2017;15 doi: 10.1186/s12916-017-0839-z
127. Thibault H, Galan P, Selz F, et al. The immune response in iron-deficient young children: effect of iron supplementation on cell-mediated immunity. *European journal of pediatrics* 1993;152(2):120-4. [published Online First: 1993/02/01]

128. Untoro J, Karyadi E, Wibowo L, et al. Multiple micronutrient supplements improve micronutrient status and anemia but not growth and morbidity of Indonesian infants: a randomized, double-blind, placebo-controlled trial. *The Journal of nutrition* 2005;135(3):639s-45s. [published Online First: 2005/03/01]
129. van den Hombergh JD, E.: Smit, Y. Does iron therapy benefit children with severe malaria-associated anaemia? A clinical trial with 12 weeks supplementation of oral iron in young children from the Turiani Division, Tanzania. *Journal of tropical pediatrics* 1996;42(4):220-7. [published Online First: 1996/08/01]
130. Verhoef H, West CE, Nzyuko SM, et al. Intermittent administration of iron and sulfadoxine-pyrimethamine to control anaemia in Kenyan children: a randomised controlled trial. *Lancet (London, England)* 2002;360(9337):908-14. doi: 10.1016/S0140-6736(02)11027-0
131. Wang YJW, Q.: Yang, L. C.: Zhang, X. R.: Zeng, C. M.: Yang, X. G.: Liu, J. [The influence of preventive iron supplementation to iron nutritional status in breastfed infants]. [Chinese]. *Zhonghua yu fang yi xue za zhi [Chinese journal of preventive medicine]* 2012; 46(4).
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/710/CN-00920710/frame.html>.
132. Wasantwisut E, Winichagoon P, Chitchumroonchokchai C, et al. Iron and zinc supplementation improved iron and zinc status, but not physical growth, of apparently healthy, breast-fed infants in rural communities of northeast Thailand. *The Journal of nutrition* 2006;136(9):2405-11. [published Online First: 2006/08/22]
133. Wieringa FT, Berger J, Dijkhuizen MA, et al. Combined iron and zinc supplementation in infants improved iron and zinc status, but interactions reduced efficacy in a multicountry trial in southeast Asia. *Journal of Nutrition* 2007;137(2):466-71.
134. Yalçın S, Yurdakök K, Açıkgöz D, et al. Short-term developmental outcome of iron prophylaxis in infants. *Pediatrics International* 2000;42(6):625-30. doi: 10.1046/j.1442-200X.2000.01299.x
135. Yip RR, J. D.: Lonnerdal, B.: Keen, C. L.: Dallman, P. R. Does iron supplementation compromise zinc nutrition in healthy infants? *The American journal of clinical nutrition* 1985;42(4):683-7. [published Online First: 1985/10/01]
136. Yurdakök K, Temiz F, Yalçın SS, et al. Efficacy of Daily and Weekly Iron Supplementation on Iron Status in Exclusively Breast-Fed Infants. *Journal of pediatric hematology/oncology* 2004;26(5):284-88. doi: 10.1097/00043426-200405000-00005
137. Zavaleta N, Respicio G, Garcia T. Efficacy and acceptability of two iron supplementation schedules in adolescent school girls in Lima, Peru. *Journal of Nutrition* 2000;130(2):462S-64S.
138. Zhao X, Lu Q, Wang S, et al. [Efficiency of NaFeEDTA fortified soy sauce on anemia prevention]. *Wei sheng yan jiu = Journal of hygiene research* 2004;33(2):202-4. [published Online First: 2004/06/24]
139. Ziegler EE, Nelson SE, Jeter JM. Iron status of breastfed infants is improved equally by medicinal iron and iron-fortified cereal. *American Journal of Clinical Nutrition* 2009;90(1):76-87. doi: 10.3945/ajcn.2008.27350
140. Ziegler EE, Nelson SE, Jeter JM. Iron supplementation of breastfed infants from an early age. *The American journal of clinical nutrition* 2009;89(2):525-32. doi: 10.3945/ajcn.2008.26591 [published Online First: 2008/12/17]
141. Zlotkin SA, K. Y.: Schauer, C.: Yeung, G. Use of microencapsulated iron(II) fumarate sprinkles to prevent recurrence of anaemia in infants and young children at high risk. *Bulletin of the World Health Organization* 2003; 81(2).
<http://onlinelibrary.wiley.com/o/cochrane/clcentral/articles/518/CN-00437518/frame.html>:
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2572396/pdf/12756979.pdf>.
142. Zlotkin S, Newton S, Aimone AM, et al. Effect of iron fortification on malaria incidence in infants and young children in Ghana: a randomized trial. *Jama* 2013;310(9):938-47. doi: 10.1001/jama.2013.277129 [published Online First: 2013/09/05]

Appendix 5. Characteristics of included studies.

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Adish, 1997	Ethiopia	24 mo - 5 yr	30 mg elemental Fe as ferrous sulfate daily for 3 months	Placebo; or Placebo + Vitamin A (factorial trial)	n=399 total participants analyzed; n=196 treatment, n=203 control	Hemoglobin, Diarrhea, Respiratory infection
Aggarwal et al, 2005	India	50 days - 80 days	3 mg/kg elemental Fe as ferric ammonium citrate daily for 8 weeks	Placebo	n=26 total participants analyzed; n=13 treatment, n=13 control	Hemoglobin, Ferritin
Aguayo et al, 2000	Bolivia	6 yr - 12 yr	3 mg/kg elemental Fe as ferrous sulfate 1 time per week for 18 weeks	Placebo	n=64 total participants analyzed; n=33 treatment, n=31 control	Hemoglobin, Anemia, HAZ, WAZ
Akman et al, 2004	Turkey	6 mo - 30 mo	6 mg/kg elemental Fe as ferrous glisine-sulphate daily for 3 months	Control	n=40 total participants analyzed; n=21 treatment, n=19 control	Hemoglobin, Ferritin
Angeles et al, 1993	Indonesia	24 mo - 6 yr	30 mg elemental Fe as ferrous sulfate + Vitamin C daily for 2 months	Placebo + Vitamin C	n=76 total participants analyzed; n=39 treatment, n=37 control	Hemoglobin, Ferritin, HAZ, WHZ, WAZ, Diarrhea, Respiratory infection
Angulo-Barro et al, 2016	China	~1 mo (mean age)	1 mg/kg elemental Fe as iron protein succinylate daily for 7.5 months	Placebo; or Placebo + Maternal iron (factorial trial)	n=1196 total participants analyzed; n=610 treatment, n=586 control	Cognitive
Arcanjo et al, 2011	Brazil	~5 yr (mean age)	50 mg elemental Fe as ferrous sulfate 1 time per week for 14 weeks	Placebo	n=99 total participants analyzed; n=50 treatment, n=49 control	Hemoglobin, Anemia
Arcanjo et al, 2013	Brazil	12 mo - 24 mo	25 mg elemental iron (formulation not specified) 1 time per week for 4 months; 12.5 mg elemental iron (formulation not specified) daily for 4 months	Placebo	n=176 total participants analyzed; n=120 treatment, n=56 control	Hemoglobin, Anemia

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Aukett et al, 1986	United Kingdom	17 mo - 19 mo	24 mg elemental Fe as ferrous sulfate + Vitamin C daily for 2 months	Placebo + Vitamin C	n=97 total participants analyzed; n=48 treatment, n=49 control	Hemoglobin, Ferritin, Anemia, Iron deficiency
Ayoya et al, 2009	Mali	7 yr - 13 yr	60 mg elemental Fe as ferrous sulfate + Praziquantal 5 times per week for 12 weeks	Control + Praziquantal	n=202 total participants analyzed; n=105 treatment, n=97 control	Hemoglobin, Ferritin, Anemia, Iron deficiency anemia, Malaria
Ayoya et al, 2012	Mali	7 yr - 13 yr	60 mg elemental Fe as ferrous sulfate + Praziquantal 5 times per week for 12 weeks	Control + Praziquantal	n=202 total participants analyzed; n=105 treatment, n=97 control	Cognitive
Ballin et al, 1992	Israel	16 yr - 18 yr	105 mg elemental Fe as iron polystyrene sulfonate adsorbate daily for 2 months	Placebo	n=59 total participants analyzed; n=29 treatment, n=30 control	Iron deficiency
Baqui et al, 2003	Bangladesh	~6 mo (mean age)	20 mg elemental Fe as ferrous sulfate 1 time per week for 6 months	Placebo + Riboflavin; or Placebo + Zinc + Riboflavin (factorial trial)	n=249 total participants analyzed; n=125 treatment, n=124 control	Hemoglobin, Ferritin, Diarrhea, Respiratory infection
Barclay et al, 1991	United Kingdom	~1 mo (mean age)	13.8 mg elemental Fe as NaFeEDTA daily for 16 weeks; 7 mg elemental Fe as NaFeEDTA daily for 16 weeks	Control + Mutivitamin	n=55 total participants analyzed; n=36 treatment, n=19 control	Hemoglobin, Ferritin
Baumgartner et al, 2012	South Africa	6 yr - 12 yr	50 mg elemental Fe as ferrous sulfate 4 times per week for 8.5 months	Placebo + Vitamin C; or Placebo + n-3 fatty acids + Vitamin C (factorial trial)	n=294 total participants analyzed; n=145 treatment, n=149 control	Hemoglobin, Ferritin, HAZ, WHZ, WAZ

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Berger et al, 1997	Bolivia	3 yr - 8 yr	3-4 mg/kg elemental Fe as ferrous sulfate 1 time per week for 16 weeks; 3-4 mg/kg elemental Fe as ferrous sulfate 5 times per week for 16 weeks	Placebo	n=173 total participants analyzed; n=116 treatment, n=57 control	Hemoglobin, Anemia
Berger et al, 2000	Togo	6 mo - 3 yr	2-3 mg/kg elemental Fe as iron betainate daily for 3 months	Placebo	n=163 total participants analyzed; n=84 treatment, n=79 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Diarrhea, Respiratory infection, Malaria
Berger et al, 2006	Vietnam	4 mo - 7 mo	10 mg elemental Fe as ferrous sulfate 7 times per week for 6 months	Placebo + Vitamin A; or Placebo + Vitamin A + Zinc (factorial trial)	n=770 total participants analyzed; n=384 treatment, n=386 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia, HAZ, WHZ, WAZ, Stunting, Wasting, Diarrhea, Respiratory infection
Berglund et al, 2010	Sweden	~1 mo (mean age)	1 mg/kg elemental Fe as ferrous succinate daily for 4.5 months; 2 mg/kg elemental Fe as ferrous succinate daily for 4.5 months	Placebo	n=243 total participants analyzed; n=160 treatment, n=83 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Bhatia et al, 1993	India	~4 yr (mean age)	40 mg elemental iron (formulation not specified) daily for 6 months	Placebo	n=156 total participants analyzed; n=84 treatment, n=72 control	Hemoglobin
Black et al, 2004	Bangladesh	~7 mo (mean age)	20 mg elemental Fe as ferrous sulfate 1 time per week for 6 months	Placebo + Riboflavin + Vitamin A; or Placebo + Zinc + Riboflavin + Vitamin A (factorial trial)	n=186 total participants analyzed; n=92 treatment, n=94 control	Cognitive

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Bora et al, 2019	India	0 mo	2 mg/kg elemental Fe as ferrous ascorbate daily for 6 months	Placebo	n=180 total participants analyzed; n=90 treatment, n=90 control	Hemoglobin, Ferritin
Bruner et al, 1996	USA	13 yr - 19 yr	260 mg elemental Fe as ferrous sulfate daily for 8 weeks	Placebo	n=73 total participants analyzed; n=37 treatment, n=36 control	Hemoglobin, Ferritin
Burman et al, 1972	United Kingdom	~3 mo (mean age)	10 mg elemental Fe as colloidal ferric hydroxide daily for 21 months	Placebo	n=192 total participants analyzed; n=86 treatment, n=106 control	Hemoglobin
Buzina-Suboticanec et al, 1998	Croatia	8 yr - 10 yr	100 mg elemental Fe as ferriglycine sulfate 6 times per week for 10 weeks	Placebo	n=60 total participants analyzed; n=31 treatment, n=29 control	Hemoglobin, Cognitive
Charoenlarp et al, 1980	Thailand	6 yr - 13 yr	40 mg elemental Fe as ferrous sulfate 5 times per week for 5 months	Placebo	n=72 total participants analyzed; n=33 treatment, n=39 control	Hemoglobin, Helminth Infection
Chen et al, 2011	China	3 yr - 7 yr	12 mg elemental Fe as NaFeEDTA + Vitamin A 5 times per week for 26 weeks	Placebo + Vitamin A	n=132 total participants analyzed; n=71 treatment, n=61 control	Hemoglobin, Ferritin
Chen et al, 2013	China	3 yr - 7 yr	1.5 mg/kg elemental Fe as ferrous sulfate 5 times per week for 6 months	Placebo; or Placebo + Vitamin A (factorial trial)	n=387 total participants analyzed; n=188 treatment, n=199 control	Diarrhea, Respiratory infection
Chen et al, 2014	China	3 yr - 6 yr	1-2 mg/kg elemental Fe as ferrous sulfate 5 times per week for 6 months	Control; or Control + Vitamin A (factorial trial)	n=387 total participants analyzed; n=188 treatment, n=199 control	Hemoglobin, Ferritin, Anemia, Iron deficiency
Cheng et al, 2001	China	7 yr - 12 yr	2.6 mg elemental Fe as daily for 1 month	Placebo	n=108 total participants analyzed; n=53 treatment, n=55 control	Hemoglobin

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Choe et al, 1999	Korea	10 yr - 18 yr	6 mg/kg elemental Fe as ferrous sulfate + H. pylori therapy (bismuth subcitrate, amoxicillin, metronidazole) daily for 10 weeks	Placebo + H. pylori therapy (bismuth subcitrate, amoxicillin, metronidazole)	n=11 total participants analyzed; n=6 treatment, n=5 control	Hemoglobin, Ferritin
Chwang et al, 1988	Indonesia	8 yr - 14 yr	2 mg/kg elemental Fe as ferrous sulfate 5 times per week for 12 weeks	Placebo	n=119 total participants analyzed; n=59 treatment, n=60 control	Hemoglobin, HAZ, WAZ
Das et al, 1984	India	6 mo - 7 yr	5 mg elemental Fe as neoferrum infants drop 5 times per week for 12 weeks; 10 mg elemental Fe as neoferrum infants drop 5 times per week for 12 weeks; 20 mg elemental Fe as neoferrum infants drop 2 times per week for 12 weeks; 40 mg elemental Fe as neoferrum infants drop 2 times per week for 12 weeks; 20 mg elemental Fe as neoferrum infants drop 1 time per week for 12 weeks; 40 mg elemental Fe as neoferrum infants drop 1 time per week for 12 weeks	Placebo	n=151 total participants analyzed; n=115 treatment, n=36 control	Hemoglobin
de Silva et al, 2003	Sri Lanka	5 yr - 11 yr	60 mg elemental Fe as ferrous sulfate daily for 8 weeks	Placebo	n=363 total participants analyzed; n=261 treatment, n=102 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Diarrhea, Respiratory infection

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Desai et al, 2003	Kenya	2 mo - 3 yr	3.83 mg/kg elemental Fe as ferrous sulfate daily for 12 weeks	Placebo; or Placebo + Antimalarial (sulfadoxine-pyrimethamine at 4 and 8 weeks) (factorial trial)	n=491 total participants analyzed; n=256 treatment, n=235 control	Hemoglobin, Anemia, Malaria
Devaki et al, 2008	India	15 yr - 19 yr	100 mg elemental Fe as iron(iii) hydroxide polymaltose complex 6 times per week for 8 months	Placebo	n=60 total participants analyzed; n=30 treatment, n=30 control	Hemoglobin, Ferritin
Dewey et al, 2002	Sweden	~6 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 5 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 5 months	Placebo	n=172 total participants analyzed; n=136 treatment, n=36 control	HAZ, WAZ, Diarrhea
Dijkhuizen et al, 2001	Indonesia	~4 mo (mean age)	10 mg elemental Fe as ferrous sulfate 5 times per week for 6 months	Placebo; or Placebo + Zinc (factorial trial)	n=360 total participants analyzed; n=172 treatment, n=188 control	Hemoglobin, Ferritin, Anemia, Iron deficiency anemia, HAZ, WHZ, WAZ, Stunting, Wasting
Dijkhuizen et al, 2008	Indonesia, Thailand, Vietnam	4 mo - 6 mo	10 mg elemental Fe as ferrous sulfate + Zinc to half the population 5-7 times per week for 6 months	Placebo	n=2451 total participants analyzed; n=1212 treatment, n=1239 control	HAZ, WHZ, WAZ, Stunting, Wasting, Underweight

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Domellöf et al, 2001	Sweden	~6 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 5 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 1 mg/kg elemental Fe as ferrous sulfate daily for 5 months	Placebo	n=171 total participants analyzed; n=136 treatment, n=35 control	Hemoglobin, Ferritin
Dossa et al, 2001	Benin	3 yr - 6 yr	60 mg elemental Fe as ferrous sulfate daily for 3 months	Placebo; or Placebo + Albendazole (factorial trial)	n=140 total participants analyzed; n=70 treatment, n=70 control	Hemoglobin, HAZ, WHZ
Dossa et al, 2001	Benin	18 mo - 30 mo	66 mg elemental Fe as ferrous fumarate daily for 6 weeks	Placebo	n=74 total participants analyzed; n=35 treatment, n=39 control	Hemoglobin
Eftekhari et al, 2006	Iran	14 yr - 19 yr	60 mg elemental Fe as ferrous sulfate 5 times per week for 12 weeks	Placebo; or Placebo + Iodine (factorial trial)	n=94 total participants analyzed; n=47 treatment, n=47 control	Hemoglobin, Ferritin
Elwood et al, 1970	United Kingdom	14 yr - 15 yr	10 mg elemental Fe as ferrous fumarate daily for 36 months; 30 mg elemental Fe as ferrous fumarate daily for 36 months	Placebo + Riboflavin	n=601 total participants analyzed; n=386 treatment, n=215 control	Hemoglobin
Engstrom et al, 2008	Brazil	5 mo - 7 mo	12.5 mg elemental Fe as ferrous sulfate daily for 24 weeks; 25 mg elemental Fe as ferrous sulfate 1 time per week for 24 weeks	Placebo	n=391 total participants analyzed; n=297 treatment, n=94 control	Hemoglobin, Anemia

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Ermis et al, 2002	Turkey	~5 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate every other day for 4 months; 2 mg/kg elemental Fe as ferrous sulfate every other day for 4 months; 2 mg/kg elemental Fe as ferrous sulfate every other day for 4 months	Placebo	n=107 total participants analyzed; n=84 treatment, n=23 control	Hemoglobin, Ferritin, Iron deficiency, Iron deficiency anemia
Fahmida et al, 2007	Indonesia	3 mo - 6 mo	10 mg elemental Fe as ferrous sulfate + Zinc + Vitamin A daily for 6 months	Placebo + Zinc + Vitamin A	n=377 total participants analyzed; n=186 treatment, n=191 control	Hemoglobin, Ferritin, Anemia, HAZ, WHZ, WAZ, Stunting, Wasting, Underweight
Fallahi et al, 2007	Iran	~11 yr (mean age)	20 mg elemental Fe as ferrous sulfate + Zinc 6 times per week for 4 months	Placebo + Zinc	n=54 total participants analyzed; n=26 treatment, n=28 control	Hemoglobin, Ferritin
Franz et al, 2000	Germany	~0 mo (mean age)	2-4 mg/kg elemental Fe as ferrous sulfate daily for 8 weeks	Control	n=133 total participants analyzed; n=68 treatment, n=65 control	Ferritin, Iron deficiency
Friel et al, 2003	Canada	~1 mo (mean age)	7.5 mg elemental Fe as ferrous sulfate daily for 5 months	Placebo	n=49 total participants analyzed; n=28 treatment, n=21 control	Hemoglobin, Ferritin, Iron deficiency, Iron deficiency anemia
Fujiu et al, 2004	Japan	~1 mo (mean age)	4 mg/kg elemental iron (formulation not specified) + Recombinant human erythropoietin daily for 8 weeks	Control + Recombinant human erythropoietin	n=24 total participants analyzed; n=12 treatment, n=12 control	Hemoglobin, Ferritin
Gebresellasie, 1996	Ethiopia	5 yr - 15 yr	60 mg elemental Fe as ferrous sulfate daily for 12 weeks	Placebo	n=480 total participants analyzed; n=239 treatment, n=241 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Malaria

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Geltman et al, 2001	USA	~6 mo (mean age)	10 mg elemental iron (formulation not specified) + Multivitamin (A, B, C, D, E) daily for 3 months	Placebo + Multivitamin (A, B, C, D, E)	n=240 total participants analyzed; n=117 treatment, n=123 control	Anemia, Iron deficiency
Geltman et al, 2004	USA	~6 mo (mean age)	10 mg elemental Fe as + Multivitamin daily for 3 months	Placebo + Multivitamin	n=284 total participants analyzed; n=138 treatment, n=146 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Gokcay et al, 2012	Turkey	~6 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate + Vitamin D daily for 6 months	Control + Vitamin D	n=105 total participants analyzed; n=51 treatment, n=54 control	Hemoglobin, Anemia
Gopaldas et al, 1985	India	8 yr - 16 yr	26.2 mg elemental Fe as ferrous sulfate daily for 2 months; 37.4 mg elemental Fe as ferrous sulfate daily for 2 months	Placebo	n=210 total participants analyzed; n=140 treatment, n=70 control	Hemoglobin, Anemia
Gopaldas et al, 1985	India	8 yr - 16 yr	30 mg elemental Fe as ferrous sulfate daily for 2 months; 40 mg elemental Fe as ferrous sulfate daily for 2 months	Placebo	n=48 total participants analyzed; n=32 treatment, n=16 control	Cognitive
Greisen et al, 1986	Zambia	5 yr - 16 yr	66 mg elemental Fe as ferrous fumarate 5 times per week for 6 weeks	Placebo	n=430 total participants analyzed; n=218 treatment, n=212 control	Hemoglobin
Hacihamdioglu et al, 2013	Turkey	~4 mo (mean age)	10 mg elemental Fe as ferrous sulfate daily for 2 months	Control	n=53 total participants analyzed; n=27 treatment, n=26 control	Hemoglobin, Ferritin, Iron deficiency
Harvey et al, 1989	Papua New Guinea	8 yr - 13 yr	130 mg elemental Fe as ferrous sulfate 5 times per week for 16 weeks	Placebo	n=298 total participants analyzed; n=156 treatment, n=142 control	Hemoglobin, Malaria

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Hathirat et al, 1992	Thailand	9 yr - 12 yr	93.75 mg elemental Fe as ferrous sulfate 5 times per week for 16 weeks	Placebo	n=1772 total participants analyzed; n=885 treatment, n=887 control	Hemoglobin, Ferritin
Hess et al, 2002	Ivory Coast	5 yr - 15 yr	60 mg elemental Fe as ferrous sulfate 4 times per week for 16 weeks	Placebo	n=166 total participants analyzed; n=85 treatment, n=81 control	Hemoglobin, Ferritin, Anemia, Iron deficiency
Hettiarachchi et al, 2008	Sri Lanka	12 yr - 17 yr	50 mg elemental Fe as ferrous fumarate 5 times per week for 24 weeks	Placebo; or Placebo + Zinc (factorial trial)	n=774 total participants analyzed; n=392 treatment, n=382 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, HAZ, WAZ
Hieu et al, 2012	Vietnam	6 yr - 10 yr	1-2 mg/kg elemental Fe as ferrous fumarate 1 time per week for 21-23 weeks	Control	n=221 total participants analyzed; n=95 treatment, n=126 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Hop et al, 2005	Vietnam	6 mo - 12 mo	10 mg elemental Fe as ferrous fumarate 7 times per week for 6 months	Placebo	n=152 total participants analyzed; n=75 treatment, n=77 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, HAZ, WHZ, WAZ
Htet et al, 2019	Myanmar	~16 yr (mean age)	60 mg elemental iron (formulation not specified) 1 time per week for 12 weeks	Placebo + Folic acid; or Placebo + Vitamin A + Folic acid (factorial trial)	n=391 total participants analyzed; n=192 treatment, n=199 control	Hemoglobin, Ferritin
Idjradinata et al, 1993	Indonesia	12 mo - 18 mo	3 mg/kg elemental Fe as ferrous sulfate daily for 4 months	Placebo	n=119 total participants analyzed; n=60 treatment, n=59 control	Hemoglobin, Ferritin
Idjratinata et al, 1994	Indonesia	12 mo - 18 mo	3 mg/kg elemental Fe as ferrous sulfate daily for 16 weeks	Placebo	n=44 total participants analyzed; n=22 treatment, n=22 control	HAZ, WAZ

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Irigoyen et al, 1991	USA	~6 mo (mean age)	3 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 6 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Placebo	n=225 total participants analyzed; n=150 treatment, n=75 control	Hemoglobin, Ferritin, Diarrhea
Jalambo et al, 2018	Palestine	15 yr - 20 yr	200 mg elemental Fe as ferrous fumarate 1 time per week for 3 months	Control	n=87 total participants analyzed; n=45 treatment, n=42 control	Hemoglobin, Ferritin
Kapur et al, 2003	India	9 mo - 3 yr	20 mg elemental Fe as ferium, m/s emcure pharm india 1 time per week for 8 weeks	Placebo; or Placebo + Nutrition education (factorial trial)	n=232 total participants analyzed; n=116 treatment, n=116 control	Hemoglobin, Ferritin
Kashyap et al, 1987	India	8 yr - 15 yr	60 mg elemental Fe as ferrous sulfate daily for 4 months	Placebo	n=166 total participants analyzed; n=83 treatment, n=83 control	Hemoglobin, Anemia
Kashyap et al, 1987	India	8 yr - 15 yr	60 mg elemental Fe as ferrous sulfate daily for 4 months	Placebo	n=130 total participants analyzed; n=65 treatment, n=65 control	Cognitive
Kianfar et al, 2000	Iran	~16 yr (mean age)	50 mg elemental Fe as ferrous sulfate daily for 3 months; 50 mg elemental Fe as ferrous sulfate 2 times per week for 3 months; 50 mg elemental Fe as ferrous sulfate 1 time per week for 3 months	Control	n=268 total participants analyzed; n=194 treatment, n=74 control	Hemoglobin, Ferritin
Kordas et al, 2005	Mexico	~7 yr (mean age)	30 mg elemental Fe as ferrous fumarate daily for 6 months	Placebo	n=527 total participants analyzed; n=271 treatment, n=256 control	Cognitive

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Kusumastuti et al, 2018	Indonesia	25 mo - 5 yr	7.5 mg elemental iron (formulation not specified) daily for 3 months	Placebo; or Placebo + Zinc (factorial trial)	n=68 total participants analyzed; n=34 treatment, n=34 control	HAZ, WAZ
Lambert et al, 2002	New Zealand	12 yr - 18 yr	105 mg elemental Fe as ferrous sulfate daily for 8 weeks	Placebo	n=116 total participants analyzed; n=57 treatment, n=59 control	Hemoglobin, Ferritin
Latham et al, 1990	Kenya	~8 yr (mean age)	80 mg elemental Fe as ferrous sulfate daily for 15 weeks	Placebo	n=55 total participants analyzed; n=29 treatment, n=26 control	Hemoglobin, Malaria, Helminth Infection
Lawless et al, 1994	Kenya	6 yr - 12 yr	47 mg elemental Fe as ferrous sulfate daily for 14 weeks	Placebo	n=86 total participants analyzed; n=44 treatment, n=42 control	Hemoglobin, Ferritin, HAZ, WHZ, WAZ, Diarrhea, Malaria
Lee et al, 1997	Korea	12 yr - 16 yr	60 mg elemental Fe as ferrous sulfate daily for 9 weeks	Placebo	n=15 total participants analyzed; n=9 treatment, n=6 control	Hemoglobin, Ferritin
Leenstra et al, 2009	Kenya	12 yr - 19 yr	120 mg elemental Fe as ferrous sulfate + Vitamin A 1 time per week for 3 months	Placebo + Vitamin A	n=249 total participants analyzed; n=124 treatment, n=125 control	Hemoglobin, Ferritin, Anemia, Malaria
Lind et al, 2003	Indonesia	~6 mo (mean age)	10 mg elemental Fe as ferrous sulfate daily for 6 months	Placebo + Vitamin C; or Placebo + Vitamin C + Zinc (factorial trial)	n=549 total participants analyzed; n=272 treatment, n=277 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Lind et al, 2004	Indonesia	~6 mo (mean age)	10 mg elemental Fe as ferrous sulfate daily for 6 months	Placebo + Vitamin C; or Placebo + Zinc + Vitamin C (factorial trial)	n=656 total participants analyzed; n=324 treatment, n=332 control	HAZ, WHZ, WAZ, Stunting, Wasting, Underweight
Loría et al, 1979	Mexico	~2 mo (mean age)	1-2 mg/kg elemental Fe as ferrous sulfate daily for 16 months	Placebo; or Placebo + Prenatal iron (factorial trial)	n=92 total participants analyzed; n=48 treatment, n=44 control	Hemoglobin

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Lozoff et al, 2016	China	~2 mo (mean age)	1 mg/kg elemental Fe as iron proteinsuccinylate oral solution daily for 7.5 months	Placebo; or Placebo + Maternal iron (factorial trial)	n=1276 total participants analyzed; n=648 treatment, n=628 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia, HAZ, WHZ, WAZ
López de Romaña et al, 2005	Peru	6 mo - 12 mo	10 mg elemental Fe as ferrous fumarate 7 times per week for 6 months	Placebo	n=146 total participants analyzed; n=74 treatment, n=72 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, HAZ, WHZ, WAZ, Diarrhea, Respiratory infection
Majumdar et al, 2003	India	6 mo - 24 mo	2 mg/kg elemental iron (formulation not specified) daily for 4 months	Placebo	n=100 total participants analyzed; n=50 treatment, n=50 control	Hemoglobin, Ferritin
Malan et al, 2015	South Africa	6 yr - 12 yr	50 mg elemental Fe as ferrous sulfate 4 times per week for 8.5 months	Placebo + Vitamin C; or Placebo + n-3 fatty acids + Vitamin C (factorial trial)	n=296 total participants analyzed; n=146 treatment, n=150 control	Diarrhea, Respiratory infection
Massaga et al, 2003	Tanzania	3 mo - 4 mo	7.5 mg elemental Fe as ferric ammonium citrate daily for 6 months	Placebo; or Placebo + Amodaquine (antimalarial) (factorial trial)	n=291 total participants analyzed; n=146 treatment, n=145 control	Anemia, Malaria
Mebrahtu et al, 2004	Tanzania	4 mo - 6 yr	10 mg elemental Fe as ferrous sulfate + Placebo & Anthelmintic (Mebendazole) to half daily for 12 months	Placebo + Placebo & Anthelmintic (Mebendazole) to half	n=538 total participants analyzed; n=273 treatment, n=265 control	Malaria
Mejía et al, 1988	Guatemala	12 mo - 9 yr	3 mg/kg elemental Fe as ferrous sulfate daily for 2 months	Placebo; or Placebo + Vitamin A (factorial trial)	n=99 total participants analyzed; n=54 treatment, n=45 control	Hemoglobin

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Menendez et al, 1997	Tanzania	~2 mo (mean age)	12 mg elemental Fe as ferrous glycine sulphate daily for 16 weeks	Placebo; or Placebo + Malaria prophylaxis (Deltaprim) (factorial trial)	n=832 total participants analyzed; n=417 treatment, n=415 control	Malaria
Menendez et al, 2004	Tanzania	~2 mo (mean age)	12 mg elemental Fe as ferrous glycine sulphate daily for 4 months	Placebo; or Placebo + Malaria prophylaxis (Deltaprim) (factorial trial)	n=601 total participants analyzed; n=307 treatment, n=294 control	Ferritin, Iron deficiency
Metallinos-Katsaras et al, 2004	Greece	3 yr - 5 yr	15 mg elemental Fe as ferrous fumarate + Multivitamin (Vitamins A, B, C, D, E) 5 times per week for 2 months	Placebo + Multivitamin (Vitamins A, B, C, D, E)	n=48 total participants analyzed; n=31 treatment, n=17 control	Hemoglobin, Ferritin
Mitra et al, 1997	Bangladesh	2 mo - 4 yr	15 mg elemental Fe as ferrous gluconate + Multivitamin (A, C, D) daily for 15 months	Placebo + Multivitamin (A, C, D)	n=289 total participants analyzed; n=141 treatment, n=148 control	Diarrhea, Respiratory infection
Mozaffari-Koshravi et al, 2010	Iran	14 yr - 17 yr	150 mg elemental Fe as ferrous sulfate 1 time per week for 16 weeks	Control	n=193 total participants analyzed; n=96 treatment, n=97 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Mwanri et al, 2000	Tanzania	9 yr - 12 yr	40 mg elemental Fe as ferrous sulfate 3 times per week for 12 weeks	Placebo; or Placebo + Vitamin A (factorial trial)	n=136 total participants analyzed; n=68 treatment, n=68 control	Hemoglobin, Anemia
Nagpal et al, 2004	India	4 mo - 6 mo	2 mg/kg elemental Fe as ferric ammonium citrate daily for 8 weeks	Placebo	n=43 total participants analyzed; n=19 treatment, n=24 control	Hemoglobin, Ferritin
Nair et al, 2017	India	4 mo - 6 mo	2 mg/kg elemental Fe as colloidal iron daily for 7 months	Control	n=44 total participants analyzed; n=22 treatment, n=22 control	Hemoglobin, Anemia

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Nchito et al, 2009	Zambia	~10 yr (mean age)	60 mg elemental Fe as ferrous dextran 5 times per week for 10 months	Placebo; or Placebo + Multivitamin (factorial trial)	n=215 total participants analyzed; n=118 treatment, n=97 control	Helminth Infection
Northrop-Clewes et al, 1996	Pakistan	~14 mo (mean age)	15 mg elemental Fe as ferrous sulfate daily for 12 weeks	Placebo	n=191 total participants analyzed; n=95 treatment, n=96 control	Hemoglobin, Ferritin
Olsen et al, 2000	Kenya	4 yr - 16 yr	60 mg elemental Fe as ferrous dextran 2 times per week for 12 months	Placebo	n=200 total participants analyzed; n=108 treatment, n=92 control	Helminth Infection
Olsen et al, 2006	Kenya	~9 yr (mean age)	60 mg elemental Fe as ferrous dextran 2 times per week for 12 months	Placebo	n=200 total participants analyzed; n=108 treatment, n=92 control	Hemoglobin, Ferritin
Paganini et al, 2017	Kenya	6 mo - 9.5 mo	5 mg elemental Fe as NaFeEDTA and ferrous fumarate + Micronutrient powder daily for 4 months	Placebo + Micronutrient powder	n=97 total participants analyzed; n=49 treatment, n=48 control	Hemoglobin, Ferritin, Anemia, Iron deficiency anemia
Palupi et al, 1997	Indonesia	24 mo - 6 yr	30 mg elemental Fe as ferrous sulfate 1 time per week for 9 weeks	Placebo	n=194 total participants analyzed; n=96 treatment, n=98 control	Hemoglobin, Anemia, HAZ, WHZ, WAZ, Helminth Infection
Paracha et al, 1993	Pakistan	8 yr - 11 yr	76 mg elemental Fe as ferrous gluconate + Multivitamin daily for 11 weeks	Control + Multivitamin	n=119 total participants analyzed; n=61 treatment, n=58 control	Hemoglobin, Ferritin
Perrone et al, 1999	Italy	4 yr - 12 yr	12 mg elemental Fe as iron polymaltosate + Zinc daily for 12 months	Control + Zinc	n=20 total participants analyzed; n=10 treatment, n=10 control	Ferritin, HAZ, WAZ
Prasetyani et al, 2017	Indonesia	5 yr - 19 yr	100 mg elemental Fe as iron(iii)-hydroxide polymaltose complex 6 times per week for 8 weeks	Placebo	n=578 total participants analyzed; n=290 treatment, n=288 control	Hemoglobin, Ferritin, Malaria

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Reeves et al, 1985	USA	11 mo - 14 mo	3 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Placebo	n=179 total participants analyzed; n=77 treatment, n=102 control	Diarrhea
Rezaeian et al, 2014	Iran	14 yr - 19 yr	50 mg elemental Fe as ferrous sulfate 2 times per week for 16 weeks	Placebo	n=200 total participants analyzed; n=100 treatment, n=100 control	Hemoglobin
Richard et al, 2006	Peru	6 mo - 16 yr	15 mg elemental Fe as ferrous sulfate daily for 7 months	Placebo; or Placebo + Zinc (factorial trial)	n=758 total participants analyzed; n=378 treatment, n=380 control	Hemoglobin, HAZ, WAZ, Diarrhea, Respiratory infection, Malaria
Rosado et al, 1997	Mexico	18 mo - 3 yr	20 mg elemental Fe as ferrous sulfate 5 times per week for 12 months	Placebo; or Placebo + Zinc (factorial trial)	n=217 total participants analyzed; n=108 treatment, n=109 control	Hemoglobin, Ferritin, Iron deficiency, HAZ, WHZ, WAZ, Diarrhea, Respiratory infection
Rosado et al, 2006	Mexico	6 yr - 8 yr	30 mg elemental Fe as ferrous fumarate 5 times per week for 6 months	Placebo; or Placebo + Zinc (factorial trial)	n=517 total participants analyzed; n=265 treatment, n=252 control	Hemoglobin, Ferritin
Roschnik et al, 2004	Philippines	7 yr - 13 yr	108 mg elemental Fe as ferrous sulfate 1 time per week for 10 weeks	Control	n=1510 total participants analyzed; n=708 treatment, n=802 control	Hemoglobin, Anemia
Sarker et al, 2008	Bangladesh	24 mo - 6 yr	3 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Placebo; or Placebo + H-pylori treatment (factorial trial)	n=197 total participants analyzed; n=99 treatment, n=98 control	Hemoglobin, Ferritin
Seshadri et al, 1989	India	8 yr - 16 yr	30 mg elemental Fe as ferrous sulfate daily for 2 months; 40 mg elemental Fe as ferrous sulfate daily for 2 months	Placebo	n=113 total participants analyzed; n=97 treatment, n=16 control	Hemoglobin, Anemia
Smith et al, 1989	Gambia	6 mo - 6 yr	3-6 mg/kg elemental Fe as ferrous sulfate daily for 12 weeks	Placebo	n=186 total participants analyzed; n=97 treatment, n=89 control	Malaria

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Smuts et al, 2005	South Africa	6 mo - 12 mo	10 mg elemental iron (formulation not specified) daily for 6 months	Placebo	n=99 total participants analyzed; n=49 treatment, n=50 control	Hemoglobin, Ferritin, HAZ, WHZ, WAZ, Diarrhea, Respiratory infection
Smuts et al, 2014	South Africa	6 yr - 12 yr	50 mg elemental Fe as ferrous sulfate 4 times per week for 8.5 months	Placebo + Vitamin C; or Placebo + n-3 fatty acids + Vitamin C (factorial trial)	n=86 total participants analyzed; n=43 treatment, n=43 control	Cognitive
Soemantri et al, 1989	Indonesia	8 yr - 12 yr	2 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Placebo	n=130 total participants analyzed; n=71 treatment, n=59 control	Hemoglobin, Cognitive
Soewondo et al, 1989	Indonesia	~5 yr (mean age)	50 mg elemental Fe as ferrous sulfate daily for 8 weeks	Placebo	n=176 total participants analyzed; n=77 treatment, n=99 control	Hemoglobin, Ferritin
Stoltzfus et al, 2001	Tanzania (Zanzibar)	12 mo - 4 yr	10 mg elemental Fe as ferrous sulfate daily for 12 months	Placebo	n=538 total participants analyzed; n=277 treatment, n=261 control	Hemoglobin, Ferritin, Helminth Infection, Cognitive
Stoltzfus et al, 2004	Tanzania	6 mo - 29 mo	10 mg elemental Fe as ferrous sulfate daily for 12 months	Placebo; or Placebo + Mebendazole (factorial trial)	n=145 total participants analyzed; n=74 treatment, n=71 control	Anemia, Stunting
Sungthong et al, 2002	Thailand	6 yr - 14 yr	60 mg elemental Fe as ferrous sulfate daily for 16 weeks; 60 mg elemental Fe as ferrous sulfate 1 time per week for 16 weeks	Placebo	n=396 total participants analyzed; n=274 treatment, n=122 control	Hemoglobin, Ferritin, Iron deficiency anemia, HAZ, WAZ

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Sungthong et al, 2004	Thailand	6 yr - 14 yr	60 mg elemental Fe as ferrous sulfate 5 times per week for 16 weeks; 60 mg elemental Fe as ferrous sulfate 1 time per week for 16 weeks	Placebo	n=391 total participants analyzed; n=269 treatment, n=122 control	Cognitive
Teshome et al, 2017	Kenya	12 mo - 3 yr	3 mg elemental Fe as NaFeEDTA daily for 1 month; 12.5 mg elemental Fe as ferrous fumarate daily for 1 month	Placebo + Anti-malarial (dihydroartemisinin-piperaquine); anti-helminthic (albendazole, praziquantel); micronutrient powder (13 nutrients)	n=315 total participants analyzed; n=210 treatment, n=105 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Malaria
Thibault et al, 1993	France	6 mo - 4 yr	38.6 mg elemental Fe as hydroxyproline iron daily for 2 months	Placebo	n=70 total participants analyzed; n=32 treatment, n=38 control	Hemoglobin, Ferritin
Untoro et al, 2005	Indonesia	6 mo - 12 mo	10 mg elemental Fe as ferrous sulfate 7 times per week for 23 weeks	Placebo	n=134 total participants analyzed; n=69 treatment, n=65 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, HAZ, WHZ, WAZ, Diarrhea, Respiratory infection
van den Hombergh et al, 1996	Tanzania	0 mo - 30 mo	40 mg elemental Fe as ferrous sulfate + Malaria treatment + folic acid daily for 12 weeks	Placebo + Malaria treatment + folic acid	n=95 total participants analyzed; n=48 treatment, n=47 control	Hemoglobin, WAZ, Malaria
Verhoef et al, 2002	Kenya	2 mo - 3 yr	3 mg/kg elemental Fe as ferrous fumarate 2 times per week for 12 weeks	Placebo; or Placebo + Antimalarial (sulfadoxine-	n=328 total participants analyzed; n=164 treatment, n=164 control	Hemoglobin, Iron deficiency, Malaria

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
				pyrimethamine) (factorial trial)		
Wang et al, 2012	China	~4 mo (mean age)	1 mg/kg elemental Fe as for 2 months	Control	n=60 total participants analyzed; n=26 treatment, n=34 control	Hemoglobin, Ferritin
Wasantwisut et al, 2006	Thailand	4 mo - 6 mo	10 mg elemental Fe as ferrous sulfate daily for 6 months	Placebo + Vitamin C, A; or Placebo + Zinc + Vitamins C, A (factorial trial)	n=609 total participants analyzed; n=305 treatment, n=304 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia, HAZ, WHZ, WAZ
Wieringa et al, 2007	Indonesia, Thailand, Vietnam	4 mo - 6 mo	10 mg elemental Fe as ferrous sulfate 5-7 times per week for 6 months	Placebo; or Placebo + Zinc (factorial trial)	n=2049 total participants analyzed; n=1017 treatment, n=1032 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Yalçın et al, 2000	Turkey	~6 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Control	n=16 total participants analyzed; n=7 treatment, n=9 control	Hemoglobin, Ferritin, Anemia, Iron deficiency, Iron deficiency anemia
Yip et al, 1985	USA	11 mo - 13 mo	30 mg/kg elemental Fe as ferrous sulfate daily for 3 months	Placebo	n=291 total participants analyzed; n=146 treatment, n=145 control	Hemoglobin, Ferritin
Yurdakök et al, 2004	Turkey	~4 mo (mean age)	1 mg/kg elemental Fe as ferrous sulfate daily for 3 months; 7 mg/kg elemental Fe as ferrous sulfate 1 time per week for 3 months	Control	n=53 total participants analyzed; n=37 treatment, n=16 control	Hemoglobin, Ferritin
Zavaleta et al, 2000	Peru	12 yr - 19 yr	60 mg elemental Fe as ferrous sulfate 2 times per week for 17 weeks	Placebo	n=198 total participants analyzed; n=101 treatment, n=97 control	Anemia

Author & Year	Country	Age	Intervention	Comparison group	Sample size	Outcomes
Zhao et al, 2004	China	3 yr - 6 yr	4 mg elemental Fe as NaFeEDTA for 18 months	Placebo	n=213 total participants analyzed; n=120 treatment, n=93 control	Hemoglobin, Anemia
Ziegler et al, 2009	USA	4 mo - 4 mo	7.5 mg elemental Fe as ferrous sulfate daily for 5 months	Control	n=98 total participants analyzed; n=42 treatment, n=56 control	Hemoglobin, Ferritin
Ziegler et al, 2009	USA	~1 mo (mean age)	7 mg elemental Fe as ferrous sulfate + Multivitamins (A, C, D) daily for 4.5 months	Placebo + Multivitamins (A, C, D)	n=63 total participants analyzed; n=31 treatment, n=32 control	Hemoglobin, Ferritin, Iron deficiency, Iron deficiency anemia
Zlotkin et al, 2003	Ghana	8 mo - 20 mo	40 mg elemental Fe as ferrous fumarate daily for 6 months; 12.5 mg elemental Fe as ferrous sulfate daily for 6 months	Placebo	n=241 total participants analyzed; n=161 treatment, n=80 control	Hemoglobin, Ferritin, Anemia
Zlotkin et al, 2013	Ghana	6 mo - 4 yr	12.5 mg elemental Fe as microencapsulated ferrous fumarate + Ascorbic acid + vitamin a + zinc daily for 5 months	Placebo + Ascorbic acid + vitamin a + zinc	n=1815 total participants analyzed; n=900 treatment, n=915 control	Hemoglobin, Anemia, Iron deficiency, Malaria

Appendix 6. Modification of iron effects after control for baseline anemia.

	Hemoglobin (g/L) *	Ferritin (ng/mL) *	Anemia †	Iron deficiency †	Iron deficiency anemia †
Weekly frequency					
Frequent (3-7 times/week)	ref	ref	ref	ref	ref
n	95	60	48	29	15
Intermittent (1-2 times/week)					
n	23	16	9	4	3
Estimate	-3.1	-15.0	0.98	1.12	1.61
95% CI	(-5.9, -0.3)	(-24.7, -5.3)	(0.65, 1.48)	(0.24, 3.19)	(0.29, 8.84)
p-interaction	0.03	0.003	0.94	0.83	0.56
Duration					
1-3 months					
n	74	43	23	13	5
Estimate	ref	ref	ref	ref	ref
4-6 months					
n	39	30	30	20	13
Estimate	0.2	8.2	1.07	0.55	1.54
95% CI	(-2.4, 2.9)	(-1.0, 17.5)	(0.76, 1.52)	(0.25, 1.25)	(0.26, 9.28)
≥7 months					
n	8	4	6	0	0
Estimate	-1.3	40.7	1.38	n/e	n/e
95% CI	(-6.0, 3.5)	(1.7, 79.6)	(0.85, 2.24)		
Global p-interaction	0.83	0.04	0.40	0.15	0.61
Dose‡					
Low (1st tertile)					
n	36	26	21	9	8
Estimate	ref	ref	ref	ref	ref
Moderate (2nd tertile)					
n	27	21	22	15	9
Estimate	4.9	14.5	0.82	1.01	0.30
95% CI	(1.8, 8.0)	(5.9, 23.1)	(0.59, 1.13)	(0.40, 2.57)	(0.14, 0.69)
High (3rd tertile)					
n	31	20	8	6	1
Estimate	2.1	6.4	1.07	0.96	0.24
95% CI	(-1.0, 5.1)	(-2.4, 15.1)	(0.68, 1.70)	(0.24, 3.93)	(0.01, 6.80)
Global p-interaction	0.008	0.006	0.33	1.00	0.008

* Difference in the weighted mean effect of iron in the index category compared to the weighted mean effect of iron in the reference category, adjusting for baseline anemia.

† Ratio of the prevalence ratio for iron in the index category relative to the prevalence ratio for iron in the reference category, adjusting for baseline anemia.

‡ Age-adjusted dose tertiles; see Appendix 3.

Abbreviations: CI, confidence interval; ref, reference category; n/e, not estimable

Appendix 7. Effect modification of iron by baseline anemia, child age, child sex, WHO region, and iron formulation.

	Hemoglobin (g/L) *	Ferritin (ng/mL) *	Anemia †	Iron deficiency †	Iron deficiency anemia †
Baseline anemia					
All anemic					
n	33	17	9	5	1
Estimate	10.6	13.1	0.35	0.19	0.15
95% CI	(7.3, 13.9)	(8.0, 18.2)	(0.26, 0.47)	(0.06, 0.61)	(0.02, 1.26)
Mixed anemic and non-anemic					
n	54	30	40	23	13
Estimate	6.5	22.7	0.63	0.26	0.13
95% CI	(5.4, 7.5)	(17.4, 27.9)	(0.55, 0.71)	(0.19, 0.36)	(0.08, 0.22)
All non-anemic					
n	28	20	10	5	4
Estimate	5.1	18.7	0.62	0.20	0.21
95% CI	(3.5, 6.7)	(13.5, 23.9)	(0.41, 0.93)	(0.11, 0.36)	(0.06, 0.70)
Missing baseline anemia data					
n	52	40	10	15	9
Estimate	3.3	16.6	0.82	0.61	0.48
95% CI	(2.4, 4.2)	(13.4, 19.8)	(0.70, 0.95)	(0.48, 0.78)	(0.29, 0.78)
Global p-interaction	<0.001	0.28	0.002	0.13	0.06
Child age					
0 to 5 months					
n	39	33	18	19	17
Estimate	5.2	25.1	0.57	0.23	0.17
95% CI	(3.7, 6.7)	(19.7, 30.5)	(0.48, 0.70)	(0.15, 0.35)	(0.10, 0.31)
6 to 23 months					
n	31	23	23	15	5
Estimate	5.1	14.4	0.77	0.38	0.32
95% CI	(3.3, 6.9)	(8.7, 20.0)	(0.67, 0.90)	(0.27, 0.53)	(0.14, 0.69)
2 to <5 years					
n	25	10	6	4	0
Estimate	5.7	19.5	0.76	0.31	n/e
95% CI	(3.8, 7.7)	(11.3, 27.7)	(0.53, 1.09)	(0.14, 0.67)	
5 to <12 years					
n	29	13	13	6	4
Estimate	8.5	17.9	0.48	0.27	0.19
95% CI	(5.7, 11.4)	(10.9, 24.9)	(0.38, 0.60)	(0.11, 0.69)	(0.06, 0.56)
≥12 years					
n	21	20	5	3	1
Estimate	6.6	10.9	0.49	0.30	0.09
95% CI	(4.4, 8.7)	(9.1, 12.7)	(0.31, 0.79)	(0.11, 0.80)	(0.01, 1.64)
Global p-interaction	0.18	0.01	0.047	0.73	0.57
Child sex					

All female					
n	22	17	5	2	1
Estimate	6.6	9.3	0.39	0.74	0.09
95% CI	(4.2, 9.1)	(7.7, 10.9)	(0.28, 0.55)	(0.46, 1.19)	(0.01, 1.64)
All male					
n	6	0	2	0	0
Estimate	12.1	n/e	0.54	n/e	n/e
95% CI	(10.1, 14.2)		(0.46, 0.64)		
Mixed female and male					
n	100	70	58	38	25
Estimate	5.5	20.1	0.63	0.33	0.21
95% CI	(4.6, 6.4)	(16.8, 23.5)	(0.57, 0.70)	(0.26, 0.41)	(0.13, 0.32)
Missing baseline sex data					
n	39	20	4	8	1
Estimate	7.6	21.5	0.61	0.17	0.11
95% CI	(4.9, 10.3)	(12.3, 30.6)	(0.43, 0.86)	(0.09, 0.31)	(0.01, 1.99)
Global p-interaction	0.07	0.04	0.41	0.14	0.85
WHO region					
Africa					
n	26	11	20	10	2
Estimate	6.1	17.6	0.70	0.28	0.35
95% CI	(3.9, 8.3)	(13.1, 22.1)	(0.60, 0.82)	(0.16, 0.49)	(0.21, 0.56)
Americas					
n	29	13	13	7	3
Estimate	4.6	19.3	0.63	0.75	0.66
95% CI	(2.6, 6.7)	(13.1, 25.5)	(0.43, 0.91)	(0.51, 1.09)	(0.20, 2.14)
Eastern Mediterranean					
n	12	11	1	1	1
Estimate	8.6	8.9	0.14	0.79	0.09
95% CI	(6.1, 11.2)	(5.8, 12.0)	(0.03, 0.62)	(0.48, 1.29)	(0.01, 1.64)
Europe					
n	21	19	5	10	6
Estimate	4.4	17.4	0.32	0.28	0.16
95% CI	(2.8, 6.0)	(11.8, 23.1)	(0.14, 0.70)	(0.20, 0.41)	(0.06, 0.45)
South-East Asia					
n	56	34	20	12	10
Estimate	8.2	25.4	0.57	0.23	0.13
95% CI	(6.5, 9.7)	(19.4, 31.5)	(0.48, 0.67)	(0.16, 0.33)	(0.09, 0.18)
Western Pacific					
n	18	14	10	8	5
Estimate	5.4	14.1	0.58	0.38	0.53
95% CI	(3.6, 7.3)	(8.3, 19.9)	(0.44, 0.76)	(0.25, 0.57)	(0.29, 0.96)
Global p-interaction	0.06	0.06	0.20	0.44	0.007
Iron formulation					
Ferrous sulfate					
n	110	66	46	27	20
Estimate	7.2	21.8	0.56	0.21	0.12

95% CI	(6.1, 8.3)	(18.5, 25.1)	(0.49, 0.64)	(0.15, 0.30)	(0.09, 0.17)
Ferrous fumarate					
n	15	11	8	9	1
Estimate	4.8	16.7	0.69	0.29	0.40
95% CI	(2.2, 7.4)	(12.7, 20.8)	(0.53, 0.91)	(0.18, 0.47)	(0.08, 2.01)
Other or unspecified					
n	41	29	14	12	5
Estimate	4.2	10.3	0.78	0.56	0.81
95% CI	(2.8, 5.6)	(5.3, 15.3)	(0.62, 0.97)	(0.44, 0.73)	(0.58, 1.14)
Global p-interaction	0.02	0.005	0.13	0.03	<0.001

* Weighted mean difference

† Pooled risk ratio

Abbreviations: CI, confidence interval; n/e, not estimable

Appendix 8. Comparison of iron supplementation effects within factorial trials, using double placebo as a common referent group.

	Hemoglobin (g/L) *	Ferritin (ng/mL) *	Anemia †	Iron deficiency †	Iron deficiency anemia †
Zinc					
n	10	9	6	6	5
Double placebo	ref	ref	ref	ref	ref
Iron + Placebo	6.6 (3.6, 9.6)	28.8 (22.2, 35.4)	0.41 (0.33, 0.50)	0.15 (0.09, 0.24)	0.08 (0.05, 0.15)
Zinc + Placebo	0.1 (-1.5, 1.6)	-0.4 (-2.0, 1.2)	0.93 (0.78, 1.10)	1.06 (0.95, 1.17)	1.04 (0.81, 1.34)
Iron + Zinc	4.2 (1.9, 6.5)	21.0 (16.7, 25.3)	0.59 (0.46, 0.75)	0.19 (0.12, 0.31)	0.16 (0.09, 0.27)
Vitamin A					
n	6	3	2	1	0
Double placebo	ref	ref	ref	ref	n/a
Iron + Placebo	7.6 (3.0, 12.2)	8.4 (5.3, 11.4)	0.27 (0.16, 0.45)	0.34 (0.18, 0.66)	n/a
Vitamin A + Placebo	6.0 (3.2, 8.8)	-4.0 (-17.6, 9.5)	0.49 (0.36, 0.67)	1.24 (0.83, 1.84)	n/a
Iron + Vitamin A	10.5 (6.0, 14.9)	0.6 (-4.1, 5.3)	0.25 (0.07, 0.94)	0.63 (0.38, 1.07)	n/a

* Weighted mean difference

† Pooled risk ratio

Abbreviations: CI, confidence interval

Appendix 9. Effect heterogeneity p-values for anthropometric, infectious, and development outcomes.*†

	Modifiers							
	Frequency	Duration	Dose for age tertile	Baseline percent anemic	Age category	Baseline percent female	Factorial trial	Formulation
<i>Anthropometry</i>								
Height-for-age Z score	0.255	0.598	0.835	0.013	0.256	n/e	0.048	0.272
Weight-for-height Z score	0.970	0.665	0.291	n/e	0.311	n/e	0.559	0.843
Weight-for-age Z score	0.524	0.508	0.861	0.709	0.250	n/e	0.363	0.847
Stunting	n/e	0.452	0.393	n/e	0.378	0.378	0.700	n/e
Wasting	n/e	n/e	0.522	n/e	n/e	n/e	0.336	n/e
<i>Infections</i>								
Diarrhea (cumulative incidence)	0.136	0.282	0.274	0.358	0.884	n/e	0.642	0.904
Diarrhea (incidence rate)	0.962	0.471	0.435	n/e	0.446	n/e	0.760	0.760
Respiratory illness (cumulative incidence)	0.006	0.018	0.032	0.125	n/e	n/e	0.129	0.683
Respiratory illness (incidence rate)	0.902	0.866	0.750	n/e	0.887	n/e	0.421	0.421
Malaria (prevalence)	n/e	n/e	0.996	0.910	n/e	n/e	0.225	0.622
Malaria (incidence rate)	n/e	0.469	0.965	0.871	n/e	n/e	0.463	0.467
Hookworm (prevalence)	0.968	0.812	0.880	n/e	n/e	n/e	n/e	0.823
Ascaris lumbricoides (prevalence)	0.390	0.456	0.647	n/e	n/e	n/e	n/e	0.932
Trichuris trichiura (prevalence)	0.501	0.501	0.431	n/e	n/e	n/e	n/e	0.584
<i>Development</i>								
Bayley Mental Index	0.651	0.892	0.888	0.020	n/e	n/e	0.280	0.454
Bayley Psychomotor Index	0.807	0.432	0.738	0.023	n/e	n/e	0.672	0.634

* Categories used for modifiers: Frequency (1-2 times per week vs. 3-7 times per week); Dose for age tertile (1st vs. 2nd vs. 3rd); Baseline percent anemic (<20% anemic vs. 20-39% anemic vs. ≥40% anemic); Age category (0-5 mo vs. 6-23 mo vs. 24-59 mo vs. 5-11 y vs. 12-19 y); Baseline percent female (All participants female vs. no participants female vs. some participants female); Factorial trial (Yes vs. no); Formulation (Ferrous sulfate vs. ferrous fumarate vs. other)

† P-values calculated from meta-regression Wald tests.


Abbreviations: n/e, not estimable.

Appendix 10. Risk of bias within included studies.

Study	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of partici-pants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)
Adish 1997	Unclear	Unclear	Low	Low	Low
Aggarwal 2005	Low	Unclear	Low	Low	High
Aguayo 2000	Low	Unclear	Low	Low	Low
Akman 2004	Low	Unclear	High	Low	Low
Angeles 1993	Unclear	Unclear	Unclear	Unclear	Low
Angulo-Barro 2016	Low	Low	Low	Low	Unclear
Arcanjo 2011	High	Unclear	Low	Low	Unclear
Arcanjo 2013	High	Unclear	Low	Low	Unclear
Aukett 1986	Unclear	Low	Low	Low	Low
Ayoya 2009	Unclear	Unclear	High	Low	High
Ayoya 2012	Unclear	Unclear	High	Low	High
Ballin 1992	High	High	Low	Low	Unclear
Baqui 2003	Unclear	Unclear	Unclear	Unclear	Low
Barclay 1991	Unclear	Unclear	High	High	Unclear
Baumgartner 2012	Low	Unclear	Low	Low	High
Berger 1997	Unclear	Unclear	Low	Unclear	Low
Berger 2000	Unclear	Unclear	Low	Low	Unclear
Berger 2006	Low	Low	Low	Low	High
Berglund 2010	Low	Unclear	Low	Low	High
Bhatia 1993	High	Unclear	Unclear	Unclear	Unclear
Black 2004	Unclear	Unclear	Low	Low	Low
Bora 2019	Low	Low	Unclear	Unclear	Low
Bruner 1996	Low	Unclear	Low	Low	High
Burman 1972	High	Unclear	High	High	High
Buzina-Suboticanec 1998	High	Unclear	Low	Low	Unclear
Charoenlarp 1980	Unclear	Unclear	Unclear	Unclear	Unclear
Chen 2011	Unclear	High	Unclear	Low	Unclear
Chen 2013	Low	Unclear	Low	Low	Low
Chen 2014	Low	Unclear	High	High	Low
Cheng 2001	Unclear	Unclear	Unclear	Unclear	Unclear
Choe 1999	Low	Unclear	Low	Low	Unclear
Chwang 1988	Unclear	Unclear	Low	Low	Low
Das 1984	High	Unclear	Unclear	Unclear	Low
de Silva 2003	Unclear	Unclear	Low	Low	Unclear
Desai 2003	Low	Low	Low	Low	High
Devaki 2008	Unclear	Unclear	Unclear	Unclear	Low
Dewey 2002	Unclear	Unclear	Low	Low	Low
Dijkhuizen 2001	Low	Low	Low	Low	Low
Dijkhuizen 2008	Low	Low	Low	Low	Unclear
Domellöf 2001	Unclear	Unclear	Low	Low	Unclear
Dossa 2001	Low	Unclear	Low	Low	Low
Dossa 2001	Unclear	Unclear	Low	Low	Unclear
Eftekhari 2006	Unclear	Unclear	Low	Low	High
Elwood 1970	Unclear	Unclear	Unclear	Unclear	Unclear
Engstrom 2008	Unclear	Unclear	High	High	Unclear
Ermis 2002	Unclear	Unclear	Unclear	Unclear	Unclear
Fahmida 2007	Unclear	Unclear	Low	Low	Low

Fallahi 2007	Unclear	Unclear	Low	Unclear	Low
Franz 2000	Unclear	Unclear	Unclear	Unclear	High
Friel 2003	Unclear	Unclear	Low	Low	High
Fujiu 2004	Unclear	Unclear	Unclear	Unclear	Low
Gebresellasie 1996	Low	Low	Low	Low	Low
Geltman 2001	Unclear	Low	Low	Low	Unclear
Geltman 2004	Low	Unclear	Low	Low	Unclear
Gokcay 2012	Unclear	Unclear	High	Unclear	High
Gopaldas 1985	Unclear	Unclear	Low	Low	Low
Gopaldas 1985	Unclear	Unclear	Low	Low	Low
Greisen 1986	Unclear	Unclear	Unclear	Unclear	Unclear
Hacıhamdioglu 2013	Unclear	Unclear	High	Unclear	Unclear
Harvey 1989	Unclear	Unclear	Low	Low	Low
Hathirat 1992	Unclear	Unclear	Low	Unclear	Unclear
Hess 2002	Unclear	Unclear	Low	Low	Low
Hettiarachchi 2008	Unclear	Unclear	Low	Low	Low
Hieu 2012	Low	Low	Low	Low	Unclear
Hop 2005	Low	Low	Low	Low	Unclear
Htet 2019	Low	Low	Low	Low	Low
Idjradinata 1993	Low	Unclear	Low	Low	Low
Idjratinata 1994	Low	Unclear	Unclear	Unclear	Unclear
Irigoyen 1991	Unclear	Unclear	Low	Low	High
Jalambo 2018	Unclear	Unclear	High	Low	Low
Kapur 2003	Low	Unclear	Low	Unclear	Unclear
Kashyap 1987	Unclear	Unclear	Low	Unclear	Unclear
Kashyap 1987	Unclear	Unclear	Low	Unclear	Unclear
Kianfar 2000	Unclear	Unclear	High	Unclear	Unclear
Kordas 2005	Low	Low	Low	Low	High
Kusumastuti 2018	Low	Low	Low	Low	Unclear
Lambert 2002	Unclear	Unclear	Low	Low	Unclear
Latham 1990	Unclear	Unclear	Low	Unclear	Low
Lawless 1994	Low	Unclear	Low	Low	Low
Lee 1997	Unclear	Unclear	Low	Unclear	Unclear
Leenstra 2009	Low	Unclear	Low	Low	Low
Lind 2003	Unclear	Low	Low	Low	Low
Lind 2004	Low	Low	Low	Low	Low
López de Romaña 2005	Low	Low	Low	Low	Low
Loría 1979	Unclear	Unclear	Unclear	Unclear	Unclear
Lozoff 2016	Low	Unclear	Low	Low	Low
Majumdar 2003	Unclear	Unclear	Low	Unclear	High
Malan 2015	Low	Unclear	Low	Low	High
Massaga 2003	Low	Low	Low	Low	Low
Mebrahtu 2004	Unclear	Low	Low	Low	Unclear
Mejía 1988	Low	Low	High	Unclear	Unclear
Menendez 1997	Low	Low	Low	Low	High
Menendez 2004	Low	Low	Low	Low	High
Metallinos-Katsaras 2004	Unclear	Unclear	Low	Low	Unclear
Mitra 1997	Unclear	Unclear	Low	Low	Low
Mozaffari-Koshravi 2010	High	Unclear	High	Unclear	Low
Mwanri 2000	Low	Unclear	Low	Low	Low
Nagpal 2004	Low	Unclear	Low	Low	High
Nair 2017	High	High	Unclear	Unclear	Low
Nchito 2009	Unclear	Unclear	Low	Low	High

Northrop-Clewes 1996	Unclear	Unclear	Low	Low	Unclear
Olsen 2000	Unclear	Unclear	Low	Low	Low
Olsen 2006	Low	Low	Unclear	Unclear	Unclear
Paganini 2017	Low	Low	Low	Low	Low
Palupi 1997	Unclear	Unclear	Low	Low	Unclear
Paracha 1993	Unclear	Unclear	Low	Unclear	Unclear
Perrone 1999	Unclear	Unclear	High	Unclear	Unclear
Prasetyani 2017	Low	Unclear	Low	Low	Low
Reeves 1985	High	Unclear	Unclear	Unclear	High
Rezaeian 2014	High	Unclear	Low	Low	Low
Richard 2006	Unclear	Unclear	Low	Low	Low
Rosado 1997	Unclear	Unclear	Low	Low	Low
Rosado 2006	Low	Unclear	Low	Low	Unclear
Roschnik 2004	Low	Unclear	High	Unclear	Unclear
Sarker 2008	Low	Low	Low	Low	Unclear
Seshadri 1989	Unclear	Unclear	Low	Low	Low
Smith 1989	Unclear	Unclear	Low	Low	Unclear
Smuts 2005	Unclear	Low	Low	Low	High
Smuts 2014	Low	Unclear	Low	Low	Unclear
Soemantri 1989	Unclear	Unclear	Low	Unclear	Unclear
Soewondo 1989	Unclear	Unclear	Low	Low	Unclear
Stoltzfus 2001	Low	Low	Low	Low	Unclear
Stoltzfus 2004	Unclear	Unclear	Low	Low	High
Sungthong 2002	Low	Low	Low	Low	Low
Sungthong 2004	Low	Low	Low	Low	Low
Teshome 2017	Low	Low	Low	Low	Unclear
Thibault 1993	Unclear	Unclear	Low	Low	Unclear
Untoro 2005	Unclear	Unclear	Low	Low	Low
van den Hombergh 1996	Unclear	Unclear	Unclear	High	Low
Verhoef 2002	Low	Low	Low	Low	Low
Wang 2012	Unclear	Low	Low	Unclear	High
Wasantwisut 2006	Low	Low	Low	Low	Low
Wieringa 2007	Low	Low	Low	Low	High
Yalçin 2000	Unclear	Unclear	High	High	High
Yip 1985	High	Unclear	Unclear	Unclear	Unclear
Yurdakök 2004	Unclear	Unclear	High	High	Low
Zavaleta 2000	Unclear	Unclear	Low	Low	Low
Zhao 2004	Unclear	Unclear	Low	Low	High
Ziegler 2009	Unclear	Unclear	High	High	Unclear
Ziegler 2009	Unclear	Unclear	Low	Low	Unclear
Zlotkin 2003	Low	Low	High	Low	Low
Zlotkin 2013	Low	Unclear	Low	Low	Low

Appendix 11. Cochrane risk of bias assessment of included studies (n=129).

Appendix 12. Effect of oral iron supplementation versus placebo among children and adolescents aged <20 years among trials judged to not be at “high” risk of bias.

	n*	Estimate type	Estimate of effect (95% CI)	p-value	I ² (%)
Hemoglobin (g/L)	105	WMD	6.5 (5.4, 7.6)	<0.001	95.5
Serum ferritin (ng/mL)	68	WMD	20.9 (17.4, 24.3)	<0.001	99.5
Anemia	42	RR	0.58 (0.51, 0.65)	<0.001	87.1
Iron deficiency	31	RR	0.31 (0.24, 0.41)	<0.001	91.1
Iron deficiency anemia	17	RR	0.27 (0.17, 0.44)	<0.001	79.9

* Number of trial arms randomized to iron

Abbreviations: CI, confidence interval; IRR, incidence rate ratio; PR, prevalence ratio; RR, risk ratio; SMD, standardized mean difference; WMD, weighted mean difference

Appendix 13. Assessment of small study bias for outcomes reported by ≥ 10 groups randomized to iron.