

SUPPLEMENTARY DIGITAL MATERIAL 2

Supplementary Table II.—Group scores, Between and within group differences for Kinesiophobia and Catastrophizing#.

<b>Secondary Outcomes</b>	<b>Baseline</b>	<b>End of Treatment 4wk</b>	<b>Follow-up 2mo</b>	<b>4wk - Baseline</b>	<b>2mo - Baseline</b>	<b>2mo - 4wk</b>
<b>Kinesiophobia</b>						
<b>TSK(17-68)<sup>b</sup></b>						
Group 1 (n=30)	42.2 (39.68, 44.72)	26.5 (24.75, 28.19)	21.3 (19.50, 23.10)	-15.7 (-18.59, 12.89)*	-20.9 (-24.42, -17.37)*	-5.2 (-6.47, 3.86)*
Group 2 (n=30)	39.2 (36.82, 41.58)	26.8 (24.72, 28.88)	22 ( 20.45, 23.62)	-12.4 (-14.66, 10.14)*	-17.2 (-20.05, 14.29)*	-4.7 (-6.52, 3.01)*
Between Groups dif.	3 (-0.39, 6.39)	-0.33 (-2.98, 2.31)	0.7 (-3.08, 1.62)			
<b>Catastrophizing</b>						
<b>PCS(0-52)<sup>b</sup></b>						
Group 1 (n=30)	16.8 (14.35, 19.25)	3.6 (2.38, 4.88)	1.5 (0.59, 2.41)	-13.2 (-15.78, 10.55)*	-15.3 (-18.11, -12.48)*	-2.1 (-2.79, 1.48)*
Group 2 (n=30)	14.5 (12.29, 16.70)	4.1 (2.81, 5.39)	1.5 (0.68, 2.25)	-10.4 (-12.10, -7.80)*	-13.0 (-15.72, -10.35)*	-2.6 (-3.74, 1.53)*
Between Groups dif.	2.3 (-0.92, 5.52)	-4.7 (-2.22, 1.29)	0.03 (-1.14, 1.21)			
<b>Rumination PCS (0-16)<sup>b</sup></b>						
Group 1 (n=30)	6.5 (5.70, 7.30)	1.3 (0.73, 1.80)	0.4 (0.08, 0.65)	-5.2 (-6.29, -4.18)*	-6.1 (-7.16, -5.11)*	-0.9 (-1.44, 0.36)*
Group 2 (n=30)	5.0 (4.29, 5.71)	1.4 (0.78, 2.09)	0.4 (0.06, 0.67)	-3.5 (-4.48, -2.66)*	-4.6 (-5.49, -3.78)*	-1.0 (-1.71, -)

Between Groups dif.	1.5 (0.45, 2.55)*	-0.7 (-0.10, 0.66)	0.0 (-0.41, 0.41)			0.42)*
<b>Magnification PCS (0-12)<sup>b</sup></b>						
Group 1 (n=30)	3.8 (3.32, 4.28)	1.2 (0.83, 1.51)	0.5 (0.21, 0.79)	-2.6 (-3.14, -2.12)*	-3.3 (-3.86, -2.74)*	-0.7 (-0.97, -0.36)*
Group 2 (n=30)	3.7 (2.94, 4.52)	1.2 (0.91, 1.62)	0.7 (0.42, 1.04)	-2.5 (-3.46, -1.47)*	-3.0 (-3.95, -2.05)*	-0.5 (-0.93, -0.13)*
Between Groups dif.	0.7(-0.84, 0.97)	-0.1 (-0.58, 0.38)	-0.2 (-0.65, 0.18)			
<b>Helplessness PCS (0-24)<sup>b</sup></b>						
Group 1 (n=30)	6.5 (5.06, 7.94)	1.3 (0.58, 2.08)	0.6 (0.01, 1.33)	-5.1 (-6.56, -3.77)*	-5.8 (-7.44, -4.22)*	-0.6 (-0.99, -0.34)*
Group 2 (n=30)	5.6 (4.60, 6.93)	1.5 (1.01, 2.06)	0.3 (0.06, 0.67)	-4.2 (-5.53, -2.94)*	-5.4 (-6.81, -3.99)*	-1.1 (-1.76, -0.57)*
Between Groups dif.	0.7 (-1.08, 2.54)	-0.2 (-1.10, 0.70)	0.3 (-0.41, 1.01)			

TSK:Tampa Scale Kinesiophobia, PCS:Pain Catastrophizing Scale

b. Low score indicates better value, # Values are presented as mean (95%CI), \* P<0.05.