

Supplemental Table 4. Serum potassium distributions by category and study interval in the full study population

	Study Cohort (n = 270)			
Baseline	Mean (95% CI)		Median (IQR)	p-value
K+ value ^a	5.45 (5.38-5.53)		5.5 (5.1-5.9)	
K+ categories	n	%		
K+ < 3.5	0	0		
K+ 3.5 - 5.0	65	24.1		
K+ 5.1 - 5.4	64	23.7		
K+ 5.5 -5.9	91	33.7		
K+ >= 6.0	50	18.5		
K+ Missing	0	0		
Follow up 1-30 days.				
K+ value ^a	4.94 (4.85-5.02)		4.9 (4.6-5.4)	
K+ change from baseline ^b	-0.61 (0.49-0.73)			<0.01
K+ categories	n	%		
K+ < 3.5	1	0.4		
K+ 3.5 - 5.0	102	37.8		
K+ 5.1 - 5.4	44	16.3		
K+ 5.5 -5.9	31	11.5		
K+ >= 6.0	3	1.1		
K+ Missing	89	33.0		
Follow up 31-91 days.				
K+ value	4.88 (4.8-4.97)		4.9 (4.6-5.3)	
K+ change from baseline	-0.57 (0.45-0.68)			<0.01
K+ categories	n	%		
K+ < 3.5	1	0.4		
K+ 3.5 - 5.0	123	45.6		
K+ 5.1 - 5.4	45	16.7		
K+ 5.5 -5.9	22	8.1		
K+ >= 6.0	9	3.3		
K+ Missing	70	25.9		

Follow up 92-182 days.			
K+ value	4.89 (4.8-4.97)	4.9 (4.5-5.3)	
K+ change from baseline	-0.59 (0.47-0.71)		<0.01
K+ categories	n	%	
K+ < 3.5	2	0.7	
K+ 3.5 - 5.0	136	50.4	
K+ 5.1 - 5.4	41	15.2	
K+ 5.5 -5.9	23	8.5	
K+ >= 6.0	12	4.4	
K+ Missing	56	20.7	
<p>^a Average of K+ values that were closest to the end of the baseline period/index date or closest to the end of the follow up interval window</p> <p>^b Change of K+ values closest to the end of the follow up interval and K+ values closest to the end of baseline period/index date.</p>			