

Supplemental Table 6. Serum potassium distributions by category and study interval in patients with baseline serum potassium ≥ 5.5 mEq/L

Baseline	Study Cohort (n = 141)		
	Mean (95% CI)	Median (IQR)	p-value
K+ value ^a	5.94 (5.87-6.01)	5.8 (5.6-6.2)	
K+ categories	n	%	
K+ 5.5 -5.9	91	64.5	
K+ ≥ 6.0	50	35.5	
K+ Missing	0	0	
Follow up 1-30 days.			
K+ value ^a	4.98 (4.86-5.09)	4.9 (4.55-5.4)	
K+ change from baseline ^b	-1.0 (0.87-1.14)		<0.01
K+ categories	n	%	
K+ < 3.5	0	0	
K+ 3.5 - 5.0	57	40.4	
K+ 5.1 - 5.4	24	17.0	
K+ 5.5 -5.9	21	14.9	
K+ ≥ 6.0	2	1.4	
K+ Missing	37	26.2	
Follow up 31-91 days.			
K+ value	4.95 (4.85-5.06)	4.95 (4.6-5.3)	
K+ change from baseline	-1.0 (0.89-1.12)		<0.01
K+ categories	n	%	
K+ < 3.5	0	0	
K+ 3.5 - 5.0	65	46.1	
K+ 5.1 - 5.4	23	16.3	
K+ 5.5 -5.9	12	8.5	
K+ ≥ 6.0	6	4.3	
K+ Missing	35	24.8	
Follow up 92-182 days.			
K+ value	4.89(4.77-5.01)	4.9 (4.5-5.3)	
K+ change from baseline	-1.07 (0.93-1.21)		<0.01

K+ categories	n	%		
K+ < 3.5	1	0.7		
K+ 3.5 - 5.0	72	51.1		
K+ 5.1 - 5.4	20	14.2		
K+ 5.5 -5.9	13	9.2		
K+ >= 6.0	7	5.0		
K+ Missing	28	19.9		

^a Average of K+ values that were closest to the end of the baseline period/index date or closest to the end of the follow up interval window

^b Change of K+ values closest to the end of the follow up interval and K+ values closest to the end of baseline period/index date.